

Draft Comparative Effectiveness Review

Number xx

Pharmacotherapy for Adults With Alcohol-Use Disorders in Outpatient Settings

Prepared for:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
540 Gaither Road
Rockville, MD 20850
www.ahrq.gov

Contract No. xxx-xx-xxxx

Prepared by:

Investigators:

AHRQ Publication No. xx-EHCxxx
<Month Year>

This report is based on research conducted by the XXXXXXXXXX Evidence-based Practice Center (EPC) under contract to the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. XXX-20XX-XXXXX). The findings and conclusions in this document are those of the authors, who are responsible for its contents; the findings and conclusions do not necessarily represent the views of AHRQ. Therefore, no statement in this report should be construed as an official position of AHRQ or of the U.S. Department of Health and Human Services.

The information in this report is intended to help health care decisionmakers—patients and clinicians, health system leaders, and policymakers, among others—make well informed decisions and thereby improve the quality of health care services. This report is not intended to be a substitute for the application of clinical judgment. Anyone who makes decisions concerning the provision of clinical care should consider this report in the same way as any medical reference and in conjunction with all other pertinent information, i.e., in the context of available resources and circumstances presented by individual patients.

This report may be used, in whole or in part, as the basis for development of clinical practice guidelines and other quality enhancement tools, or as a basis for reimbursement and coverage policies. AHRQ or U.S. Department of Health and Human Services endorsement of such derivative products may not be stated or implied.

This report may periodically be assessed for the urgency to update. If an assessment is done, the resulting surveillance report describing the methodology and findings will be found on the Effective Health Care Program website at: www.effectivehealthcare.ahrq.gov. Search on the title of the report.

This document is in the public domain and may be used and reprinted without permission except those copyrighted materials that are clearly noted in the document. Further reproduction of those copyrighted materials is prohibited without the specific permission of copyright holders.

Persons using assistive technology may not be able to fully access information in this report. For assistance contact [insert program email address].

None of the investigators has any affiliations or financial involvement that conflicts with the material presented in this report.

Suggested citation: XXXXXX. Pharmacotherapy for Adults With Alcohol-Use Disorders in Outpatient Settings. Evidence Report No. XX. (Prepared by the XXXXXXXX Evidence-based Practice Center under Contract No. XXXXXXXX.) AHRQ Publication No. XXXXXX. Rockville, MD: Agency for Healthcare Research and Quality. Month Year.
www.effectivehealthcare.ahrq.gov.

Preface

The Agency for Healthcare Research and Quality (AHRQ), through its Evidence-based Practice Centers (EPCs), sponsors the development of systematic reviews to assist public- and private-sector organizations in their efforts to improve the quality of health care in the United States. These reviews provide comprehensive, science-based information on common, costly medical conditions, and new health care technologies and strategies.

Systematic reviews are the building blocks underlying evidence-based practice; they focus attention on the strength and limits of evidence from research studies about the effectiveness and safety of a clinical intervention. In the context of developing recommendations for practice, systematic reviews can help clarify whether assertions about the value of the intervention are based on strong evidence from clinical studies. For more information about AHRQ EPC systematic reviews, see www.effectivehealthcare.ahrq.gov/reference/purpose.cfm

AHRQ expects that these systematic reviews will be helpful to health plans, providers, purchasers, government programs, and the health care system as a whole. Transparency and stakeholder input are essential to the Effective Health Care Program. Please visit the Web site (www.effectivehealthcare.ahrq.gov) to see draft research questions and reports or to join an e-mail list to learn about new program products and opportunities for input.

We welcome comments on this systematic review. They may be sent by mail to the Task Order Officer named below at: Agency for Healthcare Research and Quality, 540 Gaither Road, Rockville, MD 20850, or by email to epc@ahrq.hhs.gov.

Richard Kronick, Ph.D.
Director
Agency for Healthcare Research and Quality

Stephanie Chang, M.D., M.P.H.
Director
Evidence-based Practice Program
Center for Outcomes and Evidence
Agency for Healthcare Research and Quality

Jean Slutsky, P.A., M.S.P.H.
Director, Center for Outcomes and Evidence
Agency for Healthcare Research and Quality

Carmen Kelly, Pharm.D., R.Ph.
Task Order Officer
Center for Outcomes and Evidence
Agency for Healthcare Research and Quality

Aysegul Gozu, M.D., M.P.H.
Task Order Officer
Center for Outcomes and Evidence
Agency for Healthcare Research and Quality

Acknowledgments

The authors gratefully acknowledge the continuing support of our AHRQ Task Order Officers, Carmen Kelly, Pharm.D., R.Ph. and Aysegul Gozu, M.D., M.P.H. We extend our appreciation to our Key Informants and members of our Technical Expert Panel, all of whom provided thoughtful advice and input during our research process.

Pharmacotherapy for Adults With Alcohol-Use Disorders in Outpatient Settings

Structured Abstract

Objectives: To conduct a systematic review and meta-analysis of the efficacy, comparative effectiveness, and harms of medications (both FDA-approved and others) for adults with alcohol-use disorders, and to evaluate the evidence from primary care settings.

Data sources: PubMed®, Cochrane Library, PsycINFO®, CINAHL®, EMBASE®, U.S. Food and Drug Administration Web site, ClinicalTrials.gov, and World Health Organization International Clinical Trials Registry Platform (January 1, 1970, to February 5, 2013).

Review methods: Two investigators independently selected, extracted data from, and rated risk of bias of studies. We conducted meta-analyses using random-effects models. We graded strength of evidence (SOE) based on established guidance.

Results: We included 130 studies. Most patients met criteria for alcohol dependence; mean ages were in the 40s. For acamprosate and naltrexone, numbers needed to treat (NNT) to prevent one person from returning to any drinking were 10 and 25, respectively (moderate SOE); NNT to prevent one person from returning to heavy drinking was 13 for naltrexone (moderate SOE). Our meta-analyses of 3 head-to-head trials found no statistically significant difference between the two medications for consumption outcomes (moderate SOE). With the exception of topiramate, current evidence does not establish the efficacy of medications used off-label or under investigation. No RCTs assessing acamprosate, naltrexone, or topiramate were conducted in primary care settings. We found insufficient direct evidence to conclude whether medications for alcohol dependence are effective for improving health outcomes.

Compared with placebo, patients treated with acamprosate had a higher risk of anxiety, diarrhea, and vomiting; those treated with naltrexone had a higher risk of dizziness, nausea, and vomiting. In head-to-head studies, the risk of headache was higher for naltrexone than for acamprosate. Trials of topiramate reported a significantly increased risk of paresthesias, anorexia, difficulty concentrating, dizziness, psychomotor slowing, and other adverse effects.

Evidence was insufficient to determine comparative effectiveness of medications for subgroups. Our meta-analyses for variation in naltrexone response related to *OPRM1* polymorphisms found no significant difference between AA homozygotes and those with at least one G allele.

Conclusions: Acamprosate and naltrexone have the best evidence of efficacy for improving alcohol consumption outcomes for patients with alcohol dependence. Evidence supports the efficacy of topiramate for improving some alcohol consumption outcomes, but adverse effects may limit its use clinically. Head-to-head trials have not consistently established superiority of one medication. Thus, other factors may contribute to medication choices, such as frequency of administration, potential adverse events, coexisting symptoms, and availability of treatments.

Contents

Executive Summary	ES-1
Introduction.....	1
Background.....	1
Treatments for Alcohol-Use Disorders	2
Existing Guidance.....	4
Scope and Key Questions	4
Analytic Framework	5
Methods.....	7
Topic Refinement and Review Protocol.....	7
Literature Search Strategy.....	7
Search Strategy	7
Inclusion and Exclusion Criteria.....	8
Study Selection	9
Data Extraction	10
Risk-of-Bias Assessment of Individual Studies.....	10
Data Synthesis.....	10
Strength of the Body of Evidence.....	11
Applicability	12
Peer Review and Public Commentary	12
Results.....	13
Results of Literature Searches	13
Key Question 1. Efficacy and Comparative Effectiveness for Improving Consumption	
Outcomes	14
Detailed Synthesis: Placebo-Controlled Trials of FDA-Approved Medications for Treating	
Alcohol Dependence	14
Detailed Synthesis: Placebo-Controlled Trials of Medications Used Off-label, or Those	
Under Investigation.....	33
Detailed Synthesis: Head-to-Head Trials.....	47
Key Question 2. Health Outcomes.....	53
Detailed Synthesis: Placebo-Controlled Trials of FDA-Approved Medications for Treating	
Alcohol Dependence	54
Key Question 3. Adverse Effects of Medications.....	69
Key Points	69
Detailed Synthesis.....	70
Characteristics of Included Studies	70
Key Question 4. Evidence from Primary Care Settings	75
Characteristics of Included Trials	75
Key Question 5. Subgroups	76
Detailed Synthesis.....	76
Key Question 6. Genetic Polymorphisms.....	81
Characteristics of Included Studies.....	81
Overview of Results.....	82
Detailed Results of Individual Studies.....	83
Discussion.....	85

Key Findings and Strength of Evidence	85
Efficacy and Comparative Effectiveness	85
Harms	88
Primary Care Settings	89
Subgroups and Genetic Polymorphisms	90
Findings in Relation to What Is Already Known.....	90
Applicability.....	91
Implications for Clinical and Policy Decisionmaking.....	91
Limitations of the Comparative Effectiveness Review Process	92
Limitations of the Evidence Base	92
Research Gaps.....	93
Conclusions.....	94
References.....	96

Tables

Table A. Definitions of the spectrum of alcohol misuse	ES-2
Table B. Medications that are FDA-approved for treating adults with alcohol dependence....	ES-3
Table C. Summary of findings and strength of evidence for efficacy of FDA-approved medications for alcohol dependence	ES-9
Table D. Summary of findings and strength of evidence for comparative effectiveness of acamprosate and naltrexone	ES-10
Table E. Evidence gaps for future research, by Key Question	ES-16

Table 1. Definitions of the spectrum of alcohol misuse	1
Table 2. Medications that are FDA approved for treating adults with alcohol-use disorders	3
Table 3. Eligibility criteria	8
Table 4. Definitions of the grades of overall strength of evidence	11
Table 5. Characteristics of included double-blind randomized placebo-controlled trials of acamprosate.....	15
Table 6. Characteristics of included double-blind randomized placebo-controlled trials of disulfiram	21
Table 7. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone.....	23
Table 8. Characteristics of included double-blind randomized placebo-controlled trials of baclofen.....	33
Table 9. Characteristics of included double-blind randomized placebo-controlled trials of buspirone.....	36
Table 10. Characteristics of included double-blind randomized placebo-controlled trials of citalopram	37
Table 11. Characteristics of included double-blind randomized placebo-controlled trials of fluoxetine	38
Table 12. Characteristics of included double-blind randomized placebo-controlled trials of nalmefene.....	41
Table 13. Characteristics of included double-blind randomized placebo-controlled trials of quetiapine	42

Table 14. Characteristics of included double-blind randomized placebo-controlled trials of sertraline	43
Table 15. Characteristics of included double-blind randomized placebo-controlled trials of topiramate	45
Table 16. Characteristics of included double-blind randomized placebo-controlled trials of valproic acid	46
Table 17. Characteristics of double-blind head-to-head randomized controlled trials of acamprosate versus naltrexone	48
Table 18. Characteristics of double-blind head-to-head randomized controlled trials of disulfiram versus naltrexone	49
Table 19. Characteristics of double-blind head-to-head randomized controlled trials including medications used off-label, or those under investigation.....	52
Table 20. Characteristics of included double-blind randomized placebo-controlled trials of acamprosate that report a health outcome	56
Table 21. Mortality reported in placebo-controlled trials of acamprosate.....	58
Table 22. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone that report a health outcome	59
Table 23. Characteristics of included double-blind randomized placebo-controlled trials of medications used off-label, or those under investigation.....	62
Table 24. Characteristics of head-to-head randomized controlled trials reporting a health outcome	64
Table 25. Characteristics of head-to-head randomized controlled trials including medications used off-label, or those under investigation	68
Table 26. Characteristics of studies included for KQ 3 that were not in KQ 1 or 2	70
Table 27. Results of meta-analyses for adverse events: acamprosate compared with placebo	71
Table 28. Results of meta-analyses for adverse events: naltrexone compared with placebo	74
Table 29. Results of meta-analyses for adverse events: acamprosate compared with naltrexone	74
Table 30. Characteristics of included randomized controlled trials of FDA-approved medications for treating alcohol dependence in primary care settings	76
Table 31. Characteristics of head-to-head medication studies that evaluated subgroups	78
Table 32. Characteristics of included studies that assessed the association between opioid receptor gene polymorphisms and naltrexone response	82
Table 33. Results of included studies that assessed the association between mu-opioid receptor gene polymorphisms and naltrexone response	83
Table 34. Summary of findings and strength of evidence for efficacy of FDA-approved medications for alcohol dependence	87
Table 35. Summary of findings and strength of evidence for comparative effectiveness of acamprosate and naltrexone	88
Table 36. Evidence gaps for future research, by key question	95

Figures

Figure A. Analytic framework for pharmacotherapy for adults with alcohol-use disorders in outpatient settings	ES-5
Figure B. Disposition of articles	ES-8

Figure 1. Analytic framework for pharmacotherapy for adults with alcohol-use disorders in outpatient settings	6
Figure 2. Disposition of articles	13

Appendixes

Appendix A. Search Strategy	
Appendix B. Excluded Studies	
Appendix C. Criteria Used for Evaluating Studies' Risk of Bias	
Appendix D. Strength of Evidence Assessments	
Appendix E. Placebo-Controlled Trials of Medications Used Off-label, or Those Under Investigation for Which We Found Only 1 Trial Meeting Inclusion Criteria	
Appendix F. Meta-Analyses	
Appendix G. Additional Studies of Genetic Polymorphisms Meeting Inclusion Criteria, but with Only 1 Study for a Drug-Polymorphism Pair	

Executive Summary

Background

Alcohol misuse, or unhealthy alcohol use, which includes the full spectrum from drinking above recommended limits (i.e., risky/hazardous drinking) to alcohol dependence,^{1,2} is associated with numerous health and social problems, more than 85,000 deaths per year in the United States,^{3,4} and an estimated annual cost to society of more than \$220 billion.^{5,6} Alcohol misuse is estimated to be the third leading cause of preventable mortality in the United States, following tobacco use and being overweight.⁷ For this report, we use the definitions in Table A.

Alcohol-use disorders (AUDs) include harmful use, alcohol abuse, and alcohol dependence.^{8,9} Prevalence of AUDs is higher for men than for women, with estimates indicating a lifetime risk of more than 20 percent for men.^{8,10-12} Alcohol dependence has lifetime prevalence rates of about 17 percent for men and 8 percent for women.¹³

AUDs cause substantial morbidity and mortality—that is, threefold to fourfold increased rates of early mortality.¹⁴⁻¹⁶ They are associated with hypertension, heart disease, stroke, cancer, liver cirrhosis, amnesias, cognitive impairment, sleep problems, peripheral neuropathy, gastritis and gastric ulcers, pancreatitis, decreased bone density, anemia, depression, insomnia, anxiety, suicide, and fetal alcohol syndrome.^{8,17} Excessive alcohol consumption is also a major factor in injury and violence.¹⁸ Acute alcohol-related harm can be the result of fires, drowning, falls, homicide, suicide, motor vehicle crashes, child maltreatment, and pedestrian injuries.¹⁹ In addition, AUDs can complicate the assessment and treatment of other medical and psychiatric problems.⁸

Treatments for Alcohol-Use Disorders

Treatments for AUDs continue to evolve as research on the effectiveness of various treatments is published, and new treatments are introduced and used more frequently. No single best approach has yet proven superior among the variety of available treatment options. Some common treatments for AUDs include cognitive behavioral therapy, motivational enhancement therapy, 12-step programs (e.g., Alcoholics Anonymous), and pharmacotherapy. Treatment may be delivered via individual outpatient counseling, intensive outpatient programs using group or individual methods, alcoholism treatment centers, or other approaches.

Over the past 15 to 20 years, awareness has grown that treatment may still be beneficial even if complete abstinence is not achieved. As a result, research has used other outcomes to measure the effectiveness of treatment, which can be subsumed under the concept of harm reduction.²⁰ These measures include significant increases in abstinent days or decreases in heavy drinking episodes, improved physical health, and improvements in psychosocial functioning.

Table A. Definitions of the spectrum of alcohol misuse

Term	Definition
Risky or hazardous use	Consumption of alcohol above recommended daily, weekly, or per-occasion amounts. Consumption levels that increase the risk for health consequences.
Harmful use ^{21,22}	A pattern of drinking that is already causing damage to health. The damage may be either physical (e.g., liver damage from chronic drinking) or mental (e.g., depressive episodes secondary to drinking).
Alcohol abuse ²³	<p>A. A maladaptive pattern of alcohol use leading to clinically significant impairment or distress, as manifested by at least 1 of the following occurring within a 12-month period:</p> <ol style="list-style-type: none"> (1) recurrent alcohol use resulting in a failure to fulfill major role obligations at work, school, or home (e.g., repeated absences or poor work performance related to alcohol use; alcohol-related absences, suspensions, or expulsions from school; neglect of children or household); (2) recurrent alcohol use in situations in which it is physically hazardous (e.g., driving an automobile or operating a machine when impaired); (3) recurrent alcohol-related legal problems (e.g., arrests for alcohol-related disorderly conduct); or (4) continued alcohol use despite having persistent or recurrent social or interpersonal problems caused or exacerbated by the effects of alcohol (e.g., arguments with spouse about consequences of intoxication, physical fights). <p>B. The symptoms have never met the criteria for alcohol dependence.</p>
Alcohol dependence ²³ (alcoholism, alcohol addiction)	<p>A maladaptive pattern of alcohol use, leading to clinically significant impairment or distress, as manifested by at least 3 of the following occurring at any time in the same 12-month period:</p> <ol style="list-style-type: none"> (1) tolerance, as defined by either of the following: <ol style="list-style-type: none"> (a) a need for markedly increased amounts of alcohol to achieve intoxication or desired effect; or (b) markedly diminished effect with continued use of the same amount of alcohol; (2) withdrawal, as manifested by either of the following: <ol style="list-style-type: none"> (a) the characteristic withdrawal syndrome for alcohol; or (b) alcohol (or a closely related drug) is taken to relieve or avoid withdrawal symptoms; (3) alcohol is often taken in larger amounts or over a longer period than was intended; (4) there is a persistent desire or unsuccessful efforts to cut down or control alcohol use; (5) a great deal of time is spent in activities necessary to obtain alcohol, use alcohol, or recover from its effects; (6) important social, occupational, or recreational activities are given up or reduced because of alcohol use; or (7) alcohol use is continued despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by alcohol (e.g., continued drinking despite recognition that an ulcer was made worse by alcohol consumption).

Pharmacological Interventions

Beginning in the 1950s, the pharmacotherapy for alcohol dependence consisted only of disulfiram, an aversive deterrent that produces very uncomfortable symptoms when alcohol is consumed. Since the 1990s, two oral medications (naltrexone and acamprosate) and one long-acting intramuscular formulation (of naltrexone) have been approved by the U.S. Food and Drug Administration (FDA) for alcohol dependence. Table B describes the medications available in the United States that are FDA-approved, their mechanism of action, and dosing. Many additional medications have been used off-label or studied for treatment of AUDs. These include antidepressants, mood stabilizers, anticonvulsants, alpha-adrenergic blockers, antipsychotics, and anxiolytics.

Table B. Medications that are FDA-approved for treating adults with alcohol dependence

Generic Drug Name	Mechanism	Dosing
Acamprosate	Thought to modulate hyperactive glutamatergic NMDA receptors	666 mg 3 times per day
Disulfiram	Inhibits ALDH2, causing accumulation of acetaldehyde during alcohol consumption, which produces a variety of adverse effects such as nausea, dizziness, flushing, and changes in heart rate and blood pressure	250 to 500 mg per day
Naltrexone	Opioid antagonist; competitively binds to opioid receptors and blocks the effects of endogenous opioids such as β -endorphin	Oral: 50 to 100 mg per day Intramuscular injection: 380 mg per month

Abbreviations: ALDH2 = aldehyde dehydrogenase; FDA = U.S. Food and Drug Administration; mg = milligram; NMDA = N-methyl-D-aspartate.

Despite ongoing developments and advancements in treatment approaches, alcohol dependence represents one of the most undertreated disorders in the U.S. health care system; it is estimated that only 1 in 4 individuals with alcohol dependence receives treatment.¹³ Furthermore, of those patients who receive treatment, less than 1 in 10 receives medication as part of his or her treatment. Therefore, expanding awareness and access to this relatively new treatment modality has the potential to improve outcomes and reduce the burden of this devastating illness that affects millions.

Existing Guidance

The Veterans Administration (VA), National Institute on Alcohol Abuse and Alcoholism (NIAAA), and Substance Abuse and Mental Health Services Administration (SAMHSA) all have guidelines addressing the use of pharmacotherapy for alcohol dependence.²⁴⁻²⁶ The VA guidelines recommend that oral naltrexone and/or acamprosate routinely be considered for patients with alcohol dependence (although acamprosate is currently a nonformulary medication for the VA), and that medications be offered in combination with addiction-focused counseling.

The United Kingdom's National Institute for Clinical Excellence (NICE) guidelines include the following recommendations: (1) after a successful withdrawal for people with moderate or severe alcohol dependence, to consider offering acamprosate or oral naltrexone in combination with an individual psychological intervention (cognitive behavioral therapies, behavioral therapies, or social network and environment-based therapies) focused specifically on alcohol misuse; (2) to consider offering disulfiram in combination with a psychological intervention for people who have a goal of abstinence but for whom acamprosate and oral naltrexone are not suitable, or who prefer disulfiram and understand the relative risks of taking the drug; and (3) to have specialist and competent staff administer pharmacological interventions.⁹

Scope and Key Questions

The use of medications for AUDs is associated with uncertainty and variation across providers and settings. Since the last report commissioned by the Agency for Healthcare Research and Quality (AHRQ) on medications for alcohol dependence (1999),^{27,28} there has been more than a 10-fold increase in the number of individuals studied in controlled clinical trials of naltrexone and acamprosate, and many trials of medications that are not FDA-approved. Other reasons for conducting a new review on this topic include the following: (1) to assess the comparative effectiveness of the FDA-approved medications; (2) to determine whether any agents that are not FDA-approved have evidence supporting their efficacy; (3) to evaluate the evidence on intramuscular naltrexone (Vivitrol®), a fairly recently approved medication; (4) to

evaluate whether trials provide evidence of effectiveness in primary care settings; (5) to assess whether some medications are more or less effective for adults with certain genetic polymorphisms; and (6) to inform updates to clinical practice guidelines.

Our report focuses on clinically relevant medications—those that are commonly used, those with sufficient literature for systematic review, and those of greatest interest to clinicians and to the developers of guidelines. Our report is limited to people with AUDs; it does *not* address those with risky or hazardous alcohol use (for whom medications are likely not an appropriate intervention).


The main objective of this report is to conduct a systematic review and meta-analysis of the comparative effectiveness and harms of medications for adults with AUDs. In this review, we address the following Key Questions (KQs):

- KQ 1a: Which medications are efficacious for improving consumption outcomes for adults with AUDs in outpatient settings?
- KQ 1b: How do medications for adults with AUDs compare for improving consumption outcomes in outpatient settings?
- KQ 2a: Which medications are efficacious for improving health outcomes for adults with AUDs in outpatient settings?
- KQ 2b: How do medications for adults with AUDs compare for improving health outcomes in outpatient settings?
- KQ 3a: What adverse effects are associated with medications for adults with AUDs in outpatient settings?
- KQ 3b: How do medications for adults with AUDs compare for adverse effects in outpatient settings?
- KQ 4: Are medications for treating adults with AUDs effective in primary care settings?
- KQ 5: Are any of the medications more or less effective than other medications for men or women, older adults, young adults, racial or ethnic minorities, smokers, or those with co-occurring disorders?
- KQ 6: Are any of the medications more or less effective for adults with certain genetic polymorphisms (e.g., of the mu-opioid receptor gene [*OPRM1*]) compared with adults without such polymorphisms?

Analytic Framework

We developed an analytic framework to guide the systematic review process (Figure A).

Figure A. Analytic framework for pharmacotherapy for adults with alcohol-use disorders in outpatient settings



Methods

Literature Search Strategy

To identify articles relevant to each KQ, we searched PubMed®, the Cochrane Library, PsycINFO®, CINAHL®, and EMBASE® for English-language and human-only studies published from January 1, 1970, to February 5, 2013. Searches were run by an experienced Evidence-based Practice Center (EPC) librarian and were peer-reviewed by another EPC librarian. We manually searched reference lists of pertinent reviews, trials, and background articles on this topic to look for any relevant citations that our searches might have missed.

We searched for unpublished studies relevant to this review using ClinicalTrials.gov, the World Health Organization International Clinical Trials Registry Platform, and the FDA Web site. In addition, AHRQ's Scientific Resource Center requested any unpublished studies and pertinent data from relevant pharmaceutical companies.

Eligibility Criteria

We developed inclusion and exclusion criteria with respect to populations, interventions, comparators, outcomes, timing, and setting (PICOTS) and study designs. We included studies enrolling adults with AUDs that evaluated one or more of the following medications: acamprosate, disulfiram, naltrexone, amitriptyline, aripiprazole, atomoxetine, baclofen, buspirone, citalopram, desipramine, escitalopram, fluoxetine, fluvoxamine, gabapentin, imipramine, nalmefene, olanzapine, ondansetron, paroxetine, prazosin, quetiapine, sertraline, topiramate, valproate, varenicline, and viloxazine.

Studies were required to assess at least one of the following outcomes: return to any drinking (lapse), return to heavy drinking (relapse), drinking days, heavy drinking days, drinks per drinking day, time to lapse or relapse, accidents, injuries, quality of life (QoL), function, mortality, or adverse effects. Studies were required to treat patients with a medication for a minimum of 12 weeks in an outpatient setting.

For KQs 1, 2 and 4, double-blind randomized controlled trials (RCTs) that compared one of the medications with placebo or another medication and recent systematic reviews (searches ending no earlier than 2007) were eligible. For KQ 2b, prospective cohort studies were also eligible. For KQ 3 (harms), double-blind RCTs and recent systematic reviews that compared one of the medications with placebo or with another medication were eligible. The following designs were also eligible if they compared 2 or more drugs of interest: nonrandomized controlled trials, open-label trials, secondary analyses or subgroup analyses from trials, prospective cohort studies, and case-control studies. For KQ 5 (subgroups), double-blind RCTs, recent systematic reviews, nonrandomized controlled trials, open-label trials, secondary analyses or subgroup analyses from trials, prospective cohort studies, and case-control studies were eligible, as long as the studies compared 2 or more drugs. For KQ 6 (genetic polymorphisms), double-blind RCTs, secondary analyses or subgroup analyses from trials, and prospective cohort studies comparing people with a genetic polymorphism with people without the polymorphism were eligible.

Study Selection

Two members of the research team independently reviewed each title and abstract (identified through searches) to determine eligibility. Studies marked for possible inclusion by either reviewer and those that lacked adequate information to determine eligibility underwent a full-text review. Two members of the team independently reviewed each full-text article to determine eligibility. If the reviewers disagreed, they resolved conflicts by discussion and consensus or by consulting a senior member of the team.

Data Extraction

We designed and used structured data extraction forms to gather pertinent information from each article; this included characteristics of study populations, settings, interventions, comparators, study designs, methods, and results. Trained reviewers extracted the relevant data from each included article. All data extractions were reviewed for completeness and accuracy by a second member of the team.

Risk-of-Bias Assessment of Individual Studies

To assess the risk of bias (internal validity) of studies for major outcomes of interest, we used predefined criteria based on guidance from the AHRQ *Methods Guide for Effectiveness and*

Comparative Effectiveness Reviews.²⁹ We assessed selection bias, confounding, performance bias, detection bias, and attrition bias; we included questions about adequacy of randomization, allocation concealment, similarity of groups at baseline, blinding, attrition, whether intention-to-treat (ITT) analysis was used, methods of handling missing data, and fidelity. We rated the studies as low, medium, high, or unclear risk of bias.³⁰ Two independent reviewers assessed the risk of bias for each study. Disagreements between the two reviewers were resolved by discussion and consensus or by a third member of the team.

Data Synthesis

We conducted meta-analyses using random-effects models to estimate pooled effects.³¹ For continuous outcomes, we used weighted mean differences (WMDs). For binary outcomes, we calculated risk differences (RDs) between groups. We did not include studies rated as high or unclear risk of bias in our main analyses, but did include them in sensitivity analyses. We calculated the chi-squared statistic and the I^2 statistic to assess statistical heterogeneity in effects between studies.^{32,33} We also examined potential sources of heterogeneity by analysis of subgroups defined by patient population (e.g., U.S. versus non-U.S. studies) and variation in interventions (e.g., dose). When quantitative synthesis was not appropriate (e.g., because of clinical heterogeneity, insufficient numbers of similar studies, or insufficiency or variation in outcome reporting), we synthesized the data qualitatively.

Strength of the Body of Evidence

We graded the strength of evidence (SOE) as high, moderate, low, or insufficient based on established guidance.³⁴ Developed to grade the overall strength of a body of evidence, the approach incorporates four key domains: risk of bias (includes study design and aggregate quality), consistency, directness, and precision of the evidence. It also considers optional domains. Two reviewers assessed each domain for each key outcome and determined an overall SOE grade based on domain ratings. In the event of disagreements on the domain or overall grade, they resolved differences by discussion or by consulting an experienced investigator. We graded the SOE for the following outcomes: return to any drinking, return to heavy drinking, drinking days, heavy drinking days, drinks per drinking day, accidents, injuries, QoL or function, mortality, and adverse events.


Applicability

We assessed applicability of the evidence following guidance from the *Methods Guide for Comparative Effectiveness Reviews*.³⁵ We used the PICOTS framework to explore factors that affect applicability.

Results

We included 157 published articles reporting on 130 studies; 114 were RCTs (Figure B).

Figure B. Disposition of Articles



Key Question 1. Consumption Outcomes

We found moderate SOE that both acamprosate and naltrexone are effective for improving alcohol consumption outcomes (Table C). Numbers needed to treat (NNT) to prevent 1 person from returning to any drinking were 10 and 25, respectively. For return to heavy drinking, evidence did not support the efficacy of acamprosate, whereas naltrexone was efficacious with an NNT of 13. Evidence from well-controlled trials does not adequately support the efficacy of

disulfiram compared with placebo for preventing return to any drinking or for other alcohol consumption outcomes. Some disulfiram trials reported fewer drinking days for subjects who returned to any drinking and who had a complete set of assessment interviews, and suggest that disulfiram may have a role in the treatment of alcohol dependence for some individuals.

Table C. Summary of findings and strength of evidence for efficacy of FDA-approved medications for alcohol dependence

Intervention	Outcome	N studies; N subjects ^a	Results Effect Size (95% CI) ^b	Strength of Evidence
Acamprosate	Return to any drinking	15; 4,747	RD: -0.10 (-0.15 to -0.05); NNT 10	Moderate
	Return to heavy drinking	6; 2,239	RD: -0.01 (-0.05 to 0.03)	Moderate
	Percentage drinking days	12; 4,385	WMD: -9.4 (-13.8 to -5.0)	Moderate
	Percentage heavy drinking days	0; 0	NA	Insufficient
	Drinks per drinking day	1; 116	WMD: 0.4 (-1.8 to 2.6)	Insufficient
	Accidents or injuries	0; ^c 0	NA	Insufficient
	QoL or function	1; 612	NSD	Insufficient
	Mortality	7; 2,477	7 events (ACA) vs. 5 events (placebo)	Insufficient
Disulfiram	Return to any drinking	2; 492	RD: 0.04 (-0.03 to 0.11) ^d	Low
	Return to heavy drinking	0; 0	NA	Insufficient
	Percentage drinking days	2; 290	NSD ^e	Insufficient
	Percentage heavy drinking days	0; 0	NA	Insufficient
	Drinks per drinking day	0; 0	NA	Insufficient
	Accidents or injuries	0; 0	NA	Insufficient
	QoL or function	0; 0	NA	Insufficient
	Mortality	0; 0	NA	Insufficient
Naltrexone	Return to any drinking	21; 4,232	RD: -0.04 (-0.07 to -0.01); NNT 25	Moderate
	Return to heavy drinking	21; 3,794	RD: -0.08 (-0.12 to -0.04); NNT 13	Moderate
	Percentage drinking days	19; 3,329	WMD: -4.6 (-6.6 to -2.5)	Moderate
	Percentage heavy drinking days	10; 1,423	WMD: -3.6 (-5.9 to -1.4)	Moderate
	Drinks per drinking day	11; 1,422	WMD: -0.5 (-1.0 to -0.07)	Low
	Accidents or injuries	0; 0	NA	Insufficient
	QoL or function	4; 1,513	Some conflicting results ^f	Insufficient
	Mortality	6; 1,738	1 event (NTX) vs. 2 events (placebo)	Insufficient

^a Includes only studies rated as low or medium risk of bias included in the main analyses; these numbers do not include studies rated as high or unclear risk of bias that were included in sensitivity analyses.

^b Negative effect sizes favor intervention over placebo/control.

^c One study rated as unclear risk of bias reported that one patient in the placebo group died by “accident.” No other details on the cause or nature of the accident were provided.³⁶

^d From meta-analysis of disulfiram 250 mg versus control (disulfiram 1 mg).^{37,38} Meta-analysis including studies rated as high risk of bias also found no significant difference (RD, -0.00; 95% CI, -0.10 to 0.09). Similarly, our meta-analysis found no statistically significant difference between disulfiram 250 mg per day and riboflavin (i.e., no disulfiram) (RD, -0.04; 95% CI, -0.11 to 0.03).

^e One study (N=128) reported similar percentages and no significant difference;³⁸ the other reported that disulfiram was favored among the subset of subjects (N=162 of 605 subjects) who drank and had a complete set of assessment interviews, but it did not report this outcome for the full randomized sample.³⁷ Overall, evidence was insufficient due to imprecision, inconsistency, and indirectness.

^f Unable to pool data. Two studies found no significant difference between naltrexone- and placebo-treated subjects.^{39,40} One study reported that patients receiving injectable naltrexone 380 mg per day had greater improvement on the mental health summary score than those receiving placebo at 24 weeks (8.2 versus 6.2, $p=0.044$).^{41,210} One study measured alcohol-related consequences (with the DrInC) and reported that more subjects who received placebo ($N=34$) had at least 1 alcohol-related consequence than those who received naltrexone ($N=34$): 76 percent versus 45 percent, $p=0.02$.⁴²

Abbreviations: ACA = acamprosate; CI = confidence interval; FDA = U.S. Food and Drug Administration; N = number; NA = not applicable; NNT = number needed to treat; NSD = no statistically significant difference; NTX = naltrexone; RD = risk difference; vs. = versus; WMD = weighted mean difference.

Our meta-analyses of 3 head-to-head RCTs comparing acamprosate with naltrexone,⁴³⁻⁴⁵ all rated as low risk of bias, found no statistically significant difference between the two medications for improvement in alcohol consumption outcomes (Table D). The COMBINE study was one of the 3 RCTs.⁴³ It found that patients receiving medical management with naltrexone, combined behavioral intervention (CBI), or both had better drinking outcomes than those who received placebo, but acamprosate showed no evidence of efficacy, with or without CBI.

Table D. Summary of findings and strength of evidence for comparative effectiveness of acamprosate and naltrexone

Intervention	Outcome	N studies; N subjects ^a	Results Effect Size (95% CI) ^b	Strength of Evidence
ACA vs. NTX	Return to any drinking	3; 800	RD: 0.02 (-0.03 to 0.08)	Moderate
	Return to heavy drinking	3; 800	RD: 0.01 (-0.06 to 0.07)	Moderate
	Percentage drinking days	2; 720	WMD: -2.98 (-13.4 to 7.5)	Low

^a Includes only studies rated as low or medium risk of bias included in the main analyses; these numbers do not include studies rated as high or unclear risk of bias that were included in sensitivity analyses.

^b Negative effect sizes favor acamprosate over naltrexone.

Note: Table only includes comparisons of medications with evidence of efficacy (as determined in KQ 1) and with sufficient data for synthesis. We did not include rows in this table for outcomes that we graded as insufficient SOE (percentage heavy drinking days, drinks per drinking day, accidents or injuries, QoL or function, and mortality).

Abbreviations: ACA = acamprosate; CI = confidence interval; N = number; NTX = naltrexone; QoL = quality of life; RD = risk difference; vs. = versus; WMD = weighted mean difference.

For the vast majority of medications used off-label, and those under investigation, evidence was either insufficient to determine whether they are efficacious for reducing alcohol consumption or evidence suggested that they are not efficacious for people with alcohol dependence. We found two exceptions. First, for topiramate, we found moderate SOE supporting efficacy for reducing drinking days, heavy drinking days (WMD, -11.5; 95% CI, -18.3 to -4.8), and drinks per drinking day (WMD, -1.1; 95% CI, -1.7 to -0.4)—based on the results of 2 RCTs (total $N=521$).^{46,47} The included RCTs did not report data for return to any drinking or return to heavy drinking. Second, for nalmefene, we found moderate SOE supporting efficacy for one alcohol consumption outcome—reduction in drinks per drinking day (WMD, -1.0; 95% CI, -1.8 to -0.3). However, the magnitude of benefit (reduction of 1 drink per drinking day) is not likely clinically significant, and we found insufficient evidence of efficacy for nalmefene for other consumption outcomes (return to any drinking, return to heavy drinking, and heavy drinking days) and low SOE that nalmefene is not efficacious for reducing drinking days (WMD, -1.1; 95% CI, -7.6 to 5.4). [Note: we are aware that new evidence on nalmefene has been published after our literature search and that nalmefene has since been approved in other countries; this

new evidence will be included in our update search while the report is being reviewed and any necessary changes will be made for our final report].

Key Question 2. Health Outcomes

We found insufficient direct evidence to conclude that treatment with acamprosate or naltrexone leads to improvement in health outcomes—i.e., accidents, injuries, QoL, function, or mortality. Very few trials reported any health outcomes, and the included trials were not designed or powered to assess impact on health outcomes—they typically focused on alcohol consumption outcomes. The largest pharmacotherapy trial in alcohol dependence, COMBINE, did report some evidence of improvement in QoL with naltrexone plus behavioral intervention (on the SF-12v2 physical health scale), but the difference between groups did not reach a clinically meaningful threshold.⁴⁰

Key Question 3. Harms

Evidence for many potential adverse events was insufficient to determine whether the risk was increased or not, often primarily because of lack of precision. For most of the specific adverse events, point estimates favored placebo (i.e., there were more adverse events with medications), but differences were not statistically significant. In head-to-head studies, the risk of withdrawal due to adverse events was not significantly different between acamprosate and naltrexone, whereas the risk of headache was higher for those treated with naltrexone. Compared with placebo, patients treated with acamprosate had a higher risk of anxiety, diarrhea, and vomiting, and those treated with naltrexone had a higher risk of dizziness, nausea, and vomiting. Trials of topiramate reported a significantly increased risk of many adverse events, including paresthesias, taste perversion, anorexia, difficulty with concentration/attention, nervousness, dizziness, pruritis, psychomotor slowing, and weight loss.^{46,47}

Key Question 4. Evidence from Primary Care Settings

We identified no eligible trials conducted completely in primary care settings, and no eligible trials assessing FDA-approved medications that were conducted in primary care settings. The only included trial conducted partly in primary care settings compared nalmefene with placebo in 15 sites (about half were primary care settings) in Finland.⁴⁸ See Discussion section below (under Primary Care) for more information about studies that may have applicability to primary care settings.

Key Question 5. Subgroups

We did not find any compelling evidence that naltrexone, acamprosate, or topiramate are more or less effective (compared with each other) for men or women, older adults, young adults, racial or ethnic minorities, smokers, or those with co-occurring disorders.

Key Question 6. Genetic Polymorphisms

For genetic polymorphisms, we found no studies that assessed the clinical utility of genotype-guided dosing strategies or genotype-guided medication selection, and none that randomized by genotype. All included studies were either secondary/subgroup analyses of trials or prospective cohort studies of people treated with a medication, and all assessed the association

between genotype and response to medication (i.e., clinical validity). For most polymorphism-medication pairs, we found just 1 eligible study, and we graded the SOE as insufficient.

Seven eligible studies assessed variation in naltrexone response related to mu-opioid receptor gene (*OPRM1*) polymorphisms. Our meta-analyses for return to any drinking and return to heavy drinking found no significant difference between AA homozygotes and those with at least one G allele, both without inclusion of studies rated as high or unclear risk of bias (RD, -0.03; 95% CI, -0.6 to 0.5 and RD, 0.26; 95% CI, -0.01 to 0.53, respectively) and with them (RD, 0.01; 95% CI, -0.2 to 0.2 and RD, 0.14; 95% CI, -0.03 to 0.3, respectively). Point estimates for return to heavy drinking suggest it is possible that patients with at least one G allele might be more likely to respond to naltrexone, but confidence intervals were wide; additional studies are needed to improve confidence in the estimate of the effect.

Discussion

Evidence supports the efficacy of more than one pharmacological treatment for alcohol dependence, and clinical uncertainty exists about what treatment to select for individual patients. Acamprosate and naltrexone have the best evidence supporting their efficacy, but head-to-head trials have not consistently established superiority of either medication. Thus, other factors may contribute to medication choices, such as heterogeneity of alcohol dependence, coexisting symptoms such as anxiety or insomnia, frequency of administration, cost, potential type of benefits, potential adverse events, and availability of treatments (e.g., acamprosate is currently a nonformulary medication for the VA).

For example, acamprosate is typically dosed as two 333 mg tablets given three times daily, whereas oral naltrexone is one tablet once daily, and intramuscular naltrexone is given once monthly. Acamprosate is contraindicated for people with severe renal impairment and requires dose adjustments for moderate renal impairment. Naltrexone is contraindicated for patients with acute hepatitis or liver failure (and has precautions for other hepatic disease), and for those currently using opioids or with anticipated need for opioids, and it can precipitate severe withdrawal for patients dependent on opioids. Larger doses may be required and respiratory depression may be deeper and more prolonged if opioid analgesia is needed.

Given that medications for alcohol dependence have been underutilized, entities providing health care for people with alcohol dependence may need to develop systems to optimize dissemination and implementation. For example, these could include campaigns to educate providers about the use of medications for alcohol dependence; systems to screen for unhealthy alcohol use and to provide appropriate interventions for people with unhealthy alcohol use; systems to ensure that people with alcohol dependence have access to knowledgeable providers who can prescribe medications; or systems to remind or incentivize providers to use effective medications for alcohol dependence when appropriate.

Although we found insufficient direct evidence to conclude that treatment with medications leads to improvement in health outcomes—i.e., accidents, injuries, QoL, function, or mortality—evidence from epidemiologic literature consistently relates high average alcohol consumption and heavy per-occasion use to an increased risk of health problems, such as cancers of the oral cavity, esophagus, larynx, colon, rectum, liver, and breast; liver cirrhosis; chronic pancreatitis; coronary heart disease; stroke; depression; preterm birth complications; fetal alcohol syndrome; and injuries and violence.^{1,17,49-51} Such epidemiologic evidence would suggest that improving alcohol consumption outcomes is likely to result in improved health outcomes.

Primary Care

Direct evidence in primary care settings was scant. The only included trial conducted partly in primary care settings compared nalmefene with placebo in 15 sites (about half were primary care settings) in Finland.⁴⁸ One other trial (included in KQ 1, but not in KQ 4) that compared naltrexone with placebo for 12 weeks in the United States described the use of a “primary care model.”⁵² Although the trial did not take place in a primary care setting (it was a treatment research center), and the investigators were from a department of psychiatry, the psychosocial co-intervention was delivered by a nurse practitioner with a primary care background, and the trial may have implications for how psychosocial co-interventions could be provided in primary care settings.

Two other publications that did not meet our inclusion criteria (due to the study design or comparators) may have important implications for the use of medications for alcohol dependence in primary care settings. First, a nested sequence of 3 U.S.-based RCTs compared naltrexone plus “primary care management” (PCM) with naltrexone plus cognitive behavioral therapy.⁵³ PCM was provided by nurse practitioners, physician assistants, and one internist in an initial 45-minute visit, followed by 15- to 20-minute sessions in weeks 1, 2, 3, 4, 6, 8, and 10. The study found no difference in response to treatment, as measured by avoiding persistent heavy drinking, between those who received PCM and those who received cognitive behavioral therapy (84.1 percent versus 86.5 percent). Among responders enrolled in a maintenance trial, it found higher response for those who received naltrexone and PCM than for those who received placebo and PCM (80.8 percent versus 51.9 percent, $p=0.03$). Second, a pragmatic trial with 149 general practitioners in France who were “used to managing alcohol-dependent patients in their daily practice” randomized patients ($N=422$) to acamprosate plus standard care or standard care alone.⁵⁴ Standard care in France was described as typically consisting of outpatient detoxification followed by a rehabilitation program (involving some type of psychotherapy). The trial reported better outcomes for the acamprosate group for the Alcohol-Related Problems Questionnaire score, the number of subjects with no alcohol-related problems, and for all secondary outcome measures, including QoL.

Barriers to prescribing medications for alcohol dependence in primary care may include lack of familiarity with the medications, lack of confidence in their effectiveness, or inability to provide suitable psychosocial co-interventions (e.g., due to competing demands or insufficient practice resources, personnel, or training). Like behavioral counseling interventions for risky drinking delivered in primary care, implementing the use of medications and psychosocial co-interventions for alcohol dependence in primary care might require development of support systems and additional provider and staff training.^{1,4} Further, primary care providers are typically trained to refer patients with alcohol dependence for specialized treatment. O’Malley and O’Connor recently reviewed the issues surrounding the use of medications for alcohol dependence in primary care settings.⁵⁵ They concluded that “the implementation and widespread use of medications to treat alcohol problems faces a unique set of barriers in primary care. Although primary care providers are proficient at prescribing a wide variety of medications, they generally are unfamiliar with medications for treating alcohol problems other than those used to treat alcohol withdrawal.” They referenced a body of research to support basic screening methods, brief interventions, and medication therapy that has yet to have a major impact on how primary care providers care for individuals at risk for or with alcohol problems.⁵⁶

Applicability

Most studies reported that 100 percent of subjects met criteria for alcohol dependence. We did not identify any studies that evaluated medications and reported them to be efficacious for people with AUDs who did not meet criteria for alcohol dependence (i.e., people with alcohol abuse or harmful alcohol use). The mean age of subjects was generally in the 40s, with very few studies enrolling slightly younger or older populations. Thus, it is uncertain whether the medications have similar efficacy for older (e.g., those 65 and older) or younger (e.g., in the 20s) subgroups. We did not find evidence to confirm or refute whether treatments are more or less efficacious for gender groups, racial or ethnic minorities, smokers or nonsmokers, and those with certain coexisting conditions.

Although the majority of included trials assessing the efficacy of acamprosate were conducted in Europe (15 of 20) and a minority were conducted in the United States (3 of 20), the opposite was true for naltrexone (27 of 42 in the United States and 6 of 42 in Europe). Further, the few studies of acamprosate conducted in the United States did not find it to be efficacious. It is unclear whether the different results were due to population differences or other factors. The European trials of acamprosate typically identified patients from inpatient settings or treatment programs, whereas the U.S.-based trials of acamprosate relied on advertisements and referrals. It is possible that this resulted in populations with differing alcoholism severity and differing potential for benefit. For example, studies of subjects recruited via advertisements may enroll people who have less severe disorders.

Most studies required patients to abstain for at least a few days prior to initiating medication, and the medications are generally recommended for maintenance of abstinence. Acamprosate and injectable naltrexone are only approved for use in patients who have established abstinence, though the duration of required abstinence is not set. However, some studies enrolling patients who were not yet abstinent have reported reduction in heavy drinking with naltrexone⁵⁷ or acamprosate.⁵⁸

Limitations of the Comparative Effectiveness Review Process

The scope of this review was focused on medications. We did not evaluate the effectiveness or comparative effectiveness of other interventions for AUDs (e.g., 12-step programs). We required that trials have at least 12 weeks of followup from the time of medication initiation, excluding trials of shorter duration. Some might consider this approach to omit potentially important information. However, longitudinal studies have found that treatment periods of less than 6 months' duration may yield misleading conclusions about treatment efficacy, due to fluctuations in drinking behavior that are typical of the course of alcoholism^{59,60}—suggesting that a longer duration of followup (6 months or more) might more accurately reflect the outcomes of greatest interest and importance.

Finally, publication bias and selective reporting are potential limitations. Although we searched for unpublished studies and unpublished outcomes, and did not find direct evidence of either of these biases, many of the included trials were published prior to the availability of trial registries (e.g., clinicaltrials.gov) that would allow for greater certainty in determining the potential for either type of bias.

Limitations of the Evidence Base

The evidence base was inadequate to draw conclusions for some of our questions or subquestions of interest. In particular, as described above, we found insufficient direct evidence on health outcomes, limited and varying reporting on harms, no trials conducted completely in primary care settings, and scant head-to-head evidence on differences for population subgroups.

We found insufficient direct evidence to determine whether medications are efficacious for improving health outcomes. Although evidence from epidemiologic literature consistently relates high average and heavy per-occasion alcohol use to an increased risk of health problems, it is challenging to estimate the magnitude of reduction in the risk of health problems that is derived from a reduction in consumption. For example, it is unclear how much benefit (for health outcomes) is derived from 10 percent fewer patients returning to any drinking.

Many included trials had methodological limitations introducing some risk of bias. Some had high proportions of subjects lost to followup. High attrition rates are not uncommon in studies of psychiatric conditions, but methods of handling missing data varied, and some trials did nothing to address missing data (i.e., only analyzing completers). However, many trials conducted true ITT analyses and used appropriate methods of handling missing data, such as imputing return to heavy drinking for subjects lost to followup or using multiple imputation.

Future Research

We identified numerous gaps in the evidence that future research could address. Many of these gaps are highlighted in the previous sections of this Discussion. Of note, these gaps relate only to the KQs addressed by this report, and they should not eliminate a wide range of potentially important research that falls outside of our scope. Table E summarizes the key gaps and potential future research that could address the gaps.

Conclusions

Acamprosate and naltrexone are effective for improving alcohol consumption outcomes for patients with alcohol dependence (moderate SOE). NNTs to prevent 1 person from returning to any drinking were 10 and 25, respectively; NNT to prevent 1 person from returning to heavy drinking was 13 for naltrexone. Our meta-analyses of 3 head-to-head trials found no statistically significant difference between the two medications for improvement in alcohol consumption outcomes (moderate SOE). With the exception of topiramate, for which we found moderate SOE supporting efficacy for improving some consumption outcomes, current evidence does not establish the efficacy of medications used off-label and those under investigation for people with alcohol dependence. We found insufficient direct evidence to conclude whether medications for alcohol dependence are effective for improving health outcomes. No eligible trials assessing FDA-approved medications were conducted in primary care settings. Evidence was generally insufficient to determine comparative effectiveness of acamprosate and naltrexone for subgroups.

Table E. Evidence gaps for future research, by Key Question

KQ	Evidence Gap	Potential Future Research
1	Evidence was insufficient to determine efficacy of some medications.	Future studies could evaluate medications that have some evidence (often from 1 or 2 small trials) suggesting possible efficacy (e.g., buprenorphine) or medications that have not yet been studied with some theoretical basis to support their potential efficacy.
1	We found no head-to-head studies of oral naltrexone and injectable naltrexone.	Future studies could compare the benefits or harms of oral and injectable naltrexone.
1	We found insufficient evidence evaluating medications for people with alcohol use disorders who do not meet criteria for alcohol dependence (i.e., those with alcohol abuse or harmful alcohol use).	Future studies could evaluate the efficacy of acamprosate or naltrexone in such populations.
2	We found insufficient direct evidence to conclude that treatment with acamprosate or naltrexone leads to improvement in health outcomes.	Future studies could focus on health outcomes, such as accidents, injuries, QoL, function, or mortality.
3	Relatively few studies reported information about suicide, suicidal ideation, or self-harmful behaviors.	Additional studies could be conducted to determine whether precautions about suicide, suicidal thoughts, or self-harmful behaviors are warranted.
3	Little evidence was available to determine whether naltrexone can be used for people with various liver conditions. ^a	Future studies could evaluate the use of naltrexone for people with various chronic liver conditions.
4	No eligible trials assessed the use of FDA-approved medications in primary care settings.	Future studies could evaluate the use of acamprosate and naltrexone in primary care settings.
5	Evidence on whether any medications are more or less effective than other medications for population subgroups was scant.	Future studies could compare the use of acamprosate and naltrexone for subgroups of patients (e.g., enrolling subjects who all have depression or other psychiatric conditions; comparing effectiveness for men or women or among older or younger patients).
6	Relatively few subjects contributed data to our analyses of variation in naltrexone response and <i>OPRM1</i> polymorphisms. Patients with at least one G allele may be more likely to respond to naltrexone, but confidence intervals were wide and the effect was not statistically significant.	Additional studies are likely to change our confidence in the estimate of the effect and to change the estimate.
6	No studies assessed the clinical utility of genotype-guided dosing strategies or genotype-guided medication selection, and none randomized by genotype.	If variation in naltrexone response by <i>OPRM1</i> polymorphisms becomes established, then future studies could assess the clinical utility of using genotype-guided dosing strategies. For example, studies might compare the use of genotype-guided dosing strategies (e.g., use naltrexone for patients with at least one G allele, but use acamprosate for AA homozygotes) with using naltrexone or acamprosate for all subjects.
6	Only 1 study was available for most polymorphism-medication response associations.	Future studies could explore other genotypic associations (i.e., not limiting future studies to <i>OPRM1</i> polymorphisms).

^a The FDA removed the black box warning for hepatotoxicity for injectable naltrexone, but it is unclear whether naltrexone should be used in people with various chronic liver conditions.

Abbreviations: FDA = U.S. Food and Drug Administration; *OPRM1* = mu-opioid receptor gene; QoL = quality of life.

References

- Jonas DE, Garbutt JC, Amick HR, et al. Behavioral counseling after screening for alcohol misuse in primary care: a systematic review and meta-analysis for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2012 Nov 6;157(9):645-54. PMID: 23007881.
- Saitz R. Clinical practice. Unhealthy alcohol use. *N Engl J Med*. 2005 Feb 10;352(6):596-607. PMID: 15703424.
- Mokdad AH, Marks JS, Stroup DF, et al. Actual causes of death in the United States, 2000. *JAMA*. 2004 Mar 10;291(10):1238-45. PMID: 15010446.
- Jonas DE, Garbutt JC, Brown JM, et al. Screening, Behavioral Counseling, and Referral in Primary Care to Reduce Alcohol Misuse. Comparative Effectiveness Review No. 64. (Prepared by the RTI International-University of North Carolina Evidence-based Practice Center under Contract No. 290-2007-10056-I.) AHRQ Publication No. 12-EHC055-EF. Rockville, MD: Agency for Healthcare Research and Quality; July 2012. www.effectivehealthcare.ahrq.gov/reports/final.cfm.
- Bouchery EE, Harwood HJ, Sacks JJ, et al. Economic costs of excessive alcohol consumption in the U.S., 2006. *Am J Prev Med*. 2011 Nov;41(5):516-24. PMID: 22011424.
- Harwood HJ, Fountain D, Fountain G. Economic cost of alcohol and drug abuse in the United States, 1992: a report. *Addiction*. 1999 May;94(5):631-5. PMID: 10563025.
- Centers for Disease Control and Prevention. FastStats: Alcohol Use. Updated January 27, 2012. <http://www.cdc.gov/nchs/faststats/alcohol.htm>. Accessed May 21, 2012.
- Schuckit MA. Alcohol-use disorders. *Lancet*. 2009 Feb 7;373(9662):492-501. PMID: 19168210.
- National Collaborating Centre for Mental Health, National Institute for Health & Clinical Excellence. Alcohol-use Disorders: The NICE Guidelines on Diagnosis, Assessment and Management of Harmful Drinking and Alcohol Dependence. The British Psychological Society and The Royal College of Psychiatrists 2011. <http://www.nice.org.uk/nicemedia/live/13337/53190/53190.pdf>.
- Hasin DS, Stinson FS, Ogburn E, et al. Prevalence, correlates, disability, and comorbidity of DSM-IV alcohol abuse and dependence in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2007 Jul;64(7):830-42. PMID: 17606817.
- Mertens JR, Weisner C, Ray GT, et al. Hazardous drinkers and drug users in HMO primary care: prevalence, medical conditions, and costs. *Alcohol Clin Exp Res*. 2005 Jun;29(6):989-98. PMID: 15976525.
- Teesson M, Baillie A, Lynskey M, et al. Substance use, dependence and treatment seeking in the United States and Australia: a cross-national comparison. *Drug Alcohol Depend*. 2006 Feb 1;81(2):149-55. PMID: 16043307.
- Hasin DS, Grant BF. The co-occurrence of DSM-IV alcohol abuse in DSM-IV alcohol dependence: results of the National Epidemiologic Survey on Alcohol and Related Conditions on heterogeneity that differ by population subgroup. *Arch Gen Psychiatry*. 2004 Sep;61(9):891-6. PMID: 15351767.
- Mann K, Schafer DR, Langle G, et al. The long-term course of alcoholism, 5, 10 and 16 years after treatment. *Addiction*. 2005 Jun;100(6):797-805. PMID: 15918810.
- Norstrom T. Per capita alcohol consumption and all-cause mortality in Canada, 1950-98. *Addiction*. 2004 Oct;99(10):1274-8. PMID: 15369565.
- Rivara FP, Garrison MM, Ebel B, et al. Mortality attributable to harmful drinking in the United States, 2000. *J Stud Alcohol*. 2004 Jul;65(4):530-6. PMID: 15376828.
- Corrao G, Bagnardi V, Zambon A, et al. A meta-analysis of alcohol consumption and the risk of 15 diseases. *Prev Med*. 2004 May;38(5):613-9. PMID: 15066364.
- Cherpitel CJ, Ye Y. Alcohol-attributable fraction for injury in the U.S. general population: data from the 2005 National Alcohol Survey. *J Stud Alcohol Drugs*. 2008 Jul;69(4):535-8. PMID: 18612569.
- . Alcohol-attributable deaths and years of potential life lost--United States, 2001. *MMWR*

- Morb Mortal Wkly Rep. 2004 Sep 24;53(37):866-70. PMID: 15385917.
20. O'Brien CP, McLellan AT. Myths about the treatment of addiction. *Lancet*. 1996 Jan 27;347(8996):237-40. PMID: 8551886.
 21. Isaac M, Janca A, Sartorius N. ICD-10 symptom glossary for mental disorders. Geneva: Division of Mental Health, World Health Organization; 1994.
 22. Janca A, Ustun TB, van Drimmelen J, et al. ICD-10 symptom checklist for mental disorders, version 1.1. Geneva: Division of Mental Health, World Health Organization; 1994.
 23. American Psychiatric Association. Diagnostic and statistical manual of mental disorders, 4th ed. Text rev. Washington, DC: American Psychiatric Publishing, Inc.; 2000.
 24. U.S. Department of Veterans Affairs, U.S. Department of Defense. VA/DoD Clinical Practice Guideline for Management of Substance Use Disorders (SUD). 2009. http://www.healthquality.va.gov/sud/sud_full_601f.pdf.
 25. Pettinati HM, Weiss RD, Miller WR, et al. COMBINE Monograph Series, Volume 2. Medical Management Treatment Manual: A Clinical Research Guide for Medically Trained Clinicians Providing Pharmacotherapy as Part of the Treatment for Alcohol Dependence. DHHS Publication No. (NIH) 04-5289. Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; 2004.
 26. Center for Substance Abuse and Treatment. Incorporating alcohol pharmacotherapies into medical practice. (Treatment improvement protocol (TIP); no. 49). Substance Abuse and Mental Health Services Administration (SAMHSA) Rockville, MD: 2009.
 27. West SL, Garbutt JC, Carey TS, et al. Pharmacotherapy for Alcohol Dependence. Evidence report number 3. (Contract 290-97-0011 to Research Triangle Institute, University of North Carolina, Chapel Hill) AHCPR publication no. 99-E004. Rockville, MD: Agency for Health Care Policy and Research; January 1999. <http://www.ncbi.nlm.nih.gov/books/NBK32930/>.
 28. Garbutt JC, West SL, Carey TS, et al. Pharmacological treatment of alcohol dependence: a review of the evidence. *JAMA*. 1999 Apr 14;281(14):1318-25. PMID: 10208148.
 29. Methods Guide for Effectiveness and Comparative Effectiveness Reviews. Agency for Healthcare Research and Quality Rockville (MD). <http://www.ncbi.nlm.nih.gov/books/NBK47095/>.
 30. Viswanathan M, Ansari MT, Berkman ND, et al. Assessing the Risk of Bias of Individual Studies in Systematic Reviews of Health Care Interventions. . Agency for Healthcare Research and Quality Methods Guide for Comparative Effectiveness Reviews AHRQ Publication No. 12-EHC047-EF. Rockville, MD: Agency for Healthcare Research and Quality; March 2012. www.effectivehealthcare.ahrq.gov/.
 31. Sutton AJ, Abrams KR, Jones DR, et al. Methods for Meta-Analysis in Medical Research (Wiley Series in Probability and Statistics - Applied Probability and Statistics Section). London: Wiley; 2000.
 32. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002 Jun 15;21(11):1539-58. PMID: 12111919.
 33. Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ*. 2003 Sep 6;327(7414):557-60. PMID: 12958120.
 34. Owens DK, Lohr KN, Atkins D, et al. AHRQ series paper 5: grading the strength of a body of evidence when comparing medical interventions--Agency for Healthcare Research and Quality and the Effective Health-Care Program. *J Clin Epidemiol*. 2010 May;63(5):513-23. PMID: 19595577.
 35. Atkins DC, S.; Gartlehner, G.; et al. Chapter 6: Assessing the applicability of studies when comparing medical interventions. Agency for Healthcare Research and Quality AHRQ Publication No. 11-EHC019-EF. Rockville, MD: 2011.
 36. Lhuintre JP, Moore N, Tran G, et al. Acamprosate appears to decrease alcohol intake in weaned alcoholics. *Alcohol Alcohol*. 1990;25(6):613-22. PMID: 2085344.
 37. Fuller RK, Branchey L, Brightwell DR, et al. Disulfiram treatment of alcoholism. A Veterans Administration cooperative study. *JAMA*. 1986 Sep 19;256(11):1449-55. PMID: 3528541.
 38. Fuller RK, Roth HP. Disulfiram for the treatment of alcoholism. An evaluation in 128 men. *Ann*

- Intern Med. 1979 Jun;90(6):901-4. PMID: 389121.
39. Morgenstern J, Kuerbis AN, Chen AC, et al. A randomized clinical trial of naltrexone and behavioral therapy for problem drinking men who have sex with men. *J Consult Clin Psychol*. 2012;80(5):863-75. PMID: 22612306.
 40. LoCastro JS, Youngblood M, Cisler RA, et al. Alcohol treatment effects on secondary nondrinking outcomes and quality of life: the COMBINE study. *J Stud Alcohol Drugs*. 2009 Mar;70(2):186-96. PMID: 19261230.
 41. Pettinati HM, Gastfriend DR, Dong Q, et al. Effect of extended-release naltrexone (XR-NTX) on quality of life in alcohol-dependent patients. *Alcohol Clin Exp Res*. 2009 Feb;33(2):350-6. PMID: 19053979.
 42. O'Malley SS, Robin RW, Levenson AL, et al. Naltrexone alone and with sertraline for the treatment of alcohol dependence in Alaska natives and non-natives residing in rural settings: a randomized controlled trial. *Alcohol Clin Exp Res*. 2008 Jul;32(7):1271-83. PMID: 18482155.
 43. Anton RF, O'Malley SS, Ciraulo DA, et al. Combined pharmacotherapies and behavioral interventions for alcohol dependence: the COMBINE study: a randomized controlled trial. *JAMA*. 2006 May 3;295(17):2003-17. PMID: 16670409.
 44. Kiefer F, Jahn H, Tarnaske T, et al. Comparing and combining naltrexone and acamprosate in relapse prevention of alcoholism: a double-blind, placebo-controlled study. *Arch Gen Psychiatry*. 2003 Jan;60(1):92-9. PMID: 12511176.
 45. Morley KC, Teesson M, Reid SC, et al. Naltrexone versus acamprosate in the treatment of alcohol dependence: A multi-centre, randomized, double-blind, placebo-controlled trial. *Addiction*. 2006 Oct;101(10):1451-62. PMID: 16968347.
 46. Johnson BA, Rosenthal N, Capece JA, et al. Topiramate for treating alcohol dependence: a randomized controlled trial. *JAMA*. 2007 Oct 10;298(14):1641-51. PMID: 17925516.
 47. Johnson BA, Ait-Daoud N, Bowden CL, et al. Oral topiramate for treatment of alcohol dependence: a randomised controlled trial. *Lancet*. 2003 May 17;361(9370):1677-85. PMID: 12767733.
 48. Karhuvaara S, Simojoki K, Virta A, et al. Targeted nalmefene with simple medical management in the treatment of heavy drinkers: a randomized double-blind placebo-controlled multicenter study. *Alcohol Clin Exp Res*. 2007 Jul;31(7):1179-87. PMID: 17451401.
 49. Rehm J, Baliunas D, Borges GL, et al. The relation between different dimensions of alcohol consumption and burden of disease: an overview. *Addiction*. 2010 May;105(5):817-43. PMID: 20331573.
 50. Bondy SJ, Rehm J, Ashley MJ, et al. Low-risk drinking guidelines: the scientific evidence. *Can J Public Health*. 1999 Jul-Aug;90(4):264-70. PMID: 10489725.
 51. Shalala DE. 10th Special Report to the U.S. Congress on Alcohol and Health: Highlights From Current Research: From the Secretary of Health and Human Services. Washington, DC: U.S. Department of Health and Human Services; 2000. <http://pubs.niaaa.nih.gov/publications/10report/intro>. Accessed 8 June 2012.
 52. Monterosso JR, Flannery BA, Pettinati HM, et al. Predicting treatment response to naltrexone: the influence of craving and family history. *Am J Addict*. 2001 Summer;10(3):258-68. PMID: 11579624.
 53. O'Malley SS, Rounsaville BJ, Farren C, et al. Initial and maintenance naltrexone treatment for alcohol dependence using primary care vs specialty care: a nested sequence of 3 randomized trials. *Arch Intern Med*. 2003 Jul 28;163(14):1695-704. PMID: 12885685.
 54. Kiritze-Topor P, Huas D, Rosenzweig C, et al. A pragmatic trial of acamprosate in the treatment of alcohol dependence in primary care. *Alcohol Alcohol*. 2004 Nov-Dec;39(6):520-7. PMID: 15304381.
 55. O'Malley SS, O'Connor PG. Medications for unhealthy alcohol use: across the spectrum. *Alcohol Res Health*. 2011;33(4):300-12. PMID: 23580015.
 56. D'Amico EJ, Paddock SM, Burnam A, et al. Identification of and guidance for problem drinking by general medical providers: results from a national survey. *Med Care*. 2005 Mar;43(3):229-36. PMID: 15725979.
 57. Kranzler HR, Armeli S, Tennen H, et al. Targeted naltrexone for early problem drinkers. *J Clin Psychopharmacol*. 2003 Jun;23(3):294-304. PMID: 12826991.

58. Gual A, Leher P. Acamprosate during and after acute alcohol withdrawal: a double-blind placebo-controlled study in Spain. *Alcohol Alcohol*. 2001;36(5):413-8. PMID: CN-00367117.
59. Polich JM, Armor DJ, Braiker HB. Stability and change in drinking patterns. *The Course of Alcoholism: Four Years After Treatment*. New York, NY: John Wiley & Sons; 1981:159-200.
60. Kissin B, Charnoff SM, Rosenblatt SM. Drug and placebo responses in chronic alcoholics. *Psychiatr Res Rep Am Psychiatr Assoc*. 1968 Mar;24:44-60. PMID: 4889329.

Introduction

Background

Alcohol misuse, which includes the full spectrum from drinking above recommended limits (i.e., risky or hazardous drinking) to alcohol dependence,¹⁻³ is associated with numerous health and social problems, more than 85,000 deaths per year in the United States,^{4,5} and an estimated annual cost to society of more than \$220 billion.^{6,7} Alcohol misuse is estimated to be the third leading cause of preventable mortality in the United States, following tobacco use and being overweight.⁸ Definitions of the spectrum of alcohol misuse (i.e., unhealthy alcohol use¹) continue to evolve. For the purposes of this report, we use the definitions described in Table 1.

Table 1. Definitions of the spectrum of alcohol misuse

Term	Definition
Risky or hazardous use	Consumption of alcohol above recommended daily, weekly, or per-occasion amounts. ⁵ Consumption levels that increase the risk for health consequences.
Harmful use ^{9,10}	A pattern of drinking that is already causing damage to health. The damage may be either physical (e.g., liver damage from chronic drinking) or mental (e.g., depressive episodes secondary to drinking).
Alcohol abuse ¹¹	<p>A. A maladaptive pattern of alcohol use leading to clinically significant impairment or distress, as manifested by at least 1 of the following occurring within a 12-month period:</p> <ol style="list-style-type: none">(1) recurrent alcohol use resulting in a failure to fulfill major role obligations at work, school, or home (e.g., repeated absences or poor work performance related to alcohol use; alcohol-related absences, suspensions, or expulsions from school; neglect of children or household);(2) recurrent alcohol use in situations in which it is physically hazardous (e.g., driving an automobile or operating a machine when impaired);(3) recurrent alcohol-related legal problems (e.g., arrests for alcohol-related disorderly conduct); or(4) continued alcohol use despite having persistent or recurrent social or interpersonal problems caused or exacerbated by the effects of alcohol (e.g., arguments with spouse about consequences of intoxication, physical fights). <p>B. The symptoms have never met the criteria for alcohol dependence.</p>
Alcohol dependence ¹¹ (alcoholism, alcohol addiction)	<p>A maladaptive pattern of alcohol use, leading to clinically significant impairment or distress, as manifested by at least 3 of the following occurring at any time in the same 12-month period:</p> <ol style="list-style-type: none">(1) tolerance, as defined by either of the following:<ol style="list-style-type: none">(a) a need for markedly increased amounts of alcohol to achieve intoxication or desired effect; or(b) markedly diminished effect with continued use of the same amount of alcohol;(2) withdrawal, as manifested by either of the following:<ol style="list-style-type: none">(a) the characteristic withdrawal syndrome for alcohol; or(b) alcohol (or a closely related drug) is taken to relieve or avoid withdrawal symptoms;

Table 1. Definitions of the spectrum of alcohol misuse (continued)

Term	Definition
Alcohol dependence ¹¹ (alcoholism, alcohol addiction) (continued)	(3) alcohol is often taken in larger amounts or over a longer period than was intended; (4) there is a persistent desire or unsuccessful efforts to cut down or control alcohol use; (5) a great deal of time is spent in activities necessary to obtain alcohol, use alcohol, or recover from its effects; (6) important social, occupational, or recreational activities are given up or reduced because of alcohol use; or (7) alcohol use is continued despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by alcohol (e.g., continued drinking despite recognition that an ulcer was made worse by alcohol consumption).

Alcohol-use disorders include harmful use, alcohol abuse, and alcohol dependence;^{12,13} they are relatively common in developed countries.¹² Prevalence of alcohol use disorders is higher for men than for women, with estimates indicating a lifetime risk of more than 20 percent for men.^{12,14-16} Alcohol dependence has lifetime prevalence rates of about 17 percent for men and 8 percent for women.¹⁷

Alcohol use disorders cause substantial morbidity and mortality—that is, threefold to fourfold increased rates of early mortality.¹⁸⁻²⁰ They are associated with hypertension, heart disease, stroke, cancer, liver cirrhosis, amnesias, cognitive impairment, sleep problems, peripheral neuropathy, gastritis and gastric ulcers, pancreatitis, decreased bone density, anemia, depression, insomnia, anxiety, suicide, and fetal alcohol syndrome.^{12,21} In 2009, the number of alcoholic liver disease deaths was 15,183 and the number of alcohol-induced deaths, excluding accidents and homicides, was 24,518.⁸ Excessive alcohol consumption is also a major factor in injury and violence.²² Acute alcohol-related harm can be the result of fires, drowning, falls, homicide, suicide, motor vehicle crashes, child maltreatment, and pedestrian injuries.²³ In addition, alcohol use disorders can complicate the assessment and treatment of other medical and psychiatric problems.¹²

Alcohol use disorders often begin in the teens and 20s and fluctuate over time, with periods of abstinence (perhaps following a crisis), subsequent periods of sobriety followed by temporary controlled drinking, and then enhanced likelihood of increasing intake and problems.¹² Twenty to 30 percent of people with alcohol use disorders achieve long-term remission without any formal treatment.^{12,24,25}

Some studies indicate that less than 10 percent of those with alcohol use disorders are able to achieve long periods of nonproblematic drinking.²⁶⁻³⁰ Thus, the goal of treatment in the United States has traditionally been complete abstinence, because of the belief that it is unlikely that those with alcohol use disorders can return to controlled, healthy alcohol use. However, controlled drinking and harm reduction are often goals of treatment in parts of Europe.^{12,29}

Treatments for Alcohol-Use Disorders

Treatments for alcohol use disorders continue to evolve as research on the effectiveness of various treatments is published, and new treatments, including pharmacotherapy, are introduced and used more frequently. No single best approach has yet proven superior among the variety of

available treatment options. Some common treatments for alcohol use disorders include cognitive behavioral therapy, motivational enhancement therapy, 12-step programs (e.g., Alcoholics Anonymous), and pharmacotherapy. Treatment may be delivered via individual outpatient counseling, intensive outpatient programs using group or individual methods, alcoholism treatment centers, or other approaches.

Using complete abstinence as an outcome, from 15 to 35 percent of patients have been reported to achieve 1 year of sobriety following a variety of treatment approaches.³¹ Treatment approaches reviewed have included clinical trials of disulfiram, motivational enhancement therapy, cognitive behavioral therapy, and 12-step facilitation, as well as treatment as usual within alcoholism-treatment centers. Sobriety outcomes at 3 to 5 years or longer have been reported to be in a similar range.¹²

Over the past 15 to 20 years, awareness has grown that treatment may still be beneficial even if complete abstinence is not achieved. As a result, research has used other outcomes to measure the effectiveness of treatment, which can be subsumed under the concept of harm reduction.³² These measures include significant increases in abstinent days or decreases in heavy drinking episodes, improved physical health, and improvements in psychosocial functioning. Research using these outcomes can provide additional evidence for the effectiveness of treatment for alcohol dependence.

Pharmacological Interventions for Alcohol-Use Disorders

From the 1950s until the early 1990s, the pharmacotherapy for alcohol dependence consisted only of disulfiram, an aversive deterrent that produces significant physical symptoms, such as nausea or tachycardia, when alcohol is consumed. Since the 1990s, two oral medications (naltrexone and acamprosate) and one long-acting intramuscular formulation (of naltrexone) have been approved by the U.S. Food and Drug Administration (FDA) for alcohol dependence. These medications are recommended for people with alcohol dependence, generally after a successful withdrawal from alcohol, and together with psychological intervention.¹³ Table 2 describes the medications available in the United States that are FDA approved for treatment of alcohol use disorders, their mechanism of action, and dosing. The medications are usually prescribed for 3 to 12 months, though much longer courses of treatment are not uncommon in clinical practice. In clinical trials, the FDA-approved medications have shown evidence for efficacy in enhancing abstinence, reducing relapse to heavy drinking, and reducing overall drinking behavior.³³ Many additional medications have been used off-label or studied for treatment of alcohol use disorders. These include antidepressants, mood stabilizers, anticonvulsants, alpha-adrenergic blockers, antipsychotics, and anxiolytics.

Table 2. Medications that are FDA approved for treating adults with alcohol-use disorders

Generic Drug Name	Mechanism	Dosing
Acamprosate	Thought to modulate hyperactive glutamatergic NMDA receptors	666 mg 3 times per day
Disulfiram	Inhibits ALDH2, causing accumulation of acetaldehyde during alcohol consumption, which produces a variety of adverse effects such as nausea, dizziness, flushing, and changes in heart rate and blood pressure	250 to 500 mg per day
Naltrexone	Opioid antagonist; competitively binds to opioid receptors and blocks the effects of endogenous opioids such as β -endorphin	Oral: 50 to 100 mg per day Intramuscular injection: 380 mg per month

Abbreviations: ALDH2 = aldehyde dehydrogenase; FDA = U.S. Food and Drug Administration; mg = milligram; NMDA = N-methyl-D-aspartate.

Despite ongoing developments and advancements in treatment approaches, alcohol dependence represents one of the most undertreated disorders in the U.S. health care system; it is estimated that only 1 in 4 individuals with alcohol dependence receives treatment.¹⁷ Furthermore, of those patients who receive treatment, less than 1 in 10 receives medication as part of his or her treatment. Therefore, expanding awareness and access to this relatively new treatment modality has the potential to improve health outcomes and reduce the burden of this devastating illness that affects an estimated 8 million to 9 million U.S. citizens.

Existing Guidance

The Veterans Administration (VA), National Institute on Alcohol Abuse and Alcoholism (NIAAA), and Substance Abuse and Mental Health Services Administration (SAMHSA) all have guidelines addressing the use of pharmacotherapy for alcohol dependence.³⁴⁻³⁶ The VA guidelines recommend that oral naltrexone and/or acamprosate routinely be considered for patients with alcohol dependence (although acamprosate is currently a nonformulary medication for the VA), and that medications be offered in combination with addiction-focused counseling.

In 2011, the United Kingdom's National Institute for Clinical Excellence (NICE) released a set of clinical guidelines on the identification and treatment of people with alcohol dependence and harmful alcohol use.¹³ The guidelines include the following recommendations: (1) after a successful withdrawal for people with moderate or severe alcohol dependence, to consider offering acamprosate or oral naltrexone in combination with an individual psychological intervention (cognitive behavioral therapies, behavioral therapies, or social network and environment-based therapies) focused specifically on alcohol misuse; (2) to consider offering disulfiram in combination with a psychological intervention for people who have a goal of abstinence but for whom acamprosate and oral naltrexone are not suitable, or who prefer disulfiram and understand the relative risks of taking the drug; and (3) to have specialist and competent staff administer pharmacological interventions.

Scope and Key Questions

The use of medications for alcohol-use disorders is associated with uncertainty and variation across providers and settings. In recent years, many new trials of medications for alcohol-use disorders have been published. Since the 1999 AHRQ report on medications for alcohol dependence,^{37,38} there has been more than a 10-fold increase in the number of individuals studied in controlled clinical trials of naltrexone and acamprosate, and a series of well-conducted trials have been completed with other pharmacotherapeutic agents that are not FDA-approved for treating alcohol dependence. Other reasons for conducting a new review on this topic include the following: (1) to assess the comparative effectiveness of the FDA approved medications; (2) to determine whether any agents that are not FDA approved have evidence supporting their efficacy; (3) to evaluate the evidence on intramuscular naltrexone (Vivitrol®), a fairly recently approved medication; (4) to evaluate whether or not trials provide evidence of effectiveness in primary care settings; (5) to assess whether some medications are more or less effective for adults with certain genetic polymorphisms; and (6) to inform updates to clinical practice guidelines.

We approach each Key Question (KQ) by considering the relevant Populations, Interventions, Comparators, Outcomes, Timing, and Settings (PICOTS). Our report focuses on clinically relevant medications (those that are commonly used, those with sufficient literature for systematic review, and those of greatest interest to clinicians and to the developers of

guidelines). Our report is limited to people with alcohol-use disorders; it does *not* address people with risky or hazardous alcohol use (for whom medications are likely not an appropriate intervention).


The main objective of this report is to conduct a systematic review and meta-analysis of the comparative effectiveness and harms of medications for adults with alcohol-use disorders. In this review, we address the following KQs:

- KQ 1a: Which medications are efficacious for improving consumption outcomes for adults with alcohol-use disorders in outpatient settings?
- KQ 1b: How do medications for adults with alcohol-use disorders compare for improving consumption outcomes in outpatient settings?
- KQ 2a: Which medications are efficacious for improving health outcomes for adults with alcohol-use disorders in outpatient settings?
- KQ 2b: How do medications for adults with alcohol-use disorders compare for improving health outcomes in outpatient settings?
- KQ 3a: What adverse effects are associated with medications for adults with alcohol-use disorders in outpatient settings?
- KQ 3b: How do medications for adults with alcohol-use disorders compare for adverse effects in outpatient settings?
- KQ 4: Are medications for treating adults with alcohol-use disorders effective in primary care settings?
- KQ 5: Are any of the medications more or less effective than other medications for men or women, older adults, young adults, racial or ethnic minorities, smokers, or those with co-occurring disorders?
- KQ 6: Are any of the medications more or less effective for adults with certain genetic polymorphisms (e.g., of the mu-opioid receptor gene [OPRM1]) compared with adults without such polymorphisms?

Analytic Framework

We developed an analytic framework to guide the systematic review process (Figure 1).

Figure 1. Analytic framework for pharmacotherapy for adults with alcohol-use disorders in outpatient settings



KQ 1 assesses which medications are efficacious and how they compare with one another for improving alcohol consumption outcomes. KQ 2 examines which medications are efficacious and how they compare with one another for improving health outcomes. KQ 3 examines harms. KQ 4 focuses on evidence for primary care settings. KQ 5 assesses whether the medications are more or less effective compared with each other for a variety of subgroups. KQ 6 assesses whether any of the medications are more or less effective for adults with certain genetic polymorphisms than for adults without such polymorphisms.

Methods

The methods for this comparative effectiveness review (CER) follow the methods suggested in the Agency for Healthcare Research and Quality (AHRQ) *Methods Guide for Effectiveness and Comparative Effectiveness Reviews* (<http://www.effectivehealthcare.ahrq.gov/methodsguide.cfm>).

Topic Refinement and Review Protocol

This topic was nominated by a physician affiliated with the Substance Abuse and Mental Health Services Administration (SAMHSA), which works to improve the quality and availability of substance abuse prevention, alcohol and drug addiction treatment, and mental health services. During the topic development and refinement processes, we engaged in a public process to develop a draft and final protocol for the CER process. We generated an analytic framework, preliminary Key Questions (KQs), and preliminary inclusion/exclusion criteria in the form of PICOTS (populations, interventions, comparators, outcomes, timing, settings). The processes were guided by the information provided by the topic nominator, a scan of the literature, methods and content experts, and Key Informants. We worked with six Key Informants during the topic refinement, all of whom subsequently served on the Technical Expert Panel (TEP) for this report. The TEP consisted of eight individuals in total. Key Informants and TEP members participated in conference calls and discussions through email to review the scope, analytic framework, KQs, and PICOTS; provided input on the information and categories included in evidence tables; and provided input on the data analysis plan.

The KQs were posted for public comment on AHRQ's Effective Health Care Web site from September 20 to October 18, 2012; we put them into final form after review of the comments and discussion with the TEP. The only comments we received were attempts to provide answers to the questions rather than to provide input about the draft scope, KQs, PICOTS, or analytic framework. Therefore, no changes were made based on public review. We then drafted a protocol for this CER and refined the protocol in consultation with AHRQ and the TEP before it was posted on the Effective Health Care Web site on April 29, 2013.

Literature Search Strategy

Search Strategy

To identify articles relevant to each KQ, we searched PubMed®, the Cochrane Library, PsycINFO®, CINAHL®, and EMBASE®. The full search strategy is presented in Appendix A. We used either Medical Subject Headings (MeSH) or major headings as search terms when available or key words when appropriate, focusing on terms to describe the relevant populations and interventions of interest. We reviewed our search strategy with the TEP and incorporated their input. Searches were run by an experienced information scientist serving as the Evidence-based Practice Center (EPC) librarian and were peer-reviewed by another information scientist/EPC librarian.

We limited the electronic searches to English-language, adult (18 and older), and human-only studies. Sources were searched from January 1, 1970, to February 5, 2013. This search date was selected based on the earliest publications found during the topic refinement process, the earliest study found in previous systematic reviews (which was from 1974), and expert opinion.

We manually searched reference lists of pertinent reviews, trials, and background articles on this topic to look for any relevant citations that our searches might have missed. We imported all citations into an EndNote® X4 (Thomson Reuters, New York, NY) electronic database.

We also searched for unpublished studies relevant to this review using ClinicalTrials.gov, the World Health Organization International Clinical Trials Registry Platform, and the Web site for the U.S. Food and Drug Administration (FDA). In addition, AHRQ's Scientific Resource Center requested scientific information packets from relevant pharmaceutical companies, asking for any unpublished studies or data relevant to this CER. Scientific information packets allow pharmaceutical companies to provide the EPC with published or unpublished data that they believe should be considered for the review. Any additional studies identified from the packets will be included in the post-peer/public review report.

In cases in which relevant information was unclear or not reported, we contacted authors to get additional or unpublished information. When successful, this information was included in the findings.

We will conduct an update of our literature searches (of the same databases searched initially) concurrent with the peer review process. Any literature suggested by peer reviewers or the public will be investigated and, if appropriate, incorporated into the final review. We will determine appropriateness for inclusion in the review by the same methods described in this chapter.

Inclusion and Exclusion Criteria

We developed eligibility (inclusion and exclusion) criteria with respect to PICOTS and study designs and durations for each KQ (Table 3).

Table 3. Eligibility criteria

Category	Inclusion	Exclusion
Population	Adults (age 18 years or older) with alcohol-use disorders (as defined in the Introduction) For KQ 5, co-occurring disorders include other mental health or substance use disorders (e.g., depression, cocaine addiction) and acute or chronic medical conditions (e.g., cirrhosis)	Children and adolescents under 18
Interventions	Medications approved by FDA for treating alcohol dependence (acamprosate, disulfiram, naltrexone) and the following medications, which have been used off-label or are under investigation: amitriptyline, aripiprazole, atomoxetine, baclofen, buspirone, citalopram, desipramine, escitalopram, fluoxetine, fluvoxamine, gabapentin, imipramine, nalmefene, olanzapine, ondansetron, paroxetine, prazosin, quetiapine, sertraline, topiramate, valproate, varenicline, viloxazine	Pharmacotherapy for alcohol withdrawal; any drugs not listed; combinations of medications (e.g., studies randomizing subjects to naltrexone plus ondansetron vs. placebo)
Comparators	For KQs 1 through 5, studies must compare one of the medications listed above with placebo or another medication For KQ 6, studies must compare people who have a genetic polymorphism with people who do not have the polymorphism	No comparison; nonconcordant historical controls

Table 3. Eligibility criteria (continued)

Category	Inclusion	Exclusion
Outcomes	Consumption outcomes: return to any drinking, return to heavy drinking, drinking days, heavy drinking days, drinks per drinking day, time to lapse or relapse Health outcomes: accidents, injuries, quality of life, function, mortality Adverse effects of intervention(s): withdrawals due to adverse events, nausea/vomiting, diarrhea, anorexia, palpitations, headache, dizziness, cognitive dysfunction, taste abnormalities, paresthesias (numbness, tingling), metabolic acidosis, glaucoma, vision changes, suicidal ideation, insomnia, anxiety, rash	Craving; cue reactivity
Timing/length of followup	At least 12 weeks of followup from the time of medication initiation	Less than 12 weeks
Settings	Outpatient health care (i.e., nonlaboratory) settings, including studies that begin in or recruit subjects from inpatient settings but then follow and assess subjects receiving pharmacotherapy as outpatients KQ 4 applies to primary care settings only (i.e., internal medicine, family medicine, pediatrics, obstetrics/gynecology, or college and university health clinics)	All other settings; laboratory settings; inpatient settings (if most or all of the study followed inpatients)
Publication language	English	All other languages
Admissible evidence (study design and other criteria)	Original research; eligible study designs include the following: <ul style="list-style-type: none"> • For KQs 1, 2, and 4, double-blind RCTs and recent systematic reviews were eligible • For KQ 2b (head-to-head studies reporting health outcomes), prospective cohort studies were also eligible • For KQ 3 (harms), double-blind RCTs and recent systematic reviews that compare medication with placebo or with another medication were eligible. The following designs were also eligible if they compared 2 or more drugs of interest: nonrandomized controlled trials, open-label trials, secondary analyses or subgroup analyses from trials, prospective cohort studies, and case-control studies • For KQ 5 (subgroups), double-blind RCTs, recent systematic reviews, nonrandomized controlled trials, open-label trials, secondary analyses or subgroup analyses from trials, prospective cohort studies, and case-control studies were eligible, as long as the studies compared 2 or more drugs • For KQ 6, double-blind RCTs, secondary analyses or subgroup analyses from trials, and prospective cohort studies comparing people with genetic polymorphisms with people without such polymorphisms were eligible 	Case series Case reports Nonsystematic reviews Systematic reviews with searches that ended prior to 2007 Systematic reviews that had been updated Editorials Letters to the editor Studies with historical, rather than concurrent, control groups

Abbreviations: FDA = U.S. Food and Drug Administration; KQ = Key Question; RCT = randomized controlled trial.

Study Selection

Two trained members of the research team independently reviewed each title and abstract (identified through searches) against our eligibility criteria. Studies marked for possible inclusion by either reviewer underwent a full-text review. For titles or abstracts that lacked adequate information to determine eligibility, we retrieved and reviewed the full text. Two trained members of the research team independently reviewed each full-text article and determined eligibility based on the criteria described above. If the reviewers disagreed, they resolved

conflicts by discussion and consensus or by consulting a third, senior member of the team. We recorded the principal reason that each excluded full-text publication did not satisfy the eligibility criteria (Appendix B). All results in both review stages were tracked in an EndNote® database.

Data Extraction

For studies that met our inclusion criteria, we extracted important information into evidence tables. We designed, pilot-tested, and used structured data extraction forms to gather pertinent information from each article; this included characteristics of study populations, settings, interventions, comparators, study designs, methods, and results. Trained reviewers extracted the relevant data from each included article. All data abstractions were reviewed for completeness and accuracy by a second member of the team. We recorded intention-to-treat (ITT) results if available. All data abstraction was performed using Microsoft Word® or Excel® software.

Risk-of-Bias Assessment of Individual Studies

To assess the risk of bias (internal validity) of studies for major outcomes of interest, we used predefined criteria based on guidance from the AHRQ *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*.³⁹ We assessed selection bias, confounding, performance bias, detection bias, and attrition bias; we included questions about adequacy of randomization, allocation concealment, similarity of groups at baseline, blinding, attrition, whether ITT analysis was used, methods of handling missing data, and fidelity. We rated the studies as low, medium, high, or unclear risk of bias.⁴⁰

In general terms, studies categorized as low risk of bias imply high confidence that the results represent the true treatment effects. Studies with medium risk of bias are susceptible to some risk of bias but probably not enough to invalidate the results. Studies with a medium risk of bias did not meet all criteria required for low risk of bias. These studies had some flaws in design or execution (e.g., inadequate description of methods of randomization and allocation concealment) but they provided enough information to allow readers to determine that the flaws did not likely cause major bias. Missing information often led to ratings of medium as opposed to low risk of bias. Studies assessed as high risk of bias have significant flaws stemming from serious errors in design, conduct, or analysis that may invalidate the results (e.g., high overall or differential attrition without appropriate handling of missing data).

Two independent reviewers assessed the risk of bias for each study; one of the two reviewers was always an experienced EPC investigator. Disagreements between the two reviewers were resolved by discussion and consensus or by a third member of the team. We omitted studies deemed high risk of bias by two reviewers from our main data synthesis and main analyses; we included them only in sensitivity analyses. Appendix C details the criteria used for evaluating the risk of bias of all included studies and explains the rationale for high risk of bias ratings.

Data Synthesis

We conducted quantitative synthesis using meta-analyses of outcomes reported by multiple studies that were sufficiently homogeneous to justify combining their results. To determine whether meta-analyses were appropriate, we assessed the clinical and methodological heterogeneity of the studies under consideration following established guidance.⁴¹ We did this by qualitatively assessing the PICOTS of the included studies and looking for similarities and

differences. When quantitative synthesis was not appropriate (e.g., because of clinical heterogeneity, insufficient numbers of similar studies, or insufficiency or variation in outcome reporting), we synthesized the data qualitatively.

We used random-effects models to estimate pooled effects.⁴² For continuous outcomes (e.g., scales for symptom reduction), we used weighted mean differences (WMDs). For binary outcomes we calculated risk differences (RDs) between groups. We did not include studies rated as high or unclear risk of bias in our main analyses, but did include them in sensitivity analyses. For alcohol consumption outcomes, if studies reported consumption in grams, we used a conversion factor of 13.7 grams as equivalent to a standard drink.⁴³ All quantitative analyses were conducted using Stata[®] version 11.1 (StataCorp LP, College Station, TX).

We calculated the chi-squared statistic and the I^2 statistic to assess statistical heterogeneity in effects between studies.^{44,45} An I^2 from 0 to 40 percent might not be important, 30 percent to 60 percent may represent moderate heterogeneity, 50 percent to 90 percent may represent substantial heterogeneity, and ≥ 75 percent represents considerable heterogeneity.⁴⁶ The importance of the observed value of I^2 depends on the magnitude and direction of effects and on the strength of evidence (SOE) for heterogeneity (e.g., p value from the chi-squared test, or a confidence interval for I^2). Whenever we include a meta-analysis with considerable statistical heterogeneity in this report, we attempt to provide an explanation for the heterogeneity, considering the magnitude and direction of effects.⁴⁶ We examined potential sources of heterogeneity by stratifying analyses by patient population or setting (i.e., U.S.-based trials compared with others), variation in interventions (i.e., dose and route of delivery), and duration of treatment.

Strength of the Body of Evidence

We graded SOE based on the guidance established for the Evidence-based Practice Center program.⁴⁷

Developed to grade the overall strength of a body of evidence, this approach incorporates four key domains: risk of bias (includes study design and aggregate quality), consistency, directness, and precision of the evidence. It also considers optional domains, such as a dose-response association, plausible confounding that would decrease the observed effect, strength of association (magnitude of effect), and publication bias. Table 4 defines the grades of evidence that we assigned.

Table 4. Definitions of the grades of overall strength of evidence

Grade	Definition
High	High confidence that the evidence reflects the true effect. Further research is very unlikely to change our confidence in the estimate of effect.
Moderate	Moderate confidence that the evidence reflects the true effect. Further research may change our confidence in the estimate of the effect and may change the estimate.
Low	Low confidence that the evidence reflects the true effect. Further research is likely to change our confidence in the estimate of the effect and is likely to change the estimate.
Insufficient	Evidence either is unavailable or does not permit estimation of an effect.

Source: Owens et al., 2010⁴⁷

Two reviewers assessed each domain for each key outcome and determined an overall SOE grade based on domain ratings. We generally required consistent, direct, precise evidence from studies with aggregate low risk of bias to give high SOE grades. An unfavorable assessment for any one of the four key domains (i.e., inconsistency, indirectness, imprecision, or medium aggregate risk of bias) typically resulted in downgrading to moderate SOE. Two unfavorable

assessments typically resulted in downgrading to low SOE. We allowed reviewers to include the optional domains listed above (e.g., dose-response association, publication bias) if relevant, and to upgrade or downgrade the SOE for those domains if appropriate. In the event of disagreements on the domain or overall grade, they resolved differences by consensus discussion or by consulting with a third, experienced EPC investigator.

We graded the SOE for the following outcomes: return to any drinking, return to heavy drinking, drinking days, heavy drinking days, drinks per drinking day, accidents, injuries, quality of life or function, mortality, and adverse events. Appendix D includes tables showing our assessments for each domain and the resulting SOE grades for each outcome, organized by KQ and intervention/comparison pair.

Applicability

We assessed applicability of the evidence following guidance from the *Methods Guide for Comparative Effectiveness Reviews*.⁴⁸ We used the PICOTS framework to explore factors that affect applicability. Some factors identified a priori that may limit the applicability of evidence include the age, sex, and race or ethnicity of enrolled populations; smoking status of enrolled populations; co-occurring disorders of enrolled populations; setting; type of provider prescribing the treatment; and source of subject recruitment. Regarding the source of subject recruitment, studies of subjects recruited via advertisements may enroll people that have less severe disorders, and may be less applicable to patients with more severe forms of alcohol-use disorders.

Peer Review and Public Commentary


An external peer review will be performed on this report. We will compile all comments and address each one individually, revising the text as appropriate. AHRQ will also provide review from its own staff. In addition, the Scientific Resource Center will place the draft report on the AHRQ Web site (www.effectivehealthcare.ahrq.gov/) for public review.

Results

Results of Literature Searches

Results of our searches appear in Figure 2. We included 157 published articles reporting on 130 studies. Of the included studies, 114 were randomized controlled trials. Additional details describing the included studies are provided in the relevant sections of this results chapter.

Figure 2. Disposition of articles



Key Question 1. Efficacy and Comparative Effectiveness for Improving Consumption Outcomes

For this Key Question (KQ), we describe the characteristics of included trials and then results for alcohol consumption outcomes (return to any drinking, return to heavy drinking, drinking days, heavy drinking days, drinks per drinking day) for medications for which we included multiple trials. For medications with just 1 eligible trial, we graded the strength of evidence (SOE) as insufficient (because evidence was imprecise, unknown consistency, and medium or high risk of bias); information on the characteristics and results for medications with just 1 eligible trial is provided in Appendix E.

Throughout this KQ, we include headers and sections only for consumption outcomes with sufficient data for synthesis. Negative effect sizes favor medication over placebo. Positive effect sizes favor placebo. We describe the results of sensitivity analyses that included studies rated as high or unclear risk of bias only if they changed the effect size significantly. Results of all such sensitivity analyses are provided in Appendix F.

Detailed Synthesis: Placebo-Controlled Trials of FDA-Approved Medications for Treating Alcohol Dependence

Acamprosate

Characteristics of Trials

Table 5 summarizes characteristics of the 20 trials meeting our inclusion criteria. The majority were parallel two-arm trials comparing acamprosate with placebo. Doses ranged from 1,000 to 3,000 mg per day; 1,998 mg per day (divided into 3 doses) was the most frequently used dose. Duration of treatment ranged from 12 to 52 weeks; most (16 trials) treated subjects for 12 to 26 weeks; 4 trials treated subjects for longer periods, 48 to 52 weeks. Followup to 1 year or longer was available for 8 trials.

The majority were conducted in Europe (15 trials); 3 were conducted in the United States, 1 in Brazil, and 1 in Australia. Recruitment methods varied, with trials typically identifying patients through treatment programs (e.g., inpatient detoxification, outpatient treatment), advertisements, referrals, or some combination of those.

Mean age was very similar across trials, usually in the early to mid-40s. All subjects met criteria for alcohol dependence in 19 trials; 1 trial did not report the proportion with alcohol dependence, but most subjects likely had alcohol dependence.⁴⁹ Most studies did not report information on race; 1 trial reported enrolling a majority (65 percent) of nonwhite subjects.⁵⁰ Most trials enrolled between 11 and 36 percent females; 1 trial enrolled all males,⁵¹ and 1 did not report information on sex.⁵² Just 4 trials reported information on smoking history at baseline; those trials had 46 to 81 percent smokers enrolled.^{49,53-55}

The majority of trials either did not report information about how many subjects had co-occurring psychiatric conditions or excluded subjects with other psychiatric disorders; 1 trial enrolled subjects with alcohol dependence and schizophrenia spectrum disorders.⁵⁰ Trials often included or encouraged psychological or psychosocial co-interventions.

Table 5. Characteristics of included double-blind randomized placebo-controlled trials of acamprostate

Author, Year Trial Name	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Country	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Anton, 2006 ⁵³ Donovan, 2008 ⁵⁶ COMBINE	ACA 3,000 + CBI + MM (151) ACA 3,000 + MM (152) NTX 100 + CBI + MM (155) NTX 100 + MM (154) Placebo + CBI + MM (156) Placebo + MM (153) ^a	16 (68)	U.S.	11 U.S. academic sites	Ads, community resources, clinical referrals at 11 academic sites	44	23	31	NR	As randomized; Community support group participation (like AA) encouraged	Low
Baltieri, 2004 ⁵¹	ACA 1,998 (40) Placebo (35)	12 (24)	Brazil	Outpatient	Patients seeking treatment at an outpatient clinic for treatment of drug dependence	18-60	NR	0	0	AA encouraged	Medium
Besson, 1998 ⁵⁷	ACA 1,300 to 1,998 (55) Placebo (55)	52 (108)	Switzerland	Outpatient; 3 psychiatric treatment centers	From inpatient treatment unit	42	NR	20	0	Routine counseling 100% Voluntary disulfiram 22-24%	Medium
Chick, 2000 ⁵⁸	ACA 1,998 (289) Placebo (292)	24	U.K.	Outpatient	Recruited from treatment programs	43	NR	16	0	Usual psychosocial outpatient treatment program	Medium
Geerlings, 1997 ⁵⁹	ACA 1,332 to 1,998 (128) Placebo (134)	26 (52)	Belgium, the Netherlands, and Luxembourg	Outpatient substance abuse treatment centers	Recruited from detoxification patients in same centers	40-42	NR	24	NR	ACA: benzodiazepines 5% Placebo: benzodiazepines 6%	Medium
Gual, 2001 ⁶⁰	ACA 1,998 (148) Placebo (148)	26	Spain	Outpatient; multicenter; hospitals	NR	41	NR	20 to 21	NR	NR	Medium

Table 5. Characteristics of included double-blind randomized placebo-controlled trials of acamprosate (continued)

Author, Year Trial Name	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Country	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Kiefer, 2003 ⁶¹ Kiefer, 2004 ⁶² Kiefer, 2005 ⁶³	ACA 1,998 (40) NTX 50 (40) Placebo (40) ACA 1,998 + NTX 50 (40)	12	Germany	1 site, Hamburg outpatient	Inpatient withdrawal treatment	46	NR	26	0	Group therapy	Low
Lhuintre, 1985 ⁶⁴	ACA 1,000 to 2,250 (42) Placebo (43)	13	France	Outpatient; methadone maintenance clinics	Recruited as inpatients within 48 hours of admission	40 to 43	NR	11	NR	Meprobamate 100% for first month	High
Lhuintre, 1990 ⁶⁵	ACA 1,332 (279) Placebo (290)	12	France	Outpatient; multicenter	Recruited within 48 hours of hospitalization for alcohol withdrawal	42 to 43	NR	18	NR	Psychotherapy allowed	Unclear
Mason, 2006 ⁵⁴	ACA 2,000 (258) ACA 3,000 (83) Placebo (260)	24 (32)	U.S.	21 outpatient clinics ^b	Primarily by newspaper ads	44 to 45	14 to 15	29 to 36	NR	Brief abstinence-oriented protocol-specific counseling and self-help materials 100%	Low
Morley, 2006 ⁴⁹ Morley, 2010 ⁶⁶	ACA 1,998 (55) NTX 50 (53) Placebo (61)	12	Australia	3 treatment centers with "medical care typically available at hospital based drug and alcohol treatment services"	Patients who had attended an inpatient detoxification program, outpatient treatment or followup or who responded to live or print ads	45	NR	30	Severe concurrent illness (psychiatric or other) –NOS 3	All offered 4-6 sessions of manualized compliance therapy Up-take / attendance NR	Low
Paille, 1995 ⁶⁷	ACA 1.3 g (188) ACA 2 g (173) Placebo (177)	52 (78)	France	NR ^c	Referral from alcohol specialist centers	43	NR	20	NR	Supportive psychotherapy 100% Hypnotics 6 to 7% Anxiolytics 8 to 12% Antidepressants 8 to 9%	Medium

Table 5. Characteristics of included double-blind randomized placebo-controlled trials of acamprosate (continued)

Author, Year Trial Name	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Country	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Pelc, 1996 ⁶⁸ , Pelc, 1992 ⁶⁹	ACA 1,332 to 1,998 (55) Placebo (47)	26	Belgium	Outpatient; multicenter	Post-inpatient detoxification	43	NR	31	NR	Supportive psychotherapy 100%	High
Pelc, 1997 ⁵²	ACA 1,332 (63) ACA 1,998 (63) Placebo (62)	13	Belgium, France	Outpatient; after inpatient detoxification	Inpatient referral	NR	NR	NR	NR	Counseling, social support when needed 100%	Medium
Poldrugo, 1997 ⁷⁰	ACA 1,332 to 1,998 (122) Placebo (124)	26 (52)	Italy	Inpatient for 1-2 weeks then outpatient; multicenter community based alcohol rehabilitation program	From acute inpatient withdrawal treatment	43 to 45	NR	23 to 31	0	Community-based rehabilitation program with group sessions, alcohol education, community meetings 100%	Medium
Ralevski, 2011 ⁵⁰ , Ralevski, 2001 ⁷¹	ACA 1,998 (12) Placebo (11)	12	U.S.	Outpatient; university and VA health centers	From community and through referrals from treatment facilities at a university and a VA facility	51	65	17	Schizophrenia spectrum disorders 100	Weekly skills training that incorporated CB drug relapse prevention strategies 100%	High
Sass, 1996 ⁷²	ACA 1,332 to 1,998 (136) Placebo (136)	48 (96)	Germany	Psychiatric outpatient	Outpatient referral	41 to 42	NR	22	NR	Counseling / psychotherapy 100%	Medium
Tempesta, 2000 ⁷³	ACA 1,998 (164) Placebo (166)	26 (39)	Italy	Outpatient	Recruited from outpatient internal medicine, neurology and addiction treatment programs	46	NR	17	0	Medical and behavioral counseling	Medium
Whitworth, 1996 ⁷⁴	ACA 1,332 or 1,998 (224) Placebo (224)	52 (104)	Austria	Outpatient specialty	Inpatient recruitment	42	NR	21	NR	NR	Medium

Table 5. Characteristics of included double-blind randomized placebo-controlled trials of acamprosate (continued)

Author, Year Trial Name	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Co- occurring Condition	Co-intervention	Risk of Bias
Wolwer, 2011 ⁷⁵	ACA 1,998 + IBT (124) ACA 1,998 + TAU (122) ^d Placebo + IBT (125)	24 (52)	Germany	Outpatient; 4 university hospitals 1 non-academic clinic	Recruited after inpatient detoxification	46	NR	29	NR	NR	Medium

^a Three additional treatment arms were included in COMBINE but were not relevant to our Key Questions: ACA + NTX + CBI + MM, ACA + NTX + MM, and CBI only (no pills).

^b Clinics were affiliated with academic medical centers and had investigators experienced in alcoholism treatment.

^c The article was not explicit about the setting, but patients received psychotherapy and psychiatric medication management suggesting a psychiatric outpatient setting.

^d Treatment as usual, seen once per week in an individual setting; MI techniques allowed.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: AA = Alcoholics Anonymous; ACA = acamprosate; CB = cognitive behavioral; CBI = combined behavioral intervention; IBT = integrative behavior therapy; mg = milligram; MM = medical management; N = number; NOS = not otherwise specified; NR = not reported; NTX = naltrexone; TAU = treatment as usual; U.K. = United Kingdom; U.S. = United States; VA = Veterans Affairs.

Return to Any Drinking

Eighteen of the 20 trials reported sufficient data for meta-analysis. All but 1 study⁵⁴ had point estimates trending in favor of acamprosate. Our meta-analysis of low and medium risk of bias trials found that 10 percent fewer subjects treated with acamprosate returned to any drinking than with placebo (risk difference [RD], -0.10; 95% CI, -0.15 to -0.05). Statistical heterogeneity was considerable (I^2 80.2 percent).

Differences in country and duration of treatment seem to explain much of the heterogeneity. The only 2 U.S.-based trials contributing data found no difference between acamprosate and placebo.^{53,54} Stratifying our meta-analysis by U.S. and non-U.S. studies found no difference for the 2 U.S.-based trials (RD, 0.02; 95% CI, -0.05 to 0.08), but found 12 percent fewer subjects treated with acamprosate returned to any drinking than with placebo for trials conducted in other countries (RD, -0.12; 95% CI, -0.16 to -0.08), and statistical heterogeneity decreased to the moderate range. Stratifying by duration of treatment (which corresponds to timing of outcome assessment used in analyses) found low heterogeneity among studies treating patients for 48 to 52 weeks, with an 11 percent absolute reduction in return to any drinking (RD, -0.11; 95% CI, -0.16 to -0.06; 4 trials).

Return to Heavy Drinking

Our meta-analysis found no significant difference between acamprosate and placebo (RD, -0.01; 95% CI, -0.05 to 0.03; I^2 0 percent; 6 trials).

Drinking Days

Patients treated with acamprosate had 9.4 percent fewer drinking days than those treated with placebo (weighted mean difference [WMD], -9.4; 95% CI, -13.8 to -5.0; 12 trials).

The only 2 U.S.-based trials contributing data found no difference between acamprosate and placebo.^{53,54} Stratifying our meta-analysis by U.S. and non-U.S. studies found no difference for the 2 U.S.-based trials (WMD, -2.7; 95% CI, -8.3 to 3.0), but found that patients treated with acamprosate had 11.2 percent fewer drinking days than those treated with placebo for trials conducted in other countries (WMD, -11.2; 95% CI, -15.8 to -6.6). Stratifying by duration of treatment (which corresponds to timing of outcome assessment used in our analyses) found that patients treated with acamprosate had 12.2 percent fewer drinking days than those treated with placebo over 48 to 52 weeks (WMD, -12.2; 95% CI, -16.4 to -8.1; I^2 0 percent).

Drinks per Drinking Day

Just 1 trial rated as low or medium risk of bias reported data. It found no statistically significant difference between acamprosate and placebo (WMD, 0.40; 95% CI, -1.81 to 2.61).⁴⁹

Disulfiram

Characteristics of Trials

Table 6 summarizes characteristics of the 4 trials meeting our inclusion criteria. All 4 were conducted in Veterans Administration Medical Centers. Three compared disulfiram with placebo or riboflavin (which was intended as placebo); 1 compared disulfiram with naltrexone, placebo, and the combination of naltrexone and disulfiram.⁷⁶ Doses for the intended active disulfiram arms were the same (250 mg per day) in all 4 trials. Two of the 4 trials were rated as high risk of bias, either primarily for high risk of attrition bias and inadequate handling of missing data,⁷⁷ or primarily for high risk of ascertainment bias⁷⁶; see Appendix C for details.

Duration of treatment ranged from 12 to 52 weeks. Three of the 4 trials followed subjects for 9 to 12 months. All 4 were conducted in the United States. Mean age was very similar across trials, ranging from 39 to 47 years. All subjects likely met criteria for alcohol dependence. Very few female subjects were enrolled (0 to 3 percent in the 3 trials reporting). None of the trials reported information on smoking history at baseline. One trial enrolled subjects with alcoholism who were also in methadone maintenance programs.⁷⁷ Another enrolled subjects with co-occurring psychiatric disorders.⁷⁶ Neither of the trials rated as medium risk of bias reported information on how many subjects had co-occurring psychiatric conditions.

Return to Any Drinking

Three of the 4 trials reported data. Our meta-analysis found no statistically significant difference between disulfiram 250 mg per day and disulfiram 1 mg per day or placebo, both without (RD, 0.04; 95% CI, -0.03 to 0.11) and with inclusion of the studies rated as high risk of bias (RD, -0.00; 95% CI, -0.10 to 0.09). Both medium risk of bias studies found point estimates favoring placebo/disulfiram 1 mg, but differences between groups were not statistically significant.

Our meta-analysis found no statistically significant difference between disulfiram 250 mg per day and riboflavin (i.e., no disulfiram) (RD, -0.04; 95% CI, -0.11 to 0.03). Both medium risk of bias studies found point estimates favoring disulfiram 250 mg per day, but differences between groups were not statistically significant.

The largest trial (N=605)⁷⁸ reported a significant relationship between adherence and complete abstinence in all groups (disulfiram 250 mg, disulfiram 1 mg, and no disulfiram/riboflavin). The other trial assessed as medium risk of bias similarly reported that complete abstinence correlated significantly with adherence.⁷⁹

Drinking Days

Both medium risk of bias trials reported some information about the percentage of drinking days. The smaller trial (N=128) reported no statistically significant differences among the three groups in percentage of drinking days (31 percent versus 32 percent versus 37 percent, for disulfiram 500/250, disulfiram 1, and riboflavin, respectively, p NR). The larger trial (N=605) reported this outcome only for the subset of subjects who drank and had a complete set of assessment interviews (N=162). It found that patients among this subset treated with disulfiram reported fewer drinking days than those given disulfiram 1 mg or those given riboflavin (49 percent versus 75.4 percent versus 86.5 percent, respectively, p=0.05).

Table 6. Characteristics of included double-blind randomized placebo-controlled trials of disulfiram

Author, Year Trial name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Female	% With Additional Condition	Co-intervention	Risk of Bias
Fuller, 1979 ⁷⁹	DIS 250 (43) DIS 1 (43) RIB 50 (42)	52	U.S.	Outpatient; VA hospital	Patients presenting to VA hospital requesting treatment for alcoholism or patients admitted for alcohol-related illness	43	61	0	NR	Counseling (unspecified) 100%	Medium
Fuller, 1986 ⁷⁸	DIS 250 (202) DIS 1 (204) RIB 50 (199)	52	U.S.	Outpatient; 9 VAMCs	Screened as inpatients in 7 centers and outpatients at 2	41 to 42	47	0	NR	Counseling (loosely defined) % NR	Medium
Ling, 1983 ⁷⁷	DIS 250 (41) Placebo (41)	37	U.S.	Outpatient; VA	Unclear	39	NR	NR	Heroin use 80 Marijuana use 36 Other drug use 67 Depression 83 Moderate to high depression 50	Methadone 100%	High
Petrakis, 2005 ⁷⁶ Ralevski, 2007 ⁸⁰ Petrakis, 2007 ⁸¹ Petrakis, 2006 ⁸² VA MIRECC	DIS 250 (66) NTX 50 (59) Placebo (64) NTX 50 + DIS 250 (65)	12	U.S.	Outpatient; VA	Recruited as outpatients or ad	47	26	3	Axis I disorder 100	Psychiatric treatment as usual 100%	High for DIS vs. placebo

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: DIS = disulfiram; mg = milligram; MIRECC = Mental Illness Research, Education and Clinical Center; N = number; NR = not reported; NTX = naltrexone; RIB = riboflavin; U.S. = United States; VA = Veterans Affairs; VAMC = Veterans Administration Medical Center.

Naltrexone

Characteristics of Trials

Table 7 summarizes characteristics of the 42 trials meeting our inclusion criteria. Less than half were parallel two-arm trials comparing naltrexone with placebo; most had three or more study arms. Three trials evaluated long-acting, injectable naltrexone, at doses from 150 to 400 mg per day.⁸³⁻⁸⁵ The rest administered oral naltrexone—31 trials used a dose of 50 mg per day, 6 used 100 mg per day,^{53,86-90} 1 used 150 mg per day,⁹¹ and 1 used 100 mg on Mondays and Wednesdays and 150 mg on Fridays (weekly average of 50 mg per day).⁹² Duration of treatment ranged from 12 to 52 weeks; most (36 trials) treated subjects for 12 to 17 weeks; 6 trials included treatment with naltrexone for longer periods—24 to 52 weeks.^{83,88,93-96} Two of the latter groups included comparisons of different treatment durations for 50 mg per day, either comparing 12 versus 24 weeks⁹⁶ or comparing 12 versus 52 weeks.⁹⁵

The majority were conducted in the United States only (27 trials); 6 were conducted in Europe, 3 in Australia, 2 in Brazil, 1 multinational (United States, France, and the Netherlands), and 1 each in Singapore, Iran, and Taiwan. Recruitment methods varied, with trials typically identifying patients through treatment programs (e.g., inpatient detoxification, outpatient treatment), advertisements, referrals, or some combination of those.

Mean age was very similar across trials, usually in the 40s (32 trials) or 30s (6 trials); 3 trials did not report mean age, and 1 trial enrolled older subjects (mean age 58).⁹² All subjects met criteria for alcohol dependence in the vast majority of trials. Nine trials enrolled a majority of nonwhite subjects (60 to 100 percent).^{90-92,97-102} Most trials enrolled a third or fewer females; just 1 trial enrolled a majority of women (100 percent).¹⁰³ Just 9 trials reported information on smoking history at baseline, with most of those reporting a majority of smokers (55 to 77 percent) enrolled in those trials^{49,53,95,99,104-106} and 2 reporting a minority (17 and 47 percent).^{83,89}

Eight trials reported enrolling all or a majority of subjects with co-occurring psychiatric disorders, including bipolar disorder,¹⁰⁶ schizophrenia or schizoaffective disorder,¹⁰⁷ cocaine use disorders,^{90,91,100} depression,⁸⁹ another Axis I disorder,⁷⁶ or any comorbid psychiatric disorder.¹⁰⁸ Trials generally included or encouraged psychological or psychosocial co-interventions.

Table 7. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone

Author, Year Trial name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Additional Condition	Co-intervention	Risk of Bias
Ahmadi, 2002 ¹⁰⁹ , Ahmadi, 2004 ¹¹⁰	NTX 50 (58) Placebo (58)	12	Iran	Outpatient treatment	Self-referral	43	NR	0	NR	Individual counseling 100%	Un- clear
Anton, 1999 ¹¹¹ , Anton, 2001 ¹¹²	NTX 50 (68) Placebo (63)	12	U.S.	Outpatient academic research center	Ads, referrals for treatment-seekers	41 to 44	11 to 18	27 to 31	0	CBT 100%	Med- ium
Anton, 2005 ¹¹³	NTX 50 + CBT (39) NTX 50 + MET (41) Placebo + CBT (41) Placebo + MET (39)	12	U.S.	Outpatient	Ads, referred to clinical service	43 to 45	8 to 23	21 to 27	NR	CBT and MET as randomized	Med- ium
Anton, 2006 ⁵³ Donovan, 2008 ⁵⁶ COMBINE	ACA ^a 3,000 + CBI + MM (151) ACA 3,000 + MM (152) NTX 100 + CBI + MM (155) NTX 100 + MM (154) Placebo + CBI + MM (156) Placebo + MM (153)	16 (68)	U.S.	11 U.S. academic sites	Ads, community resources, clinical referrals at 11 academic sites	44	23	31	NR	As randomized; community support group participation (like AA) encouraged	Low
Anton, 2011 ¹⁰⁴	NTX 50 (50) Placebo (50) NTX 50 + 6 weeks gabapentin, with 1,200 maximum dose (50)	16	U.S.	Outpatient	NR	43 to 47	13	18	NR	Used COMBINE's manual (CBT + MM + 12-step techniques) 100%	Med- ium

Table 7. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone (continued)

Author, Year Trial name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Additional Condition	Co-intervention	Risk of Bias
Balldin, 2003 ⁹³	NTX 50 + CBT (25) NTX 50 +ST (31) Placebo + CBT (30) Placebo + ST (32)	26	Sweden	10 sites outpatient	Newspaper, outpatient treatment	48 to 51	NR	9 to 23	0	None	Low
Baltieri, 2008 ¹⁰⁵ ; Baltieri, 2009 ¹¹⁴	TOP target 200, maximum 400 (52) NTX 50 (49) Placebo (54)	12	Brazil	Outpatient	NR	44 to 45	29	0	NR	Psychosocial 100%	High
Brown, 2009 ¹⁰⁶	NTX 50 (20) Placebo (23)	12	U.S.	Outpatient; university health center	Newspaper ads, physician referral, flyers and brochures at clinics	41	26	49	Bipolar (current depressed or mixed mood) 100 Cannabis abuse 21 Cocaine abuse 12 Amphetamine abuse 7	CBT 100%	High
Chick, 2000 ¹¹⁵	NTX 50 (90) Placebo (85)	12	U.K.	Outpatient	From patients starting outpatient alcohol rehabilitation program	43	NR	25	0	"Usual psychosocial treatment program"	Med- ium
Fogaca, 2011 ¹¹⁶	NTX 50 (20) Placebo (20) NTX 50 + PUFA (20) PUFA (20)	12	Brazil	Outpatient	Newspaper and radio ads	NR	NR	0	NR	None	High
Garbutt, 2005 ⁸³ ; Pettinati, 2009 ¹¹⁷	NTX inj 380 (208) NTX inj 190 (210) Placebo (209)	26	U.S.	Inpatient and outpatient, private and VA	NR	45	17	32	NR	BRENDA standardized ST 100%	Med- ium
Gastpar, 2002 ¹¹⁸	NTX 50 (84) Placebo (87)	12	Germany	7 centers; outpatient	Outpatient and inpatient recruitment	43	0	28	0	Psychosocial treatment	Med- ium

Table 7. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone (continued)

Author, Year Trial name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Additional Condition	Co-intervention	Risk of Bias
Guardia, 2002 ¹¹⁹	NTX 50 (101) Placebo (101)	12	Spain	7 centers, outpatient	Recruited treatment- seeking patients	NR	NR	25	NR	Psychosocial	Med- ium
Heinala, 2001 ⁹⁴	NTX 50 daily for 12 weeks then targeted + CS (34) Placebo + CS (33) NTX 50 daily for 12 weeks then targeted + ST (29) Placebo + ST (25)	32	Finland	Outpatient	Ads	46	NR	29	0	None	High
Huang, 2005 ⁹⁷	NTX 50 (20) Placebo (20)	14	Taiwan	Alcoholism treatment unit of an inpatient psychiatric hospital; 1 week inpatient, remainder outpatient	Recruited as inpatients after admission for detoxification	38 to 43	100	0	NR	Weekly individual psychotherapy sessions 100%	High
Johnson, 2004 ⁸⁴	NTX inj 400 (35) Placebo inj (5)	17	U.S., France, the Nether- lands	4 centers; outpatient	NR	43	37	27	NR	Psychosocial support 100%	High
Kiefer, 2003 ⁶¹ Kiefer, 2004 ⁶² Kiefer, 2005 ⁶³	ACA 1,998 (40) NTX 50 (40) Placebo (40) ACA 1,998 + NTX 50 (40)	12	Germany	1 site; outpatient	Inpatient withdrawal treatment	46	NR	26	0	Group therapy	Low
Killeen, 2004 ¹⁰⁸	NTX 50 + TAU (54) Placebo + TAU(43) TAU alone (48)	12	U.S.	Outpatient community substance abuse treatment center	Clinic treatment seekers	37	24	37	Comorbid psychiatric disorder 51 Additional substance use disorder 35	Several types and intensities	Med- ium

Table 7. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone (continued)

Author, Year Trial name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Additional Condition	Co-intervention	Risk of Bias
Kranzler, 2004 ⁸⁵	NTX inj 150 (185) Placebo inj (157)	12	U.S.	Outpatient	Ads, recruited as outpatients	44	17 to 18	33 to 37	NR	MET 100%	Med- ium
Kranzler, 2009 ¹²⁰	NTX 50 targeted (38) NTX 50 once daily (45) Placebo targeted (39) Placebo once daily (41)	12	U.S.	Outpatient	Media ads, local provider referral	49	3	42	Drug use disorder <1 Social phobia 3 Antisocial personality disorder 3 Dysthymic disorder <1 Agoraphobia without panic disorder <1 OCD <1 GAD <1	Brief coping skills training 100%	Med- ium
Krystal, 2001 ⁹⁵ VACS 425	NTX 50 for 12 months (209) NTX 50 for 3 months then placebo (209) Placebo (209)	12 or 52	U.S.	Multicenter, outpatient	VA clinics	49	37	3	0	12-step facilitation	Med- ium
Latt, 2002 ¹²¹	NTX 50 (56) Placebo (51)	12 (26)	Australia	4 hospitals ; outpatient	NR	45	NR	30	0	No extensive psychosocial interventions	Med- ium
Lee, 2001 ⁹⁸	NTX 50 (35) Placebo (18)	12	Singapore	Mixed: initially inpatient, discharged after 1 month from substance abuse treatment center	Direct recruitment from inpatient facility	45	≥88	0	NR	Intensive inpatient rehabilitation program; postdischarge therapy encouraged 100%	High

Table 7. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone (continued)

Author, Year Trial name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Additional Condition	Co-intervention	Risk of Bias
Longabaugh, 2009 ⁹⁶	NTX 50 for 24 weeks + BST (36) NTX 50 for 12 weeks then placebo for 12 weeks + BST (35) NTX 50 for 24 weeks + MET (33) NTX 50 for 12 weeks then placebo for 12 weeks + MET (38) ^b	12-24 (72)	U.S.	Outpatient	Newspaper ads	44 to 46	6 to 14	33 to 43	NR	None ^c	Med- ium
Monterosso, 2001 ⁸⁶	NTX 100 (121) Placebo (62)	12	U.S.	Outpatient	Ads	46	27	27	NR	BRENDAs	Med- ium
Monti, 2001 ¹²² ; Rohsenow, 2007 ¹²³ ; Rohsenow, 2000 ¹²⁴	NTX 50 (64) Placebo (64)	12 (52)	U.S.	2 weeks partial hospital (pre- medication); 52 weeks outpatient	Recruited from partial hospital program in an urban private psychiatric hospital	39	3	24	Cocaine use 23 Sedative use 8 Opiate use 4	Brief physician outpatient contacts (intensive therapy occurred prior to medication portion of trial)	Med- ium
Morgenstern, 2012 ⁸⁷	NTX 100 + MBSCt (51) NTX 100 (51) Placebo + MBSCt (50) Placebo (48)	12	U.S.	NR	Ads, community outreach	40	26	0	HIV 15 Any drug use 67	BBCET 100%	Med- ium
Morley, 2006 ⁴⁹ Morley, 2010 ⁶⁶	ACA 1,998 (55) NTX 50 (53) Placebo (61)	12	Australia	3 treatment centers with "medical care typically available at hospital based drug and alcohol treatment services"	Patients who had attended an inpatient detoxification program, outpatient treatment, or followup or who responded to live or print ads	45	NR	30	Severe concurrent illness (psychiatric or other) –NOS 3	All offered 4 to 6 sessions of manualized compliance therapy Up-take / attendance NR	Low

Table 7. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone (continued)

Author, Year Trial name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Additional Condition	Co-intervention	Risk of Bias
Morris, 2001 ¹²⁵	NTX 50 (55) Placebo (56)	12	Australia	Outpatient	Outpatient, self-referral	47	NR	0	PTSD 23 GAD 32 Panic disorder 4 MDD 6 BPD 1	Group psychoeducation and social support	Med- ium
O'Malley, 1992 ¹²⁶ ; O'Malley, 1996 ¹²⁷	NTX 50 + CS (29) NTX 50 + ST (23) Placebo + CS (25) Placebo + ST (27)	12 (38)	U.S.	Outpatient; university alcohol treatment unit	Ads and those seeking treatment at unit	41	7	26	NR	See arms	Med- ium
O'Malley, 2007 ¹⁰³	NTX 50 (57) Placebo (50) Randomization stratified by presence of eating disorder	12	U.S.	University mental health center	Newspaper ads and patients seeking substance abuse treatment	40	11	100	Eating disorder 28	CBCST 100%, based on manualized approach used in Project MATCH	Med- ium
O'Malley, 2008 ⁹⁹	NTX 50 (34) Placebo (34) NTX 50 + SERT 100 (33)	16	U.S.	Outpatient	Direct community recruitment, health clinic referral, local ads	40	70	34	NR	MM 100%	Med- ium
Oslin, 1997 ⁹²	NTX 100 on Monday and Wednesday, 150 on Friday (21) Placebo (23)	12	U.S.	Outpatient substance abuse clinic and VAMC	From a VA hospital	58	64	NR	0	Group therapy and case manager 100%	Med- ium
Oslin, 2008 ⁸⁸	NTX 100 + CBT (40) NTX 100 + BRENDA (39) NTX 100 + doctor only (41) Placebo + CBT (40) Placebo + BRENDA (40) Placebo + doctor only (40)	24	U.S.	Outpatient psychiatry clinic	Ads in local media	41	27	27	NR	None	Med- ium

Table 7. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone (continued)

Author, Year Trial name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Additional Condition	Co-intervention	Risk of Bias
Petrakis, 2004 ¹⁰⁷ ; Ralevski, 2006 ¹²⁸	NTX 50 (16) Placebo (15)	12	U.S.	At least 3 outpatient centers— MIRECC clinics	Direct recruitment from participating centers	46	19	0	Schizophrenia or schizo- affective disorder 100	CBT + psychiatric TAU Neuroleptics 52% Benzodiazepines 16% Thymoleptics 39%	Med- ium
Petrakis, 2005 ⁷⁶ Ralevski, 2007 ⁸⁰ Petrakis, 2007 ⁸¹ Petrakis, 2006 ⁸² VA MIRECC DBRCT	DIS 250 (66) NTX 50 (59) Placebo (64) NTX 50 + DIS 250 (65)	12	U.S.	Outpatient VA	Recruited as outpatient or ads	47	26	3	Axis I disorder 100	Psychiatric TAU 100%	Med- ium for NTX vs. pla- cebo
Pettinati, 2008 ⁹¹	NTX 150 (82) Placebo (82) Subjects also randomized to either CBT or BRENDA (2x2 design) ^d	12	U.S.	University- affiliated outpatient substance abuse treatment research facility	Those seeking treatment at the facility	39	76	29	Cocaine dependence 100	NR	Med- ium
Pettinati, 2010 ⁸⁹	SERT 200 (40) NTX 100 (49) Placebo (39) SERT 200 + NTX 100 (42)	14	U.S.	Outpatient	Newspaper ads, referrals from local professional or friends/family	43	35	38	Depression 100	CBT 100%	Med- ium
Schmitz, 2004 ¹⁰⁰	NTX 50 + RPT (20) NTX 50 + DC (20) Placebo + RPT (20) Placebo + DC (20)	12	U.S.	Outpatient	Ads	36	71	16	Cocaine dependence 100	RPT or DC as randomized	High

Table 7. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone (continued)

Author, Year Trial name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Country	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Additional Condition	Co-intervention	Risk of Bias
Schmitz, 2009 ⁹⁰	NTX 100 + CBT (20) NTX 100 + CBT and CM (25) Placebo + CBT (27) Placebo + CBT and CM (14)	12	U.S.	Outpatient substance abuse clinic	Media ads	34	84 to 93	13	Cocaine use disorder 100	CBT 100%	High
Volpicelli, 1995 ¹⁰¹	NTX 50 (54) Placebo (45)	12	U.S.	Substance abuse treatment unit of a VAMC	Patients in the substance abuse treatment program of a VAMC	NR	≥78	0	NR	Outpatient treatment program and group therapy 100%	Un- clear
Volpicelli, 1997 ¹⁰²	NTX 50 (48) Placebo (49)	12	U.S.	Outpatient substance abuse treatment, university/VA treatment research center	Receiving outpatient treatment	38 to 39	60 to 65	18 to 26	NR	Counseling 100%	Med- ium

^a Three additional treatment arms were included in COMBINE but were not relevant to our Key Questions: ACA + NTX + CBI + MM, ACA + NTX + MM, and CBI only (no pills).

^b Ns are numbers analyzed, numbers randomized to each group NR. Total number randomized was 174.

^c This study is not focused on NTX versus placebo comparison; it is a different design and has 4 arms, aiming to compare 12 versus 24 weeks of NTX and to compare MET versus BST (to determine whether the type of psychosocial treatment delivered in combination with duration of NTX may partially explain inconsistent findings regarding efficacy of NTX).

^d Study stratified randomization by sex and reports the results overall and separately by sex.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: AA = Alcoholics Anonymous; ACA = acamprosate; BBCET = brief behavioral compliance enhancement treatment; BPD = bipolar disorder; CB = cognitive behavioral; CBCST = cognitive behavioral coping skills therapy; CBI = combined behavioral intervention; CBT = cognitive behavioral therapy; CS = coping skills; DC = drug counseling; GAD = generalized anxiety disorder; HIV = human immunodeficiency virus; inj = injectable; MBSCT = modified behavioral self-control therapy; MDD = major depressive disorder; MET = motivational enhancement therapy; mg = milligram; MIRECC = Mental Illness Research, Education and Clinical Center; MM = medical management;

N = number; NOS = not otherwise specified; NR = not reported; NTX = naltrexone; OCD = obsessive-compulsive disorder; PTSD = post-traumatic stress disorder; PUFA = polyunsaturated fatty acid; RPT = relapse prevention therapy; SERT = sertraline; ST = supportive therapy; TAU = treatment as usual; TOP = topiramate; U.K. = United Kingdom; U.S. = United States; VA = Veterans Affairs; VACS = Veterans Affairs Cooperative Study; VAMC = Veterans Administration Medical Center.

Return to Any Drinking

Our meta-analysis of low and medium risk of bias trials found that 4 percent fewer subjects treated with naltrexone returned to any drinking than with placebo (RD, -0.04; 95% CI, -0.07 to -0.01; 21 trials). Separating U.S.- and non-U.S.-based trials found no difference in point estimates by country (both found RD, -0.04), but the effect did not reach statistical significance for the non-U.S. trials (95% CI, -0.11 to 0.03; 7 trials). Stratifying by dose and delivery method found similar effect sizes for 50 mg per day orally (RD, -0.05), 100 mg per day orally (RD, -0.03), and injectable naltrexone (RD, -0.04), although the effect did not reach statistical significance for 100 mg per day or for injectable naltrexone.

Return to Heavy Drinking

Our meta-analysis of low and medium risk of bias trials found that 8 percent fewer subjects treated with naltrexone returned to heavy drinking than with placebo (RD, -0.08; 95% CI, -0.12 to -0.04; 21 trials). Including studies rated as high risk of bias resulted in a slightly larger effect size (RD, -0.10; 95% CI, -0.14 to -0.06; 25 trials). Separating U.S.- and non-U.S.-based trials found no difference in effect size by country (both found RD, -0.08). Stratifying by dose and delivery method found a trend toward greater effect sizes for 50 mg per day (RD, -0.09) than for 100 mg per day (RD, -0.05) or injectable naltrexone (RD, -0.07). The effect did not reach statistical significance for 100 mg per day or for injectable naltrexone, but those analyses had many fewer studies and subjects (and thus less precision) and confidence intervals largely overlapped for all three dose categories.

Drinking Days

Subjects treated with naltrexone had 4.6 percent fewer drinking days than those treated with placebo (WMD, -4.6; 95% CI, -6.6 to -2.5; 19 trials). All point estimates (of the individual studies) favored naltrexone over placebo. Stratifying our meta-analysis by U.S. and non-U.S. studies found similar effect sizes for U.S.-based (WMD, -4.5) and non-U.S.-based trials (WMD, -4.7). The effect did not reach statistical significance for non-U.S.-based trials, but the analysis had fewer studies and subjects (and thus less precision) and confidence intervals overlapped.

Stratifying by dose and delivery method found a trend toward greater effect sizes for 50 mg per day (WMD, -5.4) than for 100 mg per day (WMD, -0.86); the single study of injectable naltrexone found a larger effect size (WMD, -8.6). The effect did not reach statistical significance for 100 mg per day (95% CI, -4.2 to 2.5).

Heavy Drinking Days

Subjects treated with naltrexone had 3.6 percent fewer heavy drinking days than those treated with placebo (WMD, -3.6; 95% CI, -5.9 to -1.4; 10 trials).

Drinks per Drinking Day

Subjects treated with naltrexone had 0.6 percent fewer drinks per drinking day than those treated with placebo (WMD, -0.54; 95% CI, -1.01 to -0.07; 11 trials). Stratifying our meta-analysis by U.S. and non-U.S. studies found similar effect sizes for U.S.- and non-U.S.-based trials.

Detailed Synthesis: Placebo-Controlled Trials of Medications Used Off-label, or Those Under Investigation

We found no studies meeting inclusion criteria for amitriptyline. We found 1 placebo-controlled trial for each of the following medications: aripiprazole, atomoxetine, desipramine, fluvoxamine, imipramine, olanzapine, and paroxetine. We found insufficient evidence to support the efficacy of these medications. We provide additional details about the individual trials evaluating each of these medications in Appendix E.

We found multiple placebo-controlled trials for baclofen (2), buspirone (5), citalopram (2), fluoxetine (3), nalmefene (4), quetiapine (3), sertraline (7), topiramate (4), and valproic acid (2).

Baclofen

Characteristics of Baclofen Trials

Two trials met our inclusion criteria (Table 8). Both were parallel two-arm trials comparing baclofen with placebo for 12 weeks. Mean age was 49 years in both trials. All subjects met criteria for alcohol dependence. The trials enrolled 27 percent¹²⁹ and 45 percent females.¹³⁰ Neither trial reported information on smoking history at baseline. All patients included in 1 trial had liver cirrhosis.¹²⁹ Both trials included psychological co-interventions.

Table 8. Characteristics of included double-blind randomized placebo-controlled trials of baclofen

Author, Year	Arm	Dose, mg/day (N)	Rx Duration, Weeks	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Additional Condition	Co-intervention	Risk of Bias
Addolorato, 2007 ¹²⁹	BAC (42) Placebo (42)	30	12	Italy; university treatment and research center	People contacting alcohol treatment unit	49	NR	24 to 31	Liver cirrhosis 100 Hepatitis B 15 Hepatitis C 29	Routine psychological support 100%	Medium
Garbutt, 2010 ¹³⁰	BAC (40) Placebo (40)	30	12	U.S.; out-patient, details NR	Newspaper and radio ads	49	4	45	NR	BRENDA 100%	Medium

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: BAC = baclofen; mg = milligram; N = number; NR = not reported; U.S. = United States.

Return to Any Drinking

The trial conducted in Italy reported that a lower percentage of patients treated with baclofen returned to any drinking than with placebo (29 percent [12 of 42 patients] versus 71 percent [30 of 42]; odds ratio [OR], 6.3; 95% CI, 2.4 to 16.1).¹²⁹ The trial conducted in the United States did not report numbers for rates of return to any drinking, but reported no difference between groups for time to first usage ($p=0.13$), and included a figure for percentage abstinent that shows over 90 percent of subjects returned to any drinking over the course of the trial.¹³⁰

Return to Heavy Drinking

The trial conducted in Italy reported a greater proportion of patients in the placebo group relapsing to heavy drinking than in the baclofen group (data not reported, shown in figure only,

p=0.0062). Relapse was defined as a daily alcohol intake of more than 4 drinks or an overall consumption of 14 drinks or more per week during at least 4 weeks.¹²⁹

The trial conducted in the United States found no significant difference between groups for the proportion of patients returning to heavy drinking (hazard ratio [HR], 0.924; p=0.76).¹³⁰

Drinking Days

Only the U.S.-based trial reported data for percentage of drinking days. The trial found no significant difference between groups (baclofen versus placebo: 50.1 versus 49.4, p=0.50).¹³⁰

Heavy Drinking Days

Only the U.S.-based trial reported data for percentage of heavy drinking days. The trial found no significant difference between groups (baclofen versus placebo: 25.9 versus 25.5, p=0.73).¹³⁰

Buspirone

Characteristics of Buspirone Trials

We included 5 trials comparing buspirone with placebo (Return to Any Drinking

Just 2 of the 5 trials reported data for return to any drinking;^{132,134} 1 of them was rated as high risk of bias.¹³⁴ Neither trial found a statistically significant difference between groups, and point estimates favored placebo in both trials.

Drinking Days

Two trials rated as medium risk of bias reported data.^{131,133} One trial (N=61) enrolling anxious alcoholics reported fewer drinking days for subjects treated with buspirone than for those who received placebo (at 12 weeks: 3.6 versus 13.3, p<0.10; at posttreatment follow up 26 weeks later: 9.5 versus 24.8, p<0.01).¹³¹ One trial comparing buspirone, lithium, and placebo found no significant difference between groups (over months 1 to 3: 7 percent versus 10 percent versus 8 percent, respectively).¹³³

Drinks per Drinking Day

Just 1 trial (N=61) enrolling anxious alcoholics reported drinks per day.¹³¹ It found no statistically significant difference between subjects treated with buspirone and those who received placebo over 12 weeks (0.7 versus 2.1, p NS), or at posttreatment follow up 26 weeks later (0.9 versus 4.8, p<0.10).

Table 9). Doses ranged from 40 to 60 mg per day. Duration of treatment ranged from 12 to 52 weeks. Four trials were conducted in the United States and 1 was conducted in Canada. Mean age was very similar across trials, in the early 40s. All subjects met criteria for alcohol dependence. Two studies did not report information on race; 3 reported enrolling between 0 and 18 percent nonwhite subjects across study arms. Three trials included no women; 2 included a minority of women (18 to 26 percent across study arms). None of the trials reported information on smoking history at baseline.

Two trials enrolled a majority of subjects¹³¹ or all subjects¹³² with anxiety disorders; 1 included almost half with depression.¹³³ Most trials included or encouraged psychological or psychosocial co-interventions.

Return to Any Drinking

Just 2 of the 5 trials reported data for return to any drinking;^{132,134} 1 of them was rated as high risk of bias.¹³⁴ Neither trial found a statistically significant difference between groups, and point estimates favored placebo in both trials.

Drinking Days

Two trials rated as medium risk of bias reported data.^{131,133} One trial (N=61) enrolling anxious alcoholics reported fewer drinking days for subjects treated with buspirone than for those who received placebo (at 12 weeks: 3.6 versus 13.3, $p<0.10$; at posttreatment follow up 26 weeks later: 9.5 versus 24.8, $p<0.01$).¹³¹ One trial comparing buspirone, lithium, and placebo found no significant difference between groups (over months 1 to 3: 7 percent versus 10 percent versus 8 percent, respectively).¹³³

Drinks per Drinking Day

Just 1 trial (N=61) enrolling anxious alcoholics reported drinks per day.¹³¹ It found no statistically significant difference between subjects treated with buspirone and those who received placebo over 12 weeks (0.7 versus 2.1, p NS), or at posttreatment follow up 26 weeks later (0.9 versus 4.8, $p<0.10$).

Table 9. Characteristics of included double-blind randomized placebo-controlled trials of buspirone

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Fawcett, 2000 ¹³³	Buspirone 40 (48) Placebo (52) Lithium 1,200 (56)	26	U.S.; Outpatient	Ad, referral, inpatient/out-patient programs	40	16	0	Depression 48	Supportive therapy	Medium
George, 1999 ¹³⁵	Buspirone 60 (25) Placebo (24)	52	U.S.; Outpatient	Recruited from inpatient research unit at NIAAA	42	NR	0	0	Care of psychiatrist and nurse at posthospital clinic 100%	High
Kranzler, 1994 ¹³¹	Buspirone 15-60, mean 52.5 (31) Placebo (30)	12 (38)	U.S.; Outpatient; university health center	Ads	39 to 40	0 to 10	20 to 26	GAD 37 to 46 Anxiety disorder 50 to 52 MDD 25 to 27	CBT 100%	Medium
Malcolm, 1992 ¹³²	Buspirone target 60, mean 52 (33) Placebo (34)	26	U.S.; 1-2 weeks inpatient, then outpatient; VAMC alcohol dependence treatment unit	Screened during inpatient stay for alcohol dependence treatment	42 to 44	15 to 18	0	GAD 100	None	Medium
Malec, 1996 ¹³⁴	Buspirone 40 (28) Placebo (29)	12	Canada; hospital research center	Media ad	42	NR	18	NR	None prescribed but 37% received additional treatment: AA 7% Individual psychotherapy 3%	High

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: AA = Alcoholics Anonymous; CBT = cognitive behavioral therapy; GAD = generalized anxiety disorder; MDD = major depressive disorder; mg = milligram; N = number; NIAAA = National Institute on Alcohol Abuse and Alcoholism; NR = not reported; U.S. = United States; VAMC = Veterans Administration Medical Center.

Citalopram

Characteristics of Citalopram Trials

We included 2 trials comparing citalopram 40 mg per day with placebo for 12 to 13 weeks (Table 10). Mean age was in the mid-40s for both trials. All subjects met criteria for alcohol dependence. Neither trial reported information on race. One trial enrolled all males¹³⁶ and 1 enrolled 44 percent females.¹³⁷ One did not report information on smoking history;¹³⁶ 1 included

34 percent smokers.¹³⁷ Both trials included psychological or psychosocial co-interventions. We rated both trials as high risk of bias, primarily for high risk of attrition bias and inadequate handling of missing data (see Appendix C for details).

Table 10. Characteristics of included double-blind randomized placebo-controlled trials of citalopram

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Setting	Recruitment Method	Age, Years	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Naranjo, 1995 ¹³⁷	Citalopram 40 (53) Placebo (46)	12 (20)	Canada; outpatient research center	Newspaper ad	45	44	NR	Brief psychosocial intervention 100%	High
Tiihonen, 1996 ¹³⁶	Citalopram 40 (31) Placebo (31)	13 (17)	Finland; outpatient; community-based alcohol rehabilitation center	Inpatient / outpatient referral	45 to 47	0	0	Supportive psychotherapy intervention 100%	High

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: mg = milligram; N = number; NR = not reported.

Return to Any Drinking

The trial conducted in Finland reported 25 of 31 citalopram-treated patients and 28 of 31 placebo-treated patients returned to any drinking ($p=0.10$).¹³⁶

Drinking Days

The trial conducted in Canada found similar proportions of drinking days for those who received citalopram and those who received placebo over the 12 weeks of treatment (72.7 percent versus 76.5 percent, p NS).¹³⁷

Drinks per Drinking Day

The trial conducted in Canada found similar reductions in drinks per drinking day for those who received citalopram and those who received placebo over the 12 weeks of treatment (26.1 percent versus 26.4 percent, p NS).¹³⁷

Fluoxetine

Characteristics of Fluoxetine Trials

We included 3 trials comparing fluoxetine with placebo (Table 11). Doses ranged from 20 to 60 mg per day. Duration of treatment ranged from 12 to 15 weeks. All 3 trials were conducted in the United States. Mean age ranged from 35 to 47. All subjects met criteria for alcohol dependence. For 2 trials, about half of enrolled subjects were nonwhite;^{138,139} 1 enrolled 5 percent nonwhite subjects.¹⁴⁰ One trial enrolled all males;¹³⁹ the other 2 enrolled 20 percent¹⁴⁰ or 49 percent¹³⁸ females. None of the trials reported information on smoking history at baseline. One trial only enrolled subjects with major depressive disorder and alcohol dependence.¹³⁸ Two trials included or encouraged psychological or psychosocial co-interventions.

Table 11. Characteristics of included double-blind randomized placebo-controlled trials of fluoxetine

Author, Year	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Co- occurring Condition	Co-inter- vention	Risk of Bias
Cornelius, 1997 ¹³⁸ , Cornelius, 1995 ¹⁴¹	Fluoxetine 20-40 (25) Placebo (26)	12	U.S.; inpatient psychiatric institute	Recruited as inpatient	35	53	49	MDD 100%	Usual care: psychotherapy 100%	Medium
Kabel, 1996 ¹³⁹	Fluoxetine 20-60 (15) Placebo (13)	15	U.S.; inpatient substance abuse treatment	Inpatient recruitment	47	46	0	Cocaine use 14%	NR	High
Kranzler, 1995 ¹⁴⁰	Fluoxetine 20-60, mean 47 (51) Placebo (50)	12 (38)	U.S.; outpatient clinic	Ads	40	5	20	Major depression 14%	Group psychotherapy 79% Individual psychotherapy 21%	Medium

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: MDD = major depressive disorder; mg = milligram; N = number; NR = not reported; U.S. = United States.

Return to Any Drinking

Two small trials, 1 rated as medium risk of bias (N=51) and 1 rated as high risk of bias (N=28), reporting return to any drinking found no statistically significant difference between fluoxetine and placebo.^{138,139}

Drinking Days

Both medium risk of bias trials reported drinking days. Our meta-analysis of these 2 trials found no statistically significant difference between fluoxetine and placebo (WMD, -3.2; 95% CI, -18.2 to 11.9), but statistical heterogeneity was considerable (I^2 82.7 percent). The trial enrolling subjects who all had major depressive disorder (N=51) found that subjects treated with fluoxetine had fewer drinking days than those who received placebo (WMD, -11.6; 95% CI, -22.7 to -0.5). The trial enrolling a population with 14 percent of subjects with major depression found no difference between groups, and a point estimate trending in favor of placebo (WMD, 3.8; 95% CI, -2.1 to 9.7).

Heavy Drinking Days

The trial enrolling subjects who all had major depressive disorder (N=51) reported fewer heavy drinking days for subjects treated with fluoxetine than for those who received placebo (cumulative number of days of heavy drinking: 4.8 versus 16, $p=0.04$).

Drinks per Drinking Day

Both medium risk of bias trials reported this outcome. Similar to the analysis for drinking days, our meta-analysis of these 2 trials found no statistically significant difference between fluoxetine and placebo (WMD, -1.2; 95% CI, -4.6 to 2.2), but statistical heterogeneity was considerable (I^2 78.3 percent). The trial enrolling subjects who all had major depressive disorder

(N=51) found that subjects treated with fluoxetine had fewer drinks per drinking day than those who received placebo (WMD, -3.0; 95% CI, -5.4 to -0.6). The trial enrolling a population with 14 percent of subjects with major depression found no difference between groups, and a point estimate trending in favor of placebo (WMD, 0.5; 95% CI, -1.6 to 2.6).

Nalmefene

Characteristics of Nalmefene Trials

We included 4 trials comparing nalmefene with placebo (

Table 12). Doses ranged from 5 to 80 mg per day. One trial conducted in Finland assessed targeted dosing, instructing patients to take the medication when they believed drinking to be imminent, rather than as a daily scheduled medication.¹⁴² Duration of treatment ranged from 12 to 28 weeks. Three trials were conducted in the United States and 1 in Finland. Mean age was in the 40s in all 4 trials. All subjects met criteria for alcohol dependence in 3 trials; 1 trial reported that 93 percent met criteria for alcohol dependence.¹⁴² The trials enrolled 0 to 19 percent nonwhite subjects and from 19 to 37 percent females across study arms. None of the trials reported information on smoking history at baseline. The proportion of subjects with co-occurring psychiatric conditions was either zero or was not reported in the 4 trials.

Return to Heavy Drinking

Two trials, 1 rated as medium risk of bias (N=105)¹⁴³ and 1 pilot study rated as high risk of bias (N=21),¹⁴⁴ reported return to heavy drinking. The former found that 22 percent fewer patients treated with nalmefene returned to heavy drinking than with placebo (RD, -0.22; 95% CI, -0.42 to -0.02). The pilot study found no difference between groups (RD, -0.05; 95% CI, -0.51 to 0.41).

Drinking Days

Our meta-analysis of 2 trials,^{142,143} both rated as medium risk of bias, that reported data for this outcome found no significant difference between nalmefene and placebo (WMD, -1.1; 95% CI, -7.6 to 5.4).

Heavy Drinking Days

The trial conducted in Finland that assessed targeted dosing reported a lower percentage of heavy drinking days for patients treated with targeted nalmefene than for those who received placebo (18.1 percent versus 29.7 percent, p=0.024).

Drinks per Drinking Day

All 3 trials rated as medium risk of bias reported data. Our meta-analysis found that subjects treated with nalmefene had 1 fewer drink per drinking day than those who received placebo (WMD, -1.0; 95% CI, -1.8 to -0.3).

Table 12. Characteristics of included double-blind randomized placebo-controlled trials of nalmefene

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Anton, 2004 ¹⁴⁵	NALM 5 (68) NALM 20 (66) NALM 40 (68) Placebo (68)	12	U.S.; outpatient	Ads, recruited as outpatients	44 to 46	6 to 15	22 to 33	NR	MET 100%	Medium
Karhuvaara, 2007 ¹⁴²	NALM 10 to 40 Targeted dose ^a (242) Placebo (161)	28 ^b	Finland; 15 sites ^c	Mainly by newspaper ads	49	0	19	NR	Some elements of BRENDA	Medium
Mason, 1994 ¹⁴⁴	NALM 10 (7) NALM 40 (7) Placebo (7)	12	U.S.; NR	Ads	42	10	29	0	Group therapy 0 to 14% AA 0 to 29%	High
Mason, 1999 ¹⁴³	NALM 20 or 80 (70) Placebo (35)	12	U.S.; outpatient substance abuse treatment; academic research center	Ads, press releases, other non-specified sources	42	17 to 19	31 to 37	0	CBT (used in MATCH) 100%	Medium

^a Targeted dosing; medication was taken when subjects believed drinking to be imminent, rather than as a daily scheduled medication.

^b 52 weeks total (28 weeks of initial nalmefene vs. placebo, then another randomization for nalmefene responders).

^c Sites included 5 specialist treatment clinics, 6 private general practices, 2 occupational health care offices, and 2 outpatient clinical research facilities.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: AA = Alcoholics Anonymous; CBT = cognitive behavioral therapy; MET = motivational enhancement therapy; mg = milligram; N = number; NALM = nalmefene; NR = not reported; U.S. = United States.

Quetiapine

Characteristics of Quetiapine Trials

We included 3 trials comparing quetiapine with placebo for 12 weeks (Table 13). All 3 trials were conducted in the United States. Mean age ranged from late 30s to late 40s. All subjects met criteria for alcohol dependence in 2 trials; 1 reported that 97 percent of subjects met criteria.¹⁴⁶ Just 1 trial reported information on smoking history at baseline, with 56 percent smokers enrolled.¹⁴⁷ For 2 trials, all subjects had co-occurring bipolar disorder.^{146,147} We rated all 3 trials as high risk of bias, primarily for high risk of attrition bias, high risk of selection bias, and inadequate handling of missing data (see Appendix C for details).

Table 13. Characteristics of included double-blind randomized placebo-controlled trials of quetiapine

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Brown, 2008 ¹⁴⁶	QUET titrated from 25 to 600 over 6 weeks (52) Placebo (50)	12	U.S.; NR	From community	38	39	37	Bipolar 100	NR	High
Kampman, 2007 ¹⁴⁸	QUET 400 (29) Placebo (32)	12	U.S.; outpatient	Community referrals, media ads	47	46	23	MDD 15 Antisocial personality disorder 11 PTSD 8 Panic disorder 5 Social phobia 5 GAD 3 OCD 2	BRENDA 100%	High
Stedman, 2010 ¹⁴⁷	QUET 300-800 (175) Placebo (186)	12	U.S.; outpatient; multicenter	NR	39	12	37	Bipolar 100	None	High

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: GAD = generalized anxiety disorder; MDD = major depressive disorder; mg = milligram; N = number; NR = not reported; OCD = obsessive-compulsive disorder; PTSD = post-traumatic stress disorder; QUET = quetiapine; U.S. = United States.

Return to Any Drinking

The trial that did not enroll subjects with co-occurring bipolar disorder reported that more subjects treated with quetiapine achieved complete abstinence (9 of 29 patients versus 2 of 32, $p=0.012$)—that is, fewer subjects treated with quetiapine returned to any drinking (20 of 29 versus 30 of 32).

Drinking Days

All 3 trials reported this outcome. Our meta-analysis of the 3 trials found no difference between patients treated with quetiapine and those who received placebo (WMD, -2.7; 95% CI, -12.8 to 7.5).

Heavy Drinking Days

All 3 trials reported this outcome. Our meta-analysis of the 3 trials found no difference between patients treated with quetiapine and those who received placebo (WMD, -3.1; 95% CI, -10.1 to 4.0).

Sertraline

Characteristics of Sertraline Trials

We included 7 trials comparing sertraline with placebo (Table 14). Doses ranged from 50 to 200 mg per day. Duration of treatment ranged from 12 to 26 weeks. The majority were

Table 14. Characteristics of included double-blind randomized placebo-controlled trials of sertraline

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Brady, 2005 ¹⁴⁹	SERT 150 (49) Placebo (45)	12	U.S.; outpatient	Ads, outpatient substance abuse treatment programs	37	NR	43 to 49	PTSD 100 Depressive disorder 51 Anxiety disorder 38	CBT 100%	Medium
Coskunol, 2002 ¹⁵⁰	SERT 100 (30) Placebo (29)	26	Turkey; inpatient (mean 1 month) followed by 6 months outpatient; substance abuse treatment unit	NR	44	NR	0	For eligibility, required no concurrent Axis I disorders	Thiamine 500 mg/day 100% Pyridoxone 500 mg/day 100% AA during inpatient 100%	Medium
Gual, 2003 ¹⁵¹	SERT 50-150 (44) Placebo (39)	24	Spain; 1 center; outpatient	Outpatient alcohol dependence treatment	47	NR	47	Depression/dysthymia 100	NR	Medium
Kranzler, 2011 ¹⁵² ; Kranzler, 2012 ¹⁵³	SERT 50-200 (63) Placebo (71)	12 (26)	U.S.; outpatient; university health center	Primarily ads, some clinician referrals	48	8	19	Cannabis use disorder 17.2 Cocaine use disorder 19.4 Past MDD 20.9	Coping skills training 100%	Medium
Moak, 2003 ¹⁵⁴	SERT 50-200 (38) Placebo (44)	12	U.S.; 1 site; South Carolina; outpatient	Newspaper, outpatient treatment	41	1	39	Depression/dysthymia 100	CBT	Medium
Pettinati, 2001 ¹⁵⁵	SERT 200 (50) Placebo (50)	14	U.S.; outpatient	Ads and referral	44	80	48	Depression 47	12-step facilitation	Unclear
Pettinati, 2010 ⁸⁹	SERT 200 (40) NTX 100 (49) Placebo (39) SERT 200 + NTX 100 (42)	14	U.S.; outpatient	Newspaper ads and referrals	43	35	38	Depression 100	CBT 100%	Medium

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: CBT = cognitive behavioral therapy; MDD = major depressive disorder; mg = milligram; N = number; NR = not reported; NTX = naltrexone; PTSD = post-traumatic stress disorder; SERT = sertraline; U.S. = United States.

conducted in the United States (5 trials); 1 was conducted in Turkey and 1 in Spain. Mean age was very similar across trials, usually in the 40s. All subjects met criteria for alcohol dependence in 6 trials; 1 trial reported that 99 percent of subjects met criteria for alcohol dependence.¹⁵⁴ The percentage of nonwhite subjects enrolled was not reported by 3 trials, was a small minority (1 to

8 percent) in 2 trials, was about a third in 1 trial, and was the majority (80 percent) in 1 trial. Six trials enrolled between 19 and 48 percent females; 1 enrolled all men.¹⁵⁰ Just 1 trial reported information on smoking history at baseline, with 17 percent smokers enrolled.⁸⁹ Most trials enrolled subjects with comorbidities—3 only included patients with depressive disorders,^{89,151,154} 1 only included those with PTSD;¹⁴⁹ and 1 reported that about half of subjects had depression.¹⁵⁵ Trials typically included or encouraged psychological or psychosocial co-interventions.

Return to Any Drinking

Just 1 trial reported this outcome—the trial that compared sertraline, naltrexone, sertraline plus naltrexone, and placebo.⁸⁹ It found similar proportions of patients treated with sertraline returning to any drinking as with placebo (29 of 40 patients versus 30 of 39, p NS).

Return to Heavy Drinking

Two of the trials reported this outcome—the trials conducted in Turkey and Spain.^{150,151} Our meta-analysis found no difference between patients treated with sertraline and those who received placebo (RD, -0.04; 95% CI, -0.31 to 0.23).

Drinking Days

Three of the trials reported this outcome—the trial conducted in Spain,¹⁵¹ the trial conducted in South Carolina,¹⁵⁴ and the U.S.-based trial rated as unclear risk of bias.¹⁵⁵ Our meta-analysis found no significant difference between patients treated with sertraline and those who received placebo, both without (WMD, 0.03; 95% CI, -11.0 to 11.1) and with inclusion of the trial rated as unclear risk of bias (WMD, -0.7; 95% CI, -8.2 to 6.9).

Heavy Drinking Days

Just 1 trial reported this outcome—the trial that enrolled patients with PTSD and alcohol dependence (N=94).¹⁴⁹ It reported numerically more heavy drinking days for patients treated with sertraline than for those who received placebo, but the difference was not statistically significant (number of heavy drinking days: mean, standard deviation [SD], 10.4, 2.3 versus 8.9, 2.5).

Drinks per Drinking Day

Two of the trials reported this outcome—the trial conducted in South Carolina (N=82)¹⁵⁴ and the U.S.-based trial that enrolled patients with PTSD and alcohol dependence (N=94).¹⁴⁹ Our meta-analysis found no significant difference between patients treated with sertraline and those who received placebo (WMD, -0.9; 95% CI, -2.2 to 0.5).

Topiramate

Characteristics of Topiramate Trials

We included 4 trials comparing topiramate with placebo for 12 to 14 weeks (Table 15). Two trials were conducted in the United States, 1 in Brazil, and 1 in Spain. Mean age was in the 40s in all 4 trials. All subjects met criteria for alcohol dependence in 3 trials; 1 trial did not report the proportion with alcohol dependence, but most subjects likely had alcohol dependence.¹⁵⁶ Two trials enrolled all males;^{105,157} the other 2 included from 26 to 40 percent females across study arms. The 2 non-U.S.-based trials reported information on smoking history at baseline, with 66

to 80 percent smokers enrolled in those trials.^{105,157} Three of the 4 trials offered or included psychological or psychosocial co-interventions.

Table 15. Characteristics of included double-blind randomized placebo-controlled trials of topiramate

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Baltieri, 2008 ¹⁰⁵ ; Baltieri, 2009 ¹¹⁴	TOP target 200, maximum 400 (52) NTX 50 (49) Placebo (54)	12	Brazil; outpatient	NR	44 to 45	29	0	NR	Psychosocial 100%	High
Johnson, 2003 ¹⁵⁶ ; Ma, 2006 ¹⁵⁸ ; Johnson, 2004 ¹⁵⁹	TOP 25-300 (75) Placebo (75)	12	U.S.; 1 site; Texas; outpatient	Newspaper	41	NR	28 to 40	0	None	Medium
Johnson, 2007 ¹⁶⁰ ; Johnson, 2008 ¹⁶¹	TOP 50-300, mean 171 (183) Placebo (188)	14	U.S.; 17 academic sites	From academic sites; by newspaper, radio, television ads	47 to 48	15	26 to 28	NR	BBCET 100%	Low
Rubio, 2009 ¹⁵⁷	TOP 250 (31) Placebo (32) ^a	12	Spain; outpatient	NR	42	NR	0	NR	Supportive group therapy offered	High

^a Numbers entered are those analyzed; 76 total were randomized, but dropouts were not reported by arm.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: BBCET = brief behavioral compliance enhancement treatment; mg = milligram; N = number; NR = not reported; NTX = naltrexone; TOP = topiramate; U.S. = United States.

Return to Any Drinking

Just 1 trial reported this outcome—the trial conducted in Brazil that was rated as high risk of bias.¹⁰⁵ It reported that more patients treated with topiramate returned to any drinking than with placebo (24 of 52 patients versus 15 of 54).

Drinking Days

Three of the trials reported this outcome—2 U.S.-based trial rated as low (N=371)¹⁶⁰ or medium risk of bias (N=150)¹⁵⁶ and the trial conducted in Spain (N=63) that was rated as high risk of bias.¹⁵⁷ Our meta-analysis found a lower percentage of drinking days for patients treated with topiramate than for those who received placebo both without (WMD, -8.5; 95% CI, -15.9 to -1.1) and with inclusion of the trial rated as high risk of bias (WMD, -9.7; 95% CI, -16.4 to -3.1). We were unable to include the smaller U.S.-based trial (N=150) in the meta-analysis due to differences in the type of data reported—it reported that subjects treated with topiramate had a greater percentage of days abstinent than those who received placebo (mean difference -11.6, 95% CI -3.98 to -19.3).

Heavy Drinking Days

Three of the trials reported this outcome—2 U.S.-based trials rated as low (N=371)¹⁶⁰ or medium risk of bias (N=150)¹⁵⁶ and the trial conducted in Spain (N=63) that was rated as high risk of bias.¹⁵⁷ Our meta-analysis found a lower percentage of heavy drinking days for patients treated with topiramate than for those who received placebo both without (WMD, -11.5; 95% CI, -18.3 to -4.8) and with inclusion of the trial rated as high risk of bias (WMD, -12.5; 95% CI, -17.9 to -7.2).

Drinks per Drinking Day

Three of the trials reported this outcome—2 U.S.-based trial rated as low (N=371)¹⁶⁰ or medium risk of bias (N=150)¹⁵⁶ and the trial conducted in Spain (N=63) that was rated as high risk of bias.¹⁵⁷ Our meta-analysis found that patients treated with topiramate had fewer drinks per drinking day than those treated with placebo both without (WMD, -1.1; 95% CI, -1.7 to -0.4) and with inclusion of the trial rated as high risk of bias (WMD, -1.2; 95% CI, -1.8 to -0.6).

Valproic Acid

Characteristics of Valproic Acid Trials

We included 2 trials comparing valproic acid with placebo (Table 16). Duration of treatment ranged from 12 to 24 weeks. Both trials were conducted in the United States. Mean age was very similar across trials, 38 to 40. All subjects met criteria for alcohol dependence. The trials enrolled from 25 percent¹⁶² to 54 percent¹⁶³ nonwhite subjects, and from 29 percent¹⁶² to 62 percent¹⁶³ women. Just 1 of the trials reported information on smoking history at baseline, reporting that 71 percent of subjects were smokers.¹⁶² One trial only enrolled subjects with bipolar disorder.¹⁶² Both trials included co-interventions—1 with lithium and weekly dual diagnosis (alcohol dependence and bipolar disorder) recovery counseling,¹⁶² and 1 with cognitive behavioral therapy.¹⁶³

Table 16. Characteristics of included double-blind randomized placebo-controlled trials of valproic acid

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Brady, 2002 ¹⁶³	Valproic acid 1,500 (14) Placebo (15)	12	U.S.; outpatient	Newspaper; several treatment settings	40	54	62	0	CBT	Medium
Salloum, 2005 ¹⁶²	Valproate 750+ (29) Placebo (30)	24	U.S.; outpatient substance abuse service at university clinic	Treatment seekers	38	25	29	Bipolar I disorder 100 Mixed bipolar subtype 58 Manic 21 Depressed 21 Cannabis abuse or dependence 29 Cocaine abuse 29	Lithium and weekly individual dual diagnosis recovery counseling 100%	Medium

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: CBT = cognitive behavioral therapy; mg = milligram; N = number; U.S. = United States.

Return to Any Drinking

One trial (N=29) reported no significant difference in the percentage of subjects who returned to any drinking over 12 weeks (valproic acid versus placebo: 81 versus 83, p NS).¹⁶³

Return to Heavy Drinking

Our meta-analysis found that 33 percent fewer subjects treated with valproic acid returned to heavy drinking than with placebo (RD, -0.33; 95% CI, -0.55 to -0.11; 2 trials).

Drinking Days

One trial (N=29) reported no significant difference in the percentage of drinking days over 12 weeks between subjects treated with valproic acid and those who received placebo (15.9 versus 19.6, p NS).¹⁶³

Heavy Drinking Days

Our meta-analysis found a lower percentage of heavy drinking days for patients treated with valproic acid than for those who received placebo (WMD, -8.5; 95% CI, -15.9 to -1.1; 2 trials).

Drinks per Drinking Day

Our meta-analysis found that subjects treated with valproic acid had 2.6 fewer drinks per drinking day than those who received placebo (WMD, -2.6; 95% CI, -5.0 to -0.2; 2 trials).

Detailed Synthesis: Head-to-Head Trials

Acamprosate versus Disulfiram

Characteristics of Trials

We found no studies meeting our inclusion criteria. Our searches did identify some studies comparing acamprosate with disulfiram that did not meet our inclusion criteria for this section because they were open-label studies.^{164,165}

Acamprosate versus Naltrexone

Characteristics of Trials

We included 3 trials comparing acamprosate with naltrexone (Table 17). Two used 50 mg per day doses for naltrexone;^{49,61} 1 used 100 mg per day.⁵³ Two used 1,998 mg per day doses for acamprosate;^{49,61} 1 used 3,000 mg per day.⁵³ Duration of treatment ranged from 12 to 16 weeks. One trial was conducted in the United States, 1 in Germany, and 1 in Australia. Mean age was in the mid-40s for all 3 trials. All subjects met criteria for alcohol dependence in 2 trials; 1 trial did not report the proportion with alcohol dependence, but most subjects likely had alcohol dependence.⁴⁹ Two studies did not report information on race; 1 trial reported enrolling 23 percent nonwhite subjects.⁵³ The trials enrolled a similar proportion of women (26 to 31 percent). Two trials reported information on smoking history at baseline—1 reported that 55 percent of pill-taking subjects were smokers;⁵³ 1 reported that 72 to 81 percent of subjects were smokers across study arms.⁴⁹ Trials included or encouraged psychological or psychosocial co-interventions.

Table 17. Characteristics of double-blind head-to-head randomized controlled trials of acamprosate versus naltrexone

Author, Year Trial Name	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Co- occurring Condition	Co-inter- vention	Risk of Bias
Anton, 2006 ⁵³ Donovan, 2008 ⁵⁶ COMBINE	ACA 3,000 + CBI + MM (151) ACA 3,000 + MM (152) NTX 100 + CBI + MM (155) NTX 100 + MM (154) Placebo + CBI + MM (156) Placebo + MM (153) ^a	16 (68)	U.S.; 11 academic sites	Ads, community resources, clinical referrals	44	23	31	NR	Community support group participation (like AA) encouraged	Low
Kiefer, 2003 ⁶¹ Kiefer, 2004 ⁶² Kiefer, 2005 ⁶³	ACA 1,998 (40) NTX 50 (40) Placebo (40) ACA 1,998 + NTX 50 (40)	12	Germany; 1 site in Hamburg; outpatient	From inpatient withdrawal treatment	46	NR	26	0	Group therapy	Low
Morley, 2006 ⁴⁹ Morley, 2010 ⁶⁶	ACA 1,998 (55) NTX 50 (53) Placebo (61)	12	Australia; 3 treatment centers in Sydney	Patients who had attended an inpatient detoxification program, outpatient treatment, or followup, or who responded to live or print ads	45	NR	30	Severe concurrent illness (psychiatric or other) –NOS 3	All offered 4-6 sessions of manualized compliance therapy	Low

^aThree additional treatment arms were included in COMBINE but were not relevant to our Key Questions: ACA + NTX + CBI + MM, ACA + NTX + MM, and CBI only (no pills).

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: AA = Alcoholics Anonymous; ACA = acamprosate; CBI = combined behavioral intervention; mg = milligram; MM = medical management; N = number; NA = not applicable; NOS = not otherwise specified; NTX = naltrexone; U.S. = United States.

Return to Any Drinking

Our meta-analysis found no statistically significant difference between naltrexone and acamprosate (RD, 0.02; 95% CI, -0.03 to 0.08; 3 trials).

Return to Heavy Drinking

Our meta-analysis found no statistically significant difference between naltrexone and acamprosate (RD, 0.01; 95% CI, -0.06 to 0.07; 3 trials).

Drinking Days

Two of the 3 trials reported sufficient data for meta-analysis for drinking days; neither found a statistically significant difference between treatments.^{49,53} Our meta-analysis found no statistically significant difference between naltrexone and acamprosate (WMD, -2.98; 95% CI, -13.4 to 7.5).

Heavy Drinking Days

The COMBINE study reported that analyses of alternative summary measures of drinking, including heavy drinking days per month ($p=0.006$) were consistent with those for the coprimary end points (percentage of days abstinent from alcohol and time to first heavy drinking day), all showing a significant naltrexone by combined behavioral intervention (CBI) interaction.

Drinks per Drinking Day

Two of the trials reported some information about drinks per drinking day, but not enough data for us to conduct quantitative synthesis. The trial conducted in Australia reported no statistically significant difference between acamprosate and naltrexone (mean [SD], 7.5 [6.1] versus 5.9 [6.1]; p not reported). The COMBINE study reported that analyses of alternative summary measures of drinking, including drinks per drinking day ($p=0.03$), were consistent with those for the coprimary end points (percentage of days abstinent from alcohol and time to first heavy drinking day), all showing a significant naltrexone by CBI interaction.

Disulfiram Versus Naltrexone

Characteristics of Trials

We included 1 trial comparing disulfiram with naltrexone (Table 18). It compared disulfiram, naltrexone, placebo, and the combination of disulfiram plus naltrexone for 12 weeks in Veterans Administration outpatient settings. All subjects met criteria for alcohol dependence and had co-occurring Axis I psychiatric disorders. Almost all subjects were male. The trial did not report information on smoking history at baseline.

Table 18. Characteristics of double-blind head-to-head randomized controlled trials of disulfiram versus naltrexone

Author, Year Trial Name	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	% With Co- occurring Condition	Co-inter- vention	Risk of Bias
Petrakis, 2005 ⁷⁶ Ralevski, 2007 ⁸⁰ Petrakis, 2007 ⁸¹ Petrakis, 2006 ⁸² VA MIRECC	DIS 250 (66) NTX 50 (59) Placebo (64) NTX 50 + DIS 250 (65)	12	U.S.; out- patient VA	Recruited as outpatients or by ads	47	26	3	Axis I disorder 100	Psychiatric treatment as usual 100%	High ^a

^aHigh risk of bias for disulfiram versus naltrexone; medium for naltrexone versus placebo.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: DIS = disulfiram; mg = milligram; MIRECC = Mental Illness Research, Education and Clinical Center; N = number; NTX = naltrexone; U.S. = United States; VA = Veterans Administration.

The study used a double-blind design for the comparison between naltrexone and placebo, but not for disulfiram (which was given open label). We rated the trial as high risk of bias for the comparison between disulfiram and naltrexone, primarily for high risk of ascertainment bias (see Appendix C for details; we rated it as medium risk of bias for naltrexone versus placebo).

Other studies that did not meet our inclusion criteria for this section comparing disulfiram with naltrexone were either open-label studies¹⁶⁵⁻¹⁶⁷ or were conducted in adolescents.¹⁶⁸

Return to Any Drinking

The trial reported no statistically significant difference between disulfiram and naltrexone for number of subjects achieving total abstinence (51 versus 38, $p=0.11$).

Drinking Days

The trial reported no statistically significant difference between disulfiram and naltrexone for the percentage of days abstinent (96.6 versus 95.4, $p=0.55$).

Heavy Drinking Days

The trial reported no statistically significant difference between disulfiram and naltrexone for the percentage of heavy drinking days (3.2 versus 4, $p=0.65$).

Head-to-head Trials Including Medications Used Off-label, or Those Under Investigation

Characteristics of Trials

We found 4 eligible trials (

Table 19). All 4 utilized naltrexone; none treated subjects with acamprosate or disulfiram. Off-label medications evaluated included aripiprazole, desipramine, paroxetine, sertraline, and topiramate. No 2 trials assessed the same head-to-head comparison. Duration of treatment ranged from 12 to 16 weeks. Two were conducted in the United States, 1 in Brazil, and 1 in Italy. Mean age of subjects was similar across trials (in the 40s). All subjects met criteria for alcohol dependence. One trial enrolled all males,¹⁰⁵ and 1 did not report information on sex.¹⁶⁹ The other 2 included 9 to 38 percent women.^{89,170} One trial only included subjects with both PTSD and alcohol dependence;¹⁷⁰ 1 only included those with depression and alcohol dependence.⁸⁹ Trials typically included or encouraged psychological or psychosocial co-interventions.

Aripiprazole Compared with Naltrexone

The only included trial reported no significant differences between groups for number of subjects who remained abstinent, number of subjects who relapsed, mean number of abstinent days, and heavy drinking days.¹⁶⁹

Desipramine Compared with Paroxetine

One included trial, rated as high risk of bias, randomized patients with PTSD and alcohol dependence to desipramine, paroxetine, desipramine plus naltrexone, or paroxetine plus naltrexone.¹⁷⁰ The trial found that patients treated with desipramine had fewer heavy drinking days ($p=0.009$) and drinks per drinking day ($p=0.027$) than those who received paroxetine.

Table 19. Characteristics of double-blind head-to-head randomized controlled trials including medications used off-label, or those under investigation

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks	Setting	Recruitment method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Martinotti, 2009 ¹⁶⁹	Aripiprazole 5-15 (29) NTX 50 (28)	16	Italy; out-patient; university hospital day clinic	Direct recruitment from local facility	40	NR	NR	Mood disorder 19 Anxiety disorder 11 ^a	None required	Medium
Petrakis, 2012 ¹⁷⁰	Desipramine 200 + placebo (24) ^b Paroxetine 40 + placebo (20) Desipramine 200 + NTX 50 (22) Paroxetine 40 + NTX 50 (22)	12	U.S.; out-patient; multiple mental illness centers, most from VAs	Local advertising (nonveterans); mental illness centers (veterans)	47	25	9	PTSD 100	Clinical management/compliance enhancement therapy 100%	High
Pettinati, 2010 ⁸⁹	Sertraline 200 (40) NTX 100 (49) Placebo (39) Sertraline 200 + NTX 100 (42)	14	U.S.; out-patient	Newspaper ads, referrals	43	35	38	Depression 100	CBT 100%	Medium
Baltieri, 2008 ¹⁰⁵ , Baltieri, 2009 ¹¹⁴	Topiramate target 200, maximum 400 (52) NTX 50 (49) Placebo (54)	12	Brazil; outpatient	NR	44 to 45	29	0	NR	Psychosocial 100%	High

^a Study also reported the following percentages of subjects with co-occurring disorders: impulse control disorder 5%, eating disorder 1%, somatoform disorder 1%. Personality disorders: borderline 8%, antisocial 4%, avoidant 4%, histrionic 1%, paranoid 1%, dependent 1%, passive-aggressive 1%, schizoid 1%, cannabis abuse 12%, cocaine abuse 8%, benzodiazepine abuse 1%, MDMA abuse 1%.

^b Because 2 of the 4 arms are combinations, they are not eligible/not comparisons of interest; only the head-to-head comparison of paroxetine + placebo and desipramine + placebo is eligible.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: CBT = cognitive behavioral therapy; mg = milligram; N = number; NR = not reported; NTX = naltrexone; PTSD = post-traumatic stress disorder; U.S. = United States; VA = Veterans Affairs.

Sertraline Compared with Naltrexone

The only included trial reported a higher abstinence rate for patients (all had alcohol dependence and co-occurring depression) who received the combination of sertraline and naltrexone than for those who received either naltrexone, sertraline, or placebo only (53.7 percent versus 21.3 percent versus 27.5 percent versus 23.1 percent; $p=0.001$).⁸⁹ The difference between naltrexone and sertraline given alone was not significant.

Topiramate Compared with Naltrexone

The only included trial, rated as high risk of bias, reported no significant differences between topiramate and naltrexone for proportion of abstinent subjects, cumulative abstinence duration,

time to first relapse, or heavy drinking weeks.¹⁰⁵ Significantly more subjects in the topiramate group participated in AA than in the naltrexone group (19.2 percent versus 4.1 percent, $p=0.04$).

Systematic Reviews

We included 5 systematic reviews for this KQ.¹⁷¹⁻¹⁷⁵ Two were reviews from the Cochrane Collaboration assessing acamprosate¹⁷¹ or opioid antagonists (naltrexone and nalmefene).¹⁷³ One assessed the efficacy of disulfiram,¹⁷² 1 was a sex-specific individual patient data meta-analysis of response to acamprosate,¹⁷⁴ and 1 was for the United Kingdom's National Institute for Clinical Excellence (NICE) guidelines on alcohol-use disorders.¹⁷⁵ All 5 were rated as low or medium risk of bias. In general, the 5 systematic reviews did not report findings that conflict with our results, so we describe them only briefly in this report. None of the reviews included publications from the past few years, as literature searches were typically completed 3 or more years ago.

The Cochrane Collaboration review of acamprosate for people with alcohol dependence (literature searches through January 2009) found acamprosate to be effective.¹⁷¹ It reported a 14 percent reduction in return to any drinking compared with placebo (relative risk [RR], 0.86; 95% CI, 0.81 to 0.91; number needed to treat [NNT], 9.1; 95% CI, 6.7 to 14.3). The Cochrane Collaboration review of opioid antagonists (literature searches through January 2010) reported that naltrexone reduced the risk of heavy drinking (RR, 0.83; 95% CI, 0.76 to 0.90), decreased drinking days (WMD, -3.89; 95% CI, -5.75 to -2.04), and decreased heavy drinking days (WMD, -3.25; 95% CI, -5.51 to -0.99) compared with placebo.¹⁷³ Effects of naltrexone on return to any drinking were not statistically significant (RR, 0.96; 95% CI, 0.92 to 1.00).

The sex-specific individual patient data meta-analysis (N=6,111) of response to acamprosate found a significant effect of acamprosate compared with placebo for improving rates of abstinence and no heavy drinking in both women and men.¹⁷⁴ Men and women did not differ on any measure of acamprosate efficacy.

The review for the NICE guidelines found that the evidence for both acamprosate and naltrexone supports their efficacy for improving alcohol consumption outcomes.¹⁷⁵ It reported a significant effect of acamprosate in promoting abstinence when compared with placebo (RR, 0.83; 95% CI, 0.77 to 0.88) and for the number of individuals relapsing to heavy drinking (RR, 0.90; 95% CI, 0.81 to 0.99). It found that oral naltrexone reduced rates of relapse to heavy drinking (RR, 0.83; 95% CI, 0.75 to 0.91), reduced mean drinks per drinking day (standardized mean difference [SMD], -0.28; 95% CI, -0.44 to -0.11), and reduced days of heavy drinking (SMD -0.43; 95% CI, -0.82 to -0.03) compared with placebo. Oral disulfiram was not significantly different from placebo in preventing participants from lapsing to alcohol consumption (RR 1.05; 95% CI, 0.96 to 1.15).

Key Question 2. Health Outcomes

For this key question (KQ), we describe the characteristics of included studies and then results for the included health outcomes (accidents, injuries, quality of life [QoL], function, and mortality). Throughout this KQ, we include headers and sections only for outcomes reported by the included studies.

Detailed Synthesis: Placebo-Controlled Trials of FDA-Approved Medications for Treating Alcohol Dependence

We found no placebo-controlled trials of disulfiram that reported a health outcome of interest. Below we describe 9 placebo-controlled trials of acamprosate and 9 placebo-controlled trials of naltrexone that reported a health outcome of interest (including the COMBINE study, which has comparisons between placebo and both medications). These represent a subset of the trials included in KQ 1.

Acamprosate

Characteristics of Trials

Nine placebo-controlled RCTs reported a health outcome (

Table 20). Sample sizes ranged from 110 to 612 participants in acamprosate plus placebo arms. Duration of treatment ranged from 12 to 52 weeks. Followup to 1 year or longer was available for 6 trials. No studies identified a health outcome as their primary outcome.

The mean age of patients ranged from 40 to 45. All patients enrolled in the trials had alcohol dependence. Only 1 trial reported on race: 14 to 15 percent of patients were nonwhite.⁵⁴ Females made up 18 to 36 percent of the patients across studies. Two trials reported smoking status at baseline, from 46 percent⁵⁴ to 55 percent.⁵³ No trials specified the percentage of patients who had a co-existing medical or psychiatric condition.

There was minor variation in the dosing of acamprosate across trials. Most studies used doses from 1,332 to 1,998 mg per day and determined dosing based on weight. Two studies included an arm who received 3 g per day.^{53,54} Three studies commented on the use of other pharmacotherapy to address alcohol or comorbid psychiatric disorders. One trial allowed the use of disulfiram on a voluntary basis.⁵⁷ Two other trials reported that 5 to 6 percent of patients in either treatment group were prescribed benzodiazepines.⁵⁹, and 1 trial allowed the use of “hypnotics, anxiolytics or antidepressants” in either group.⁶⁷

Two studies were conducted in the United States,^{53,54} all others were conducted in European countries. No studies were conducted in primary care settings; most were conducted in outpatient substance abuse or psychiatric treatment centers. The majority of trials recruited patients during or shortly after discharge from an inpatient substance abuse treatment center. One US trial recruited patients via newspaper advertisement⁵⁴ and 1 German trial recruited patients from outpatient substance abuse treatment centers.⁷² The COMBINE study recruited patients by advertisement and referral from 11 academic centers.⁵³

Table 20. Characteristics of included double-blind randomized placebo-controlled trials of acamprostate that report a health outcome

Author, Year Trial Name Design	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Setting	Recruitment Method	Age, Years	% Non-white	% Female	Co-intervention	Risk of Bias
Anton, 2006 ⁵³ Donovan, 2008 ⁵⁶ LoCastro, 2009 ¹⁷⁶ COMBINE	ACA 3,000 + CBI + MM (151) ACA 3,000 + MM (152) NTX 100 + CBI + MM (155) NTX 100 + MM (154) Placebo + CBI + MM (156) Placebo + MM (153) ^a	16 (68)	U.S.; 11 academic sites	Ads, community resources, clinical referrals	44	23	31	As randomized; Low community support group participation (like AA) encouraged	Low
Besson, 1998 ⁵⁷	ACA 1,300 to 1,998 (55) Placebo (55)	52 (108)	Switzerland; Outpatient; 3 psychiatric treatment centers	From inpatient treatment unit	42	NR	20	Routine counseling 100% Voluntary disulfiram 22% to 24%	Medium
Geerlings, 1997 ⁵⁹	ACA 1,332 to 1,998 (128) Placebo (134)	26 (52)	Belgium, the Netherlands, and Luxembourg; Outpatient substance abuse treatment centers	Recruited from detoxification patients in same centers	40 to 42	NR	24	ACA: benzodiazepines 5% Placebo: benzodiazepines 6%	Medium
Lhuointre, 1990 ⁶⁵	ACA 1,332 (279) Placebo (290)	12 (12)	France; Outpatient substance abuse treatment centers	Inpatient treatment centers (30 centers across France)	42 to 43	NR	18	None	Unclear
Mason, 2006 ⁵⁴	ACA 2,000 (258) ACA 3,000 (83) Placebo (260)	24 (32)	U.S.; 21 outpatient clinics ^b	Primarily by newspaper ads	44 to 45	14 to 15	29 to 36	Brief abstinence-oriented protocol-specific counseling and self-help materials 100%	Low
Paille, 1995 ⁶⁷	ACA 1.3 g (188) ACA 2 g (173) Placebo (177)	52 (78)	France; NR ^c	Referral from alcohol specialist centers	43	NR	20	Supportive psychotherapy 100% Hypnotics 6 to 7% Anxiolytics 8 to 12% Antidepressants 8 to 9%	Medium

Table 20. Characteristics of included double-blind randomized placebo-controlled trials of acamprosate that report a health outcome (continued)

Author, Year Trial Name Design	Arm Dose, mg/day (N)	Rx Duration, Weeks (Fol-lowup)	Setting	Recruitment Method	Age, Years	% Non-white	% Female	Co-inter-vention	Risk of Bias
Poldrugo, 1997 ⁷⁰	ACA 1,332 to 1,998 (122) Placebo (124)	26 (52)	Italy; Inpatient for 1-2 weeks then outpatient; multicenter community-based alcohol rehabilitation program	From acute inpatient withdrawal treatment	43 to 45	NR	23 to 31	Community-based rehabilitation program with group sessions, alcohol education, community meetings 100%	Medium
Sass, 1996 ⁷²	ACA 1,332 to 1,998 (136) Placebo (136)	48 (96)	Germany; Psychiatric outpatient	Outpatient referral	41 to 42	NR	22	Counseling / psychotherapy 100%	Medium
Whitworth, 1996 ⁷⁴	ACA 1,332 or 1,998 (224) Placebo (224)	52 (52)	Austria; Outpatient specialty	Recruited after inpatient detoxification	42	NR	21	NR	Medium

^a Three additional treatment arms were included in COMBINE but were not relevant to our Key Questions: ACA + NTX + CBI + MM, ACA + NTX + MM, and CBI only (no pills).

^b Clinics were affiliated with academic medical centers and had investigators experienced in alcoholism treatment.

^c The article was not explicit about the setting, but patients received psychotherapy and psychiatric medication management suggesting a psychiatric outpatient setting.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: AA = Alcoholics Anonymous; ACA = acamprosate; CBI = combined behavioral intervention; mg = milligram; MM = medical management; N = number; NR = not reported; NTX = naltrexone; U.S. = United States.

Eight trials were rated as low or medium risk of bias. One trial was rated as unclear risk of bias, primarily due to unclear handling of missing data and unclear masking of outcome assessors (see Appendix C for details).⁶⁵

Accidents or Injuries

We identified 1 study, rated as unclear risk of bias, reporting that one patient in the placebo group died by “accident.” No other details on the cause or nature of the accident were provided.⁶⁵

Quality of Life or Function

The COMBINE study assessed QoL using the World Health Organization Quality of Life (WHOQOL) and 12-item Short-Form Health Survey (SF-12v2) physical and mental health scores. Results were not presented for each treatment group separately.¹⁷⁶ These results are discussed in detail in the acamprosate versus naltrexone section (below). Briefly, no clinically significant differences were found across the eight combinations of pharmacological and behavioral treatments for QoL for acamprosate compared with placebo.¹⁷⁶

Mortality

Eight trials of acamprosate reported on mortality. Few deaths were reported; no study reported more than two deaths in any group. Table 21 shows the number of deaths in studies which report deaths per study arm. In the COMBINE trial, the authors reported that one fatal serious adverse event was reported during the 16-week treatment phase. Investigators classified this death as being unrelated to the study medication. No details were provided on which group the death occurred in, the nature of the adverse event, or the cause of death.⁵³

Table 21. Mortality reported in placebo-controlled trials of acamprosate

Author, Year	Study Duration, Weeks	N (Cause) Deaths, Placebo Arm	N (Cause) Deaths, Acamprosate Arm
Besson, 1998 ⁵⁷	52	1 (cardiac arrest)	0
Geerlings, 1997 ⁵⁹	26	0	0
Mason, 2006 ³⁴	26	0	0
Paille, 1995 ⁶⁷	51	2 (NR)	1.3 g arm: 2 (NR) 2.0 g arm: 2 (NR)
Poldrugo, 1997 ⁷⁰	26	1 (NR)	0
Sass, 1996 ⁷²	52	1 (suicide, by hanging)	1 (suicide, by hanging)
Whitworth, 1996 ⁷⁴	26	1 (NR)	2 (NR)

Abbreviations: N = number; NR = not reported.

Naltrexone

Characteristics of Trials

Nine RCTs comparing naltrexone with placebo reported at least one health outcome of interest (Table 22). All 9 trials were rated as low or medium risk of bias. Sample sizes ranged from 31 to 618 participants in the naltrexone plus placebo arms. Duration of treatment ranged from 12 to 26 weeks.

Mean age was similar across trials, ranging from 39 to 50. Two trials included only male patients;^{87,107} females made up 3 to 38 percent of patients in the other trials. One study did not report on the race of study participants⁹³; most of the other trials enrolled a minority of nonwhite subjects (17 to 35 percent) and 2 enrolled a majority (70 to 76 percent).^{91,99} Three studies provided information on smoking status; approximately half of participants in those trials were smokers.^{53,83,99} All trials enrolled a vast majority (93 percent or more) of patients with alcohol dependence. Three trials did not specifically include (or describe) whether study participants had any co-existing medical or psychiatric disorders.^{83,91,93} One trial was conducted among men who have sex with men; 67 percent reported any other drug use and 15 percent had HIV.⁸⁷ Four trials were conducted among populations who all had a specific psychiatric comorbidity: 1 among patients with either schizophrenia or schizoaffective disorder,¹⁰⁷ 1 among patients with cocaine dependence,⁹¹ 1 among patients with at least one other psychiatric (Axis I) disorder,⁷⁶ and 1 among patients with depression.⁸⁹

One trial evaluated the efficacy of two doses of injectable naltrexone⁸³ and the remainder randomized patients to oral naltrexone either at 50, 100,^{87,89,176} or 150 mg per day.⁹¹ Four trials described a specific behavioral or psychological co-intervention.^{83,87,93,117} Two trials conducted among those with a psychiatric comorbidity specified that patients continued medical management and usual psychiatric care^{76,107} and 1 included cognitive behavioral therapy for depressed patients.⁸⁹ No specific co-intervention was described in the trial comparing naltrexone with placebo in patients with cocaine dependence.⁹¹

Table 22. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone that report a health outcome

Author, Year Trial Name	Arm Dose, mg/day (N)	Rx Duration, Weeks (Follow- up)	Setting	Recruitment Method	Age, Years	% Non- white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Anton, 2006 ⁵³ Donovan, 2008 ⁵⁶ LoCastro, 2009 ¹⁷⁶ COMBINE	ACA 3,000 + CBI + MM (151) ACA 3,000 + MM (152) NTX 100 + CBI + MM (155) NTX 100 + MM (154) Placebo + CBI + MM (156) Placebo + MM (153) ^a	16 (68)	U.S.; 11 academic sites	Ads, community resources, clinical referrals at 11 academic sites	44	23	31	NR	As randomized; community support group participation (like AA) encouraged	Low
Balldin, 2003 ⁹³	NTX 50 (56) Placebo (62)	26	Sweden; 10 sites outpatient	Ads, outpatient treatment center	48 to 51	NR	9 to 23	0	None	Low
Garbutt, 2005 ⁸³ , Pettinati, 2009 ¹¹⁷	NTX inj 380 (208) NTX inj 190 (210) Placebo (209)	26	U.S.; Inpatient and outpatient, private and VA	NR	45	17	32	NR	BRENDA standardized supportive therapy 100%	Medium
Morgenstern, 2012 ⁸⁷	NTX 100 + MBSCT (51) NTX 100 (51) Placebo + MBSCT (50) Placebo (48)	12	U.S.; NR	Ads, community outreach	40	26	0	HIV 15 Any drug use 67	BBCET 100%	Medium
O'Malley, 2008 ⁹⁹	NTX 50 (34) Placebo (34) NTX 50 + SER 100 (33)	16	U.S.; Outpatient	Direct community recruitment, health clinic referral, local ads	40	70	34	NR	MM 100%	Medium
Petrakis, 2004 ¹⁰⁷ , Ralevski, 2006 ¹²⁸	NTX 50 (16) Placebo (15)	12	U.S.; At least 3 outpatient centers—MIRECC clinics	Direct recruitment from participating centers	46	19	0	Schizophrenia or schizoaffective disorder 100	CBT + psychiatric treatment as usual Neuroleptics 52% Benzodiazepines 16% Thymoleptics 39%	Medium
Pettinati, 2008 ⁹¹	NTX 150 (82) Placebo (82) Subjects also randomized to either CBT or BRENDA (2x2 design)	12	U.S.; University-affiliated outpatient substance abuse treatment research facility	Those seeking treatment at the facility	39	76	29	Cocaine dependence 100	NR	Medium

Table 22. Characteristics of included double-blind randomized placebo-controlled trials of naltrexone that report a health outcome (continued)

Author, Year Trial Name	Arm Dose, mg/day (N)	Rx Duration, Weeks (Follow- up)	Setting	Recruitment Method	Age, Years	% Non- white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Pettinati, 2010 ⁸⁹	SER 200 (40) NTX 100 (49) Placebo (39) SER 200 + NTX 100 (42)	14	U.S.; Outpatient	Newspaper ads, referrals from local professional or friends / family	43	35	38	Depression 100	CBT 100%	Medium
Petrakis, 2005 ⁷⁶	DIS 250 (66)	12	U.S.; Outpatient VA	Recruited as outpatients or ads	47	26	3	Axis I disorder 100	Psychiatric treatment as usual 100%	Medium
Ralevski, 2007 ⁸⁰	NXT 50 (59)									
Petrakis, 2007 ⁸¹	Placebo (64)									
Petrakis, 2006 ⁸²	NTX 50 + DIS 250 (65)									
VA MIRECC										

^a Three additional treatment arms were included in COMBINE but were not relevant to our Key Questions: ACA + NTX + CBI + MM, ACA + NTX + MM, and CBI only (no pills).

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: AA = Alcoholics Anonymous; ACA = acamprosate; CBI = combined behavioral intervention; CBT = cognitive behavioral therapy; DIS = disulfiram; inj = injection; mg = milligram; MBSCT = modified behavioral self-control therapy; MIRECC = Mental Illness Research, Education and Clinical Center; MM = medical management; N = number; NR = not reported; NTX = naltrexone; SER = sertraline; U.S. = United States; VA = Veterans Affairs.

One trial was conducted in Sweden,⁹³ all others were conducted in the United States. Most were conducted at an outpatient substance abuse or mental health center; none were conducted in primary care settings.

Mortality

Six placebo-controlled trials of naltrexone reported mortality rates; no study found more than one death in each treatment group. Three studies reported that there were no deaths in either group,^{89,91,117} 1 reported one death in each study arm without providing additional details,⁷⁶ and 1 study reported a death due to alcohol intoxication in the placebo group.⁹³ In the COMBINE trial, the authors reported that one fatal serious adverse event was reported during the 16-week treatment phase. Investigators classified this death as being unrelated to the study medication. No details were provided on which group the death occurred in, the nature of the adverse event, or the cause of death.⁵³

Quality of Life or Function

Four placebo-controlled trials of naltrexone measured QoL or some aspect of function, each trial using a different measure. One trial conducted among men who have sex with men⁸⁷ measured QoL at 13 weeks using the Short Inventory of Problems,¹⁷⁷ an alcohol-specific QoL measure used to assess negative consequences of drinking. No differences between naltrexone and placebo in end-of-treatment scores were found when using a last observation carried forward (LOCF) method to impute missing data (mean difference between groups at 13 weeks was -1.7, $p < 0.09$).⁸⁷

One study comparing injectable naltrexone with placebo measured QoL using the Medical Outcomes Study 36-item short-form health survey (SF-36).^{117,178} Data were reported separately for the overall physical and overall mental health summary scores of the SF-36. The study found no significant difference on either scale at 24 weeks between the placebo group and the injectable naltrexone 190 mg per day group. Patients receiving naltrexone 380 mg per day had greater improvement on the mental health summary score than those receiving placebo at 24 weeks (8.2 versus 6.2, $p = 0.044$), but there was no difference in improvement found on the physical health summary score (0.2 versus -0.1, $p = 0.51$).¹¹⁷

The COMBINE study assessed QoL using the WHOQOL and SF-12v2 physical and mental health scores. Results were not presented for each treatment group separately.¹⁷⁶ See the section below on acamprosate versus naltrexone for details on these results. Briefly, the results indicate that the eight combinations of pharmacological and behavioral treatments did not show clinically significant differential effects on QoL for either scale.¹⁷⁶

One placebo-controlled study of naltrexone 50 mg measured the Drinker Inventory of Consequences (DrInC) at 16 weeks.⁹⁹ The DrInC is a 50-item questionnaire designed to measure adverse consequences of alcohol abuse in five areas: interpersonal, physical, social, impulsive, and intrapersonal.¹⁷⁷ More patients in the placebo group reported one or more alcohol-related consequence than in the naltrexone group, as measured by the DrInC (76 versus 45%, $p = 0.02$).⁹⁹

Detailed Synthesis: Placebo-Controlled Trials of Medications Used Off-label, or Those Under Investigation

As described in KQ 1, we found just 1 placebo-controlled trial meeting our inclusion criteria for each of the following medications: aripiprazole, atomoxetine, desipramine, fluvoxamine, imipramine, olanzapine, and paroxetine. We found insufficient evidence to support the efficacy

of these medications. Among these studies, just 1 reported a health outcome (number of deaths for fluvoxamine and placebo).¹⁷⁹ We provide additional details about this trial in Appendix ZZ.

For the medications with multiple placebo-controlled trials (baclofen, buspirone, citalopram, fluoxetine, nalmefene, quetiapine, sertraline, topiramate, and valproic acid), 4 trials reported outcomes relevant to KQ 2 (Table 23): 1 trial of quetiapine in bipolar patients with alcohol dependence,¹⁴⁷ 2 placebo-controlled trials of sertraline in patients with co-existing depression or dysthymia,^{89,151} and 1 placebo-controlled trial of topiramate.^{160,161} Sample size ranged from 83 to 371. Duration of treatment ranged from 12 to 24 weeks.

Table 23. Characteristics of included double-blind randomized placebo-controlled trials of medications used off-label, or those under investigation

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Stedman, 2010 ¹⁴⁷	Quetiapine 300-800 (175) Placebo (186)	12	U.S.; Outpatient; 43 centers	NR	39	12	37	Bipolar 100	None	High
Gual, 2003 ¹⁵¹	SER 50-150 (44) Placebo (39)	24	Spain; Outpatient	Outpatient alcohol dependence treatment	47	NR	47	Depression/dysthymia 100	NR	Medium
Pettinati, 2010 ⁸⁹	SER 200 (40) NTX 100 (49) Placebo (39) SER 200 + NTX 100 (42)	14	U.S.; Outpatient	Newspaper ads, referrals	43	35	38	Depression 100	CBT 100%	Medium
Johnson, 2007 ¹⁶⁰ Johnson, 2008 ¹⁶¹	TOP 50-300 ^a (183) Placebo (188)	14	U.S.; 17 academic sites; outpatient	Academic sites and by newspaper, radio, television ads	47 to 48	15	26 to 28	NR	BBCET 100%	Low

^a Dose titrated over a 5-week period from 25 to a maximum of 300 mg; mean 171.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: BBCET, brief behavioral compliance enhancement treatment; CBT, cognitive behavioral therapy; mg = milligram; N = number; NR, not reported; NTX = naltrexone; SER, sertraline; TOP, topiramate; U.S., United States.

The mean age of participants was similar across trials—39 to 48 years. Twenty-six to 47 percent of patients were female, and 12 to 35 percent of patients were nonwhite in the 3 trials reporting information on race; 1 did not report information on race.¹⁵¹ Two trials reported smoking status: 1 placebo-controlled trial of sertraline enrolled 14 to 20 percent smokers,⁸⁹ and the trial of quetiapine enrolled 56 percent smokers.¹⁴⁷ Three of the trials were conducted in those with alcohol dependence; the trial of topiramate did not specify the percentage who met criteria for alcohol dependence.¹⁶⁰ Three trials were conducted in the United States and 1 in Spain.¹⁵¹

The placebo-controlled trial of quetiapine was rated as high risk of bias, primarily for high risk of attrition bias and methods of handling of missing data (see Appendix C for details).¹⁴⁷

Topiramate

Accident or Injury

One placebo-controlled trial of topiramate reported injury in a list of adverse events occurring during treatment (over 12 weeks).¹⁶⁰ Eight patients (4.4 percent) in the topiramate

group and 22 patients in the placebo group (11.7 percent) had an injury ($p=0.01$). The authors note that three separate individuals in the placebo group experienced a tibial plateau fracture. No other information is provided on the cause or nature of the injuries.¹⁶⁰

Mortality

The placebo-controlled trial of topiramate reported one death in the placebo group following a cardiac arrest associated with gastrointestinal tract bleeding and seizures.¹⁶⁰ According to the investigators, the precipitating incident could not be determined. There was no mention of deaths in the topiramate group.¹⁶⁰

Quetiapine

Mortality

Two deaths (one in each treatment group) were reported in the placebo-controlled trial of quetiapine rated as high risk of bias; one after a skull fracture caused by blunt trauma in the quetiapine group and one attributed to myocardial ischemia more than 30 days after treatment in the placebo group.¹⁴⁷ Both deaths were judged to be unrelated to the study medications by the study investigators.

Quality of Life or Function

No difference was found between the quetiapine and placebo groups in health-related QoL assessed by the Quality of Life Enjoyment and Satisfaction Questionnaire (Q-LES-Q)¹⁸⁰ at 12 weeks (mean score 46.9 versus 47.7, $p=0.63$). Functional impairment was assessed using the Sheehan Disability Score (SDS),¹⁸¹ a questionnaire that aims to assess the relationship between symptoms and impairment in work, social, and family life. No statistically significant differences between quetiapine and placebo groups were found for mean total SDS score (11.03 versus 9.17), mean SDS number of lost work days per week (1.1 versus 0.7), and SDS number of underproductive days per week over 12 weeks (1.8 versus 1.3).¹⁴⁷

Sertraline

Mortality

One placebo-controlled trial of sertraline in patients with co-existing alcohol dependence and depression reported no deaths in either treatment group at 13 weeks.⁸⁹

Quality of Life or Function

One study of patients with co-existing depression measured QoL using the SF-36 at 24 weeks. Scores were presented in a figure only (bar graph, data not reported). QoL improved during treatment for both the placebo and sertraline groups; the authors noted that the sertraline group improved more than placebo in only the mental health summary score of the SF-36 ($p=0.031$).¹⁵¹

Head-to-head Trials Including FDA-Approved Medications

Detailed Synthesis: Head-to-Head Trials

We identified 3 RCTs (Table 24) that reported at least one health outcome of interest. Two of these were rated as high risk of bias for the head-to-head comparison—one three-arm study comparing naltrexone with disulfiram or placebo,⁷⁶ and one four-arm open-label trial comparing acamprosate, disulfiram, and naltrexone.¹⁶⁵ Both trials had high risk of ascertainment bias; one did not adequately handle missing data for QoL outcomes (see Appendix C for additional details about risk of bias ratings).

Table 24. Characteristics of head-to-head randomized controlled trials reporting a health outcome

Author, Year Trial Name Design	Arm Dose, mg/day (N)	Rx Dura- tion, Weeks (Fol- lowup)	Setting	Recruitment Method	Age, Years	% Non- white	% Fe- male	Co-inter- vention	Risk of Bias
Anton, 2006 ⁵³ LoCastro, 2009 ¹⁷⁶ COMBINE DBRCT	ACA 3,000 + CBI + MM (151) ACA 3,000 + MM (152) NTX 100 + CBI + MM (155) NTX 100 + MM (154) Placebo + CBI + MM (156) Placebo + MM (153) ^a	68	U.S.; 11 sites	Ads, community resources, clinical referrals at 11 academic sites	44	23	31	Community support group participation encouraged (e.g., AA)	Low
Laaksonen, 2008 ¹⁶⁵ OLRCT	ACA 1,998 or 1,333 (81) DIS 100 to 200 (81) NTX 50 (81)	Up to 52 (119)	Finland; 6 sites in 5 cities	Volunteers seeking outpatient treatment for alcohol problems	43	0	29	Manual-based CBT ^b	High for quality of life / KQ 2
Petrakis, 2005 ⁷⁶ Ralevski, 2007 ⁸⁰ Petrakis, 2007 ⁸¹ Petrakis, 2006 ⁸² VA MIRECC DBRCT	DIS 250 (66) NTX 50 (59) Placebo (64) NTX 50 + DIS 250 (65)	12	U.S.; Outpatient VA	Recruited as outpatients or via ads	47	26	3	Psychiatric treatment as usual 100%	High for DIS vs. NTX

^a Three additional treatment arms were included in COMBINE but were not relevant to our Key Questions: ACA + NTX + CBI + MM, ACA + NTX + MM, and CBI only (no pills).

^b Co-intervention included a “Winning at last--defeating the drinking problem” booklet targeted to match medication goals (i.e., reduction in drinking or abstinence for ACA and NTX; abstinence for DIS).

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: AA = Alcoholics Anonymous; ACA = acamprosate; CBI = combined behavioral intervention; DBRCT = double-blind randomized controlled trial; DIS = disulfiram; mg = milligram; MIRECC = Mental Illness Research, Education and Clinical Center; MM = medical management; N = number; NR = not reported; NTX = naltrexone; OLRCT = open-label randomized controlled trial; U.S. = United States; VA = Veterans Affairs.

One study (COMBINE), rated as low risk of bias, reported mortality and QoL. COMBINE is a multicenter nine-arm trial that compared eight groups of patients receiving medical management with 16 weeks of naltrexone (100 mg per day) or acamprosate (3 g per day), both, and/or both placebos, with or without a combined behavioral intervention (CBI). The ninth group received CBI only and no drug or placebo. Mean age was 44 years; all patients met criteria for alcohol dependence.

Acamprosate Versus Naltrexone

Mortality

In the COMBINE trial, the authors reported that one fatal serious adverse event was reported during the 16-week treatment phase. This was classified by investigators as not related to the study medication. No details were provided on which group the death occurred in, the nature of the adverse event, or the cause of death.

One study, rated as high risk of bias, reported that one person committed suicide and two persons drowned in the acamprosate group but reported no events in the naltrexone group.¹⁶⁵

Quality of Life or Functional Status

The COMBINE study assessed QoL using the WHOQOL and SF-12v2 physical and mental health scores. Results were not presented for each treatment group separately.¹⁷⁶ To analyze the treatment effects of specific pharmacological and behavior treatment combinations on QoL, a mixed-effects general linear model was used to examine the main and interaction effects of three treatments (acamprosate, naltrexone, and CBIs) from baseline to 26 weeks and from baseline to 52 weeks (20 ANOVAs were conducted unadjusted and 20 were adjusted for percentage heavy drinking days). The results indicate that the eight combinations of pharmacological and behavioral treatments did not show differential effects on QoL for either scale. The only two significant effects reaching a p value of <0.001 (to account for multiple tests) were the two-way interaction of naltrexone by CBI for the SF-12v2 physical health score at 52 weeks for both the adjusted and unadjusted analyses. The authors conclude that this suggests CBI and naltrexone combined have a greater impact than either alone for the SF-12v2 physical health scale; however, the difference between groups was no larger than 2.1, and unlikely to suggest a clinically meaningful difference (the 95% confidence interval for the SF-12v2 physical health scale is 6.6).¹⁷⁶

One study rated as high risk of bias measured QoL with the European Quality of Life Scale (EQ-5),¹⁸² Koskenvuo Quality of Life Scale (KQL),¹⁸³ and Visual Analogue Scale (VAS).¹⁸⁴ QoL improved for both groups over the 52-week followup compared with baseline with no difference between the acamprosate or naltrexone groups.¹⁶⁵

Acamprosate Versus Disulfiram

Accident or Injury

One study, rated as high risk of bias, reported one traffic accident in the disulfiram group and none in the acamprosate group over 52 weeks.¹⁶⁵ No details of the event were described; the study coordinator determined that the event was not related to the study treatment.

Mortality

One study, rated as high risk of bias, reported that one person committed suicide and two persons drowned in the acamprosate group and reported no events in the disulfiram group.¹⁶⁵

Quality of Life

QoL was measured in one study rated high risk of bias with the EQ-5, KQL, and VAS. QoL improved for both groups over the 52-week followup compared with baseline with no difference between the acamprosate or disulfiram groups.¹⁶⁵

Disulfiram Versus Naltrexone

Accident or Injury

One study, rated as high risk of bias, reported one traffic accident in the disulfiram group and no accident or injuries in the naltrexone group.¹⁶⁵ No details of the event were described; the study coordinator determined that the event was not related to the study treatment.

Mortality

In 1 study rated high risk of bias that compared disulfiram and naltrexone among patients with co-existing depression, one person died in the naltrexone group and no deaths were reported in the disulfiram group.⁷⁶

Quality of Life

QoL was measured in 1 study rated high risk of bias with the EQ-5, KQL, and VAS. QoL improved for both groups over the 52-week followup compared with baseline with no difference between the disulfiram or naltrexone groups.¹⁶⁵

Head-to-Head Trials Including Medications Used Off-label, or Those Under Investigation

Characteristics of Trials

We identified 3 head-to-head trials of off-label medications that measured an eligible health outcome (

Table 25). One compared sertraline with naltrexone; 2 compared topiramate with naltrexone. Sample size ranged from 89 to 182 within the relevant head-to-head arms. All subjects met criteria for alcohol dependence, the average age of participants was similar across trials (43 to 48), and females made up 15 to 38 percent of participants. The trial comparing sertraline with naltrexone was conducted among patients with co-occurring depression.⁸⁹ The trials comparing topiramate with naltrexone enrolled about a quarter of subjects with personality disorders.^{185,186} Only the study comparing sertraline with naltrexone reported on smoking rates: 14 to 20 percent of participants were smokers. All studies included a psychological co-intervention.

One double-blind RCT compared sertraline 200 mg per day with naltrexone 100 mg per day⁸⁹ and 2 open-label RCTs compared topiramate 200 mg per day to naltrexone 50 mg per day.^{185,186} One study was conducted within the United States⁸⁹ and 2 were conducted in Spain.^{185,186} The trial comparing sertraline to naltrexone was rated as medium risk of bias and the 2 studies comparing topiramate to naltrexone were rated as high risk of bias.^{185,186} One study allowed titration of topiramate from 200 mg per day up to 300 to 400 mg per day based on continued alcohol consumption or craving.¹⁸⁵

Table 25. Characteristics of head-to-head randomized controlled trials including medications used off-label, or those under investigation

Author, Year Design	Arm Dose, mg/day (N)	Rx Duration, Weeks	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Pettinati, 2010 ⁸⁹ DBRCT	SER 200 (40) NTX 100 (49) Placebo (39) SER 200 + NTX 100 (42)	14	U.S.; Outpatient	Newspaper ads, referrals	43	35	38	Depression 100	CBT 100%	Medium
Florez, 2008 ¹⁸⁵ OLRCT	TOP intended 200 ^a (51) NTX 50 (51)	26	Spain; Outpatient substance abuse clinic, referrals	Recruited when presenting for treatment	47	0	15	Personality disorders 27	Therapy based on Relapse Prevention Model 100%	High
Florez, 2011 ¹⁸⁶ OLRCT	TOP 200 (91) NTX 50 (91)	26	Spain; Outpatient substance abuse clinic, referrals	Recruited and screened when presenting for treatment	47 to 48	NR	15	Personality disorders 23	BRENDA 100% At least monthly meeting with psychiatrist 100%	High

^aActual dosing: increased by 50 mg per day up to 300 or 400 mg based on consumption control or cravings.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: CBT = cognitive behavioral therapy; DBRCT = double-blind randomized controlled trial; mg = milligram; N = number; OLRCT = open-label randomized controlled trial; U.S., United States.

Sertraline Versus Naltrexone

Mortality

One trial comparing sertraline with naltrexone among patients with co-occurring depression and alcohol dependence reported no deaths in either group.⁸⁹

Topiramate Versus Naltrexone

Quality of Life or Function

One unblinded study rated as high risk of bias used the World Health Organization Psychiatric Disability Assessment Schedule (WHO/DAS) to assess alcohol dependence-related disability at 3 and 6 months.¹⁸⁶ No significant changes were found in most domains of the WHO/DAS at 3 months (personal, family, social), with one exception: patients taking topiramate had a lower disability score on the employment domain (1.64 versus 2.2, $p=0.047$). At 6 months, the topiramate group had lower disability scores for the family (0.58 versus 1.05, $p=0.035$) and social domains (0.46 versus 0.83, $p=0.154$); there was no difference between the two groups in the employment or personal domains at 6 months.¹⁸⁶ A similar study (by the same author), which dosed topiramate based on continued alcohol intake or craving, found no difference between the topiramate and naltrexone groups on any of the WHO/DAS domains at 3 or 6 months.¹⁸⁵

This same study measured QoL using the EQ-5D at 3 and 6 months.¹⁸⁶ At 3 months, the topiramate group had a small, but statistically significant, greater improvement in QoL compared with the naltrexone group (96.10 versus 94.16, $p=0.014$); there was no difference between the two groups at 6 months.¹⁸⁶ A similar study (by the same author), which dosed topiramate based

on continued alcohol intake or craving, found that patients treated with topiramate had better QoL at 3 months compared with naltrexone (96.88 versus 95.21, $p=0.014$) but no statistically significant difference was found between the two groups at 6 months.¹⁸⁵

Key Question 3. Adverse Effects of Medications

For this question, we evaluated trials included in Key Questions (KQs) 1 and 2. In addition, we searched for nonrandomized controlled trials (non-RCTs), open-label trials, single-blind trials, prospective cohort studies, and case-control studies otherwise meeting the eligibility criteria. We ultimately included 96 double-blind RCTs, eight open-label or single-blind RCTs, and one prospective cohort study. Throughout this KQ, we often describe risks of various adverse events—risks reported are absolute risk differences (RDs) between intervention and control. Because the studies were not primarily focused on harms, the reporting of harms varied across studies significantly. Limited information was reported for most of the off-label medications—insufficient for synthesis of specific adverse events or for making definitive conclusions. We therefore focus here on the FDA-approved medications and those with moderate or better evidence supporting efficacy. We do not include information on medications with insufficient evidence to support their efficacy (i.e., efficacy as determined in KQ 1).

Key Points

- Adverse events were often not collected using standardized measures, and methods for systematically capturing adverse events were often not reported.
- Selective outcome reporting could impact our results. Reporting varied across studies, with some studies only reporting adverse effects that were significantly different from placebo (or control) group, some reporting effects observed in more than some percentage of patients (e.g., 5 percent or more), and others listing effects that were considered in the study.
- While major harms were rarely reported in the studies, some minor harms (e.g., diarrhea) were reported more consistently.
- For many serious harms, the evidence was insufficient to determine comparative rates of adverse events — very little data were available.
- **Suicidality, or self-harmful behaviors:** evidence was insufficient to determine whether risk was increased with any of the medications. Overall evidence was limited to three cases of suicide attempts or suicidal ideation reported in acamprosate arms and three in placebo arms.
- **Withdrawals due to adverse events:** In head-to-head studies, the risk of withdrawals due to adverse events was not significantly different between acamprosate and naltrexone.
- **Specific adverse events:** Compared with placebo, patients treated with acamprosate had a higher risk of anxiety, diarrhea, and vomiting. Compared with placebo, those treated with naltrexone had a higher risk of dizziness, nausea, and vomiting. In head-to-head studies, patients treated with acamprosate had a slightly lower risk of headache than those treated with naltrexone.

Detailed Synthesis

In this section, we have considered harms associated with acamprosate, disulfiram, and naltrexone. Our main meta-analyses included studies of low and medium risk of bias reporting results for the specific adverse event. We conducted sensitivity analyses that also included studies rated as high or unclear risk of bias. Insufficient data were available to conduct meta-analyses of results from studies that compared disulfiram with placebo, acamprosate, naltrexone, or other controls. Therefore, we described and summarized these qualitatively when possible.

Characteristics of Included Studies

The vast majority of the included RCTs are described in KQs 1 and 2, and we do not describe them again in this KQ. Nine studies not described in KQs 1 or 2 were eligible for inclusion in this KQ. These included 7 open-label^{164,166,167,185-187} or single-blind RCTs,¹⁸⁸ 1 double-blind RCT,^{189,190} and 1 prospective cohort study.¹⁹¹ Of those 9, 5 focused on comparisons addressed in this KQ (Table 26); the other 4 focused on comparisons with medications used off-label—either topiramate¹⁸⁵⁻¹⁸⁷ or buspirone.^{189,190} All of the studies listed in Table 26 were rated as high risk of bias, primarily due to concerns with selection bias, attrition bias, measurement bias, confounding, or selective outcome reporting bias (see Appendix C for details).

Table 26. Characteristics of studies included for KQ 3 that were not in KQ 1 or 2

Author, Year Design	Arm Dose, mg/day (N)	Rx Duration, Weeks	Country Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Additional Condition	Co-intervention	Risk of Bias
Narayana, 2008 ¹⁹¹ Prospective cohort	ACA 1,332 to 1,998 (28) NTX 50 (26) TOP 100 to 125 (38)	52	India; military, outpatient	Members of the Armed Forces	38	100	0	NR	Various psychotherapies were offered	High
Nava, 2006 ¹⁶⁷ OLRCT	GHB 50 ^a (28) NTX 50 (24) DIS 200 (28)	52	Italy; outpatient	Advertisements, word of mouth, press release	38.5 to 42.7	NR	15%	0	Cognitive behavioral therapy	High
De Sousa, 2005 ¹⁶⁴ OLRCT	ACA 1,998 (50) DIS 250 (50)	35	India; outpatient, private psychiatric hospital	Patients undergoing detoxification	42 to 43	100	0	NR	Weekly supportive group psychotherapy offered	High
De Sousa, 2004 ¹⁶⁶ OLRCT	DIS 250 (50) NTX 50 (50)	52	India; outpatient	Recruited as inpatients	43 to 47	NR	0	NR	Supportive group psychotherapy	High
Rubio, 2001 ¹⁸⁸ SBRCT	ACA 1,665-1,998 (80) NTX 50 (77)	52	Spain; outpatient	Patients presenting to hospital for detoxification	44	NR	0	0	Supportive group therapy weekly; weekly visits with a psychiatrist for 3 months, then biweekly until end of study	High

^a Dose is 50 mg per kg of body weight 3 times a day.

Notes: Age (y) is the mean age in years, unless otherwise stated.

The following studies also met the inclusion criteria, but assessed harms of an off-label medication (compared with placebo) without evidence of efficacy, or compared an off-label medication without evidence of efficacy with an FDA-approved medication, and are therefore not described further in this Key Question: Florez, 2011,¹⁸⁶ Florez, 2008¹⁸⁵, De Sousa, 2008,¹⁸⁷ and Tollefson, 1991.^{189,190}

Abbreviations: ACA = acamprosate; DIS = disulfiram; FDA = U.S. Food and Drug Administration; GHB = γ -Hydroxybutyric acid; mg = milligrams; N = number; NR = not reported; NTX = naltrexone; OLRCT = open-label randomized controlled trial; SBRCT = single-blind randomized controlled trial; TOP = topiramate; U.S. = United States.

For the 5 studies not described elsewhere that focused on comparisons addressed in this KQ, 2 compared acamprosate with naltrexone,^{188,191} 2 compared naltrexone with disulfiram,^{166,167} and 1 compared acamprosate with disulfiram.¹⁶⁴ Study duration ranged from 35 to 52 weeks. Three of the studies were conducted in India, 1 in Spain, and 1 in Italy. For 3 of the 4 trials, study participants were recruited as inpatients. For the other trial, recruitment methods included advertisements, word of mouth, and a press release.¹⁶⁷ The prospective cohort study¹⁹¹ followed members of the armed forces. Mean age ranged from 38 to 47 years. Only 1 of the studies included women.¹⁶⁷ In 2 studies,^{164,191} all participants were nonwhite; race and ethnicity was not reported in the other 3 studies.

Acamprosate Compared With Placebo

Table 27 summarizes the results of our meta-analyses. The only statistically significant findings for harms from our main analyses were for anxiety, diarrhea, and vomiting. Statistical heterogeneity was considerable for the diarrhea analysis (I^2 92.7 percent), with some studies finding much higher rates of diarrhea than with placebo (with absolute risks increased as much as 33 percent). Sensitivity analyses for withdrawals due to adverse events, anxiety, diarrhea, and vomiting were also statistically significant (finding higher risk with acamprosate).

Table 27. Results of meta-analyses for adverse events: acamprosate compared with placebo

Outcome	N trials	N subjects	RD	95% CI	Heterogeneity I^2
Withdrawal due to adverse events	11	4,069	0.007	-0.003 to 0.017	18.5%
Withdrawal due to adverse events—SA	14	4,833	0.008	0.001 to 0.016	0.0%
Anxiety	1	821	0.23	0.17 to 0.29	NA
Anxiety—SA	2	844	0.19	0.06 to 0.32	47.2%
Diarrhea	11	3,264	0.090	0.019 to 0.160	92.7%
Diarrhea—SA	13	4,083	0.084	0.026 to 0.142	90.5%
Dizziness	0	0	NA	NA	NA
Headache	5	1,039	-0.003	-0.059 to 0.053	73.7%
Headache—SA	6	1,608	-0.000	-0.044 to 0.044	66.1%
Insomnia	1	116	0.04	-0.03 to 0.106	NA
Insomnia—SA	2	685	0.039	-0.009 to 0.086	0.0%
Nausea	5	1,623	0.01	-0.01 to 0.03	0.0%
Nausea—SA	6	1,693	0.01	-0.01 to 0.03	0.0%
Numbness	1	262	0.008	-0.013 to 0.029	NA
Numbness—SA	2	831	0.010	-0.010 to 0.029	0.0%
Rash	1	246	-0.008	-0.030 to 0.014	NA
Rash—SA	2	316	0.016	-0.065 to 0.097	66.9%
Suicide attempts or suicidal ideation	1	581	0.007	-0.005 to 0.019	NA
Suicide attempts or suicidal ideation—SA	3	1,173	0.002	-0.008 to 0.011	14.6%
Vomiting	3	1,782	0.02	0.01 to 0.04	0.0%
Vomiting – SA	4	1,805	0.02	0.01 to 0.04	3.5%

Note: Positive risk differences favor placebo. Sensitivity analyses include studies rated as high risk of bias.

Abbreviations: CI = confidence interval; N = number of trials or subjects contributing data; NA = not applicable; RD = risk difference; SA = sensitivity analysis.

Disulfiram Compared With Placebo or Control

Four included studies compared disulfiram with placebo or control.⁷⁶⁻⁷⁹ One of these did not report results for adverse events.⁷⁹ The other 3 did not yield sufficient quantitative data to conduct meta-analyses.

One study of disulfiram compared with placebo in patients who were all taking methadone reported that “there were no deaths, serious adverse reactions, or illnesses that could be attributed to the combined use of the drugs [disulfiram and methadone]” but did not provide details about the incidence of specific adverse events in the study population.⁷⁷

In another study, patients who received 250 mg per day of disulfiram reported “moderate or severe” drowsiness more often than those not given disulfiram (8 versus 2 percent, $p=0.03$). There was no significant difference in the incidence of drowsiness between the 250 and 1 mg per day disulfiram groups.⁷⁸ In this same study, disulfiram was discontinued by 3 patients in the 250 mg per day group and one patient in the 1 mg per day group because of increased serum alkaline phosphatase or aspartate aminotransferase. Psychiatric problems were observed in 11 patients with no statistically significant difference between the three groups.⁷⁸

Results from a four-arm study comparing disulfiram combined with naltrexone, disulfiram combined with placebo, naltrexone alone, and placebo alone showed that patients on any study medication experienced aftertaste, blurred vision, confusion, constipation, drowsiness, dry mouth, loss of appetite, nausea, or tremors more often than patients who received placebo. There were no statistically significant between-group differences for other adverse events.⁷⁶ Six of the 14 serious adverse events reported in this study occurred in the disulfiram with placebo group (4 psychiatric hospitalizations—2 for a change in mental status and 2 for suicidal ideation, 1 cardiac event, and 1 hospitalization for acute axonal neuropathy) and 3 occurred in the placebo group (1 death, 1 drug and alcohol overdose, and 1 hospitalization for pneumonia).⁷⁶

Naltrexone Compared With Placebo

Table 28 summarizes the results of our meta-analyses. We found statistically significant increased risk of withdrawal due to adverse events, dizziness, nausea, and vomiting.

Acamprosate Compared With Disulfiram

Both studies reporting results for adverse events for this comparison were rated as high risk of bias; both reported no statistically significant differences between the acamprosate and disulfiram groups.^{164,165}

One of the studies reported that six patients who received disulfiram experienced elevated alanine transaminase (ALAT) levels. Subsequently, three of the patients discontinued the medication, and three continued to receive a half dose; ALAT levels normalized within 2 to 3 weeks.¹⁶⁵ The most common adverse events reported in the study for patients treated with acamprosate were diarrhea and dermatological problems; for patients treated with disulfiram—tiredness and headache.

Table 28. Results of meta-analyses for adverse events: naltrexone compared with placebo

Outcome	N trials	N subjects	RD	95% CI	Heterogeneity I ²
Withdrawal due to adverse events	14	2,203	0.02	0.01 to 0.04	0.0%
Withdrawal due to adverse events - SA	16	2,319	0.02	0.01 to 0.04	0.0%
Anxiety	5	725	0.03	-0.01 to 0.08	0.0%
Anxiety—SA	7	940	0.03	-0.01 to 0.07	0.0%
Diarrhea	9	2,232	0.011	-0.018 to 0.041	34.8%
Diarrhea - SA	10	2,335	0.004	-0.025 to 0.034	40.8%
Dizziness	11	2,549	0.068	0.037 to 0.099	46.7%
Dizziness - SA	15	2,851	0.062	0.037 to 0.088	35.7%
Headache	14	3,102	0.010	-0.020 to 0.039	15.9%
Headache - SA	19	3,554	0.007	-0.024 to 0.038	27.1%
Insomnia	6	1,571	0.015	-0.016 to 0.046	0.0%
Insomnia - SA	10	1,964	0.018	-0.008 to 0.044	0.0%
Nausea	22	4,320	0.11	0.07 to 0.16	72.5%
Nausea - SA	30	4,928	0.10	0.07 to 0.14	68.5%
Numbness	1	246	0.032	-0.093 to 0.157	NA
Numbness - SA	2	410	-0.015	-0.108 to 0.078	46.0%
Rash	2	134	0.056	-0.128 to 0.241	44.6%
Rash - SA	3	187	0.006	-0.069 to 0.081	19.8%
Suicide	0	0	NA	NA	NA
Blurred vision	2	133	0.079	-0.172 to 0.331	46.3%
Blurred vision - SA	NA	NA	NA	NA	NA
Vomiting	7	2,103	0.05	0.03 to 0.08	0.0%
Vomiting - SA	9	2,232	0.04	0.02 to 0.07	5.1%

Note: Positive risk differences favor placebo. Sensitivity analyses include studies rated as high risk of bias.

Abbreviations: CI = confidence interval; N = number of trials or subjects contributing data; NA = not applicable; RD = risk difference; SA = sensitivity analysis.

Acamprosate Compared With Naltrexone

Table 29 summarizes the results of our meta-analyses. The only trial rated as low or medium risk reporting headache did not find a statistically significant difference between treatments, but findings trended in favor of acamprosate. Our meta-analysis including high risk of bias studies found that patients treated with acamprosate had a lower risk of headache than those treated with naltrexone.

Table 29. Results of meta-analyses for adverse events: acamprosate compared with naltrexone

Outcome	N trials	N subjects	RD	95% CI	Heterogeneity I ²
Withdrawal due to adverse events	1	612	-0.01	-0.04 to 0.02	NA
Withdrawal due to adverse events—SA	2	769	-0.01	-0.04 to 0.01	0.0%
Diarrhea	3	800	0.20	-0.02 to 0.42	92.8%
Diarrhea—SA	4	957	0.16	-0.09 to 0.40	97.1%
Dizziness	1	108	-0.02	-0.08 to 0.04	NA
Dizziness—SA	2	270	-0.09	-0.28 to 0.09	89.8%
Headache	1	108	-0.06	-0.15 to 0.03	NA
Headache—SA	3	427	-0.10	-0.17 to -0.03	39.8%
Nausea	3	800	-0.05	-0.12 to 0.03	59.8%
Nausea—SA	5	1,119	-0.08	-0.15 to 0.00	68.0%

Note: Positive risk differences favor naltrexone. Table only includes rows for outcomes with sufficient data for meta-analyses.

Abbreviations: CI = confidence interval; N = number of trials or subjects contributing data; NA, not applicable; RD = risk difference; SA = sensitivity analysis.

A prospective cohort study rated as high risk of bias comparing acamprosate with naltrexone reported that adverse events were uncommon, mild, and temporary in both groups. The most common adverse events in the naltrexone group (N=26) were anxiety (23.07 percent),

nervousness (23.07 percent), and insomnia (15.4 percent); these were not reported in the acamprosate group. The most common adverse events in the acamprosate group (N=28) were nausea (25.0 percent) and diarrhea (21.42 percent); 11 percent of those in the naltrexone group experienced nausea, but none reported diarrhea.¹⁹¹

Disulfiram Compared With Naltrexone

We found 4 studies comparing disulfiram with naltrexone and reporting on adverse events; all 4 were rated as high risk of bias.^{76,165-167} One of these reported no statistically significant difference in the incidence of adverse events between groups;¹⁶⁵ another stated that no serious adverse events occurred during the study and reported the incidence of adverse events only among those who withdrew because of adverse events.¹⁶⁷

In 1 of the studies, nausea, drowsiness, abdominal pain, and diarrhea were more common among patients receiving naltrexone than among those receiving disulfiram, but statistical significance was not reported.¹⁶⁶

A four-arm study comparing disulfiram combined with naltrexone, disulfiram combined with placebo, naltrexone alone, and placebo alone found that fever was more common in the disulfiram group than in the naltrexone group ($p=0.03$); nervousness ($p=0.005$) and restlessness ($p=0.03$) were more common in the naltrexone group than in the disulfiram group.⁷⁶

Key Question 4. Evidence from Primary Care Settings

Characteristics of Included Trials

We identified no eligible trials conducted completely in primary care settings. One included trial compared targeted nalmefene with placebo for 28 weeks in 15 sites in Finland—5 of the sites were specialist treatment clinics, 6 were private general practice offices, 2 were offices for occupational health, and 2 were specialized in conducting outpatient clinical research (Table 30).¹⁴² For targeted nalmefene dosing, patients were instructed to take the medication when they believed drinking to be imminent, rather than as a daily scheduled medication. The trial reported that 93 percent of subjects met criteria for alcohol dependence. It did not report information on smoking history at baseline. The study did not include any formal manualized psychosocial treatment, but did include some elements of BRENDA,¹⁹² including biopsychosocial assessment, feedback to subjects about assessments, simple advice to reduce drinking, and monitoring of treatment progress—with the emphasis on correct use of the study medication.

Several other published studies, including some in other sections of this report, may have implications for or some applicability to primary care settings, an issue addressed in the report Discussion.

Results for Consumption Outcomes

The trial found no significant difference in percentage of drinking days between nalmefene and placebo (WMD, -3.8; 95% CI, -9.3 to 1.7), but reported a lower percentage of heavy drinking days for patients treated with targeted nalmefene (18.1 percent versus 29.7 percent, $p=0.024$) and 1 fewer drinks per drinking day for patients treated with nalmefene (WMD, -1.0; 95% CI, -2.0 to -0.02) than for those who received placebo.

Table 30. Characteristics of included randomized controlled trials of FDA-approved medications for treating alcohol dependence in primary care settings

Author, Year	Arm Dose, mg/day (N)	Rx Duration, Weeks (Followup)	Setting	Recruitment Method	Age, Years	% Non-white	% Female	% With Co-occurring Condition	Co-intervention	Risk of Bias
Karhuvaara, 2007 ¹⁴²	Nalmefene 10 to 40 targeted dose (242) Placebo (161)	28 (52) ^a	Finland; 15 sites	Mainly by newspaper ads	49	0	19	NR	Some elements of BRENDA	Medium

^aAfter 28 weeks, nalmefene responders were invited to continue in a double-blind randomized controlled trial for an additional 24 weeks.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: FDA = U.S. Food and Drug Administration; mg = milligram; N = number; NR = not reported; U.S. = United States.

Key Question 5. Subgroups

We evaluated evidence on whether any of the medications were more or less effective than other medications for the following subgroups: men or women, older adults, young adults, racial or ethnic minorities, smokers, or those with co-occurring disorders. Only studies that compared at least two medications with each other were eligible for this Key Question (KQ). Throughout this KQ, we include headers and sections only for subgroups reported by the included studies.

Detailed Synthesis

Characteristics of Included Studies

Eleven RCTs and 1 observational study addressed this KQ (

Table 31). Studies included FDA-approved (acamprosate, disulfiram, naltrexone, topiramate) and non-FDA-approved (desipramine, paroxetine, sertraline, topiramate) medications. Treatment durations ranged from 12 weeks (7 studies) to 68 weeks. All but 1 of the studies reported concurrent psychiatric care, psychotherapy, or other psychosocial support. Studies were conducted in Australia, Brazil, Germany, and India in addition to the United States.

Mean age ranged from 32 to 47, the reported proportion nonwhite ranged from 23 to 100 percent, and the reported proportion female ranged from 0 to 72 percent. In 11 of the studies, all participants had alcohol dependence; in 1 it was not reported. Smoking rates were high (55 to 81 percent of participants) in 3 studies;^{49,55,66,105,114} all patients in 1 study¹⁹³ had cocaine dependence. Three studies included only participants with psychiatric comorbidities (Axis I disorders, depression, or PTSD).^{76,80-82,89,170} Participants were recruited from the community as well as from outpatient and inpatient contacts. Three of these studies were rated low risk of bias, 3 were rated medium, and the rest were rated high risk of bias, primarily due to concerns with attrition bias, inadequate handling of missing data, or measurement bias (see Appendix C for details).

Table 31. Characteristics of head-to-head medication studies that evaluated subgroups

Author, Year Trial Name Design	Arm Dose, mg/day (N)	Subgroup(s)	Rx Dura- tion, Weeks	Setting	Age, Years	% Non- white	% Fe- male	Co-intervention	Risk of Bias
Greenfield, 2010 ¹⁹⁴ Fucito, 2012 ⁵⁵ COMBINE DBRCT	ACA 3,000 + CBI + MM (151) ACA 3,000 + MM (152) NTX 100 + CBI + MM (155) NTX 100 + MM (154) Placebo + CBI + MM (156) Placebo + MM (153)	Men/women; smokers	68	11 U.S. academic sites	44	23	31	As randomized; community support group participation (like AA) encouraged	Low
Carroll, 1993 ¹⁹³ NA OLRCT	DIS 250 (9) NTX 50 (9)	Cocaine dependence	12	U.S.; outpatient substance abuse	32	39	72	Weekly individual psychotherapy 100%	High
De Sousa, 2004 ¹⁶⁶ NA OLRCT	DIS 250 (50) NTX 50 (50)	Men	52	India; outpatient	43 to 47	NR	0	Supportive group psychotherapy 100%	High
De Sousa, 2005 ¹⁶⁴ NA OLRCT	ACA 1,998 (50) DIS 250 (50)	Men	35	India; outpatient; private psychiatric hospital	42 to 43	100	0	Weekly supportive group psychotherapy offered	High
Kiefer, 2005 ⁶³ NA DBRCT	ACA 1,998 (40) NTX 50 (40) Placebo (40) ACA 1,998 + NTX 50 (40)	Somatic distress, depression, anxiety	12	Germany; 1 site, outpatient	46	NR	26	Group therapy	Low
Morley, 2006 ⁴⁹ Morley, 2010 ⁶⁶ NA DBRCT	ACA 1,998 (55) NTX 50 (53) Placebo (61)	Depression	12	Australia; 3 treatment centers with “medical care typically available at hospital based drug and alcohol treatment services”	45	NR	30	All offered 4-6 sessions of manualized compliance therapy Up-take / attendance NR	Low
Narayana, 2008 ¹⁹¹ Prospective cohort	ACA 1,332 to 1,998 (28) NTX 50 (26) TOP 100 to 125 (38)	Men	52	Indian military, outpatient	38	100	0	NR	High
Petrakis, 2005 ⁷⁶ Ralevski, 2007 ⁸⁰ Petrakis, 2007 ⁸¹ Petrakis, 2006 ⁸² VA MIRECC	DIS 250 (66) NTX 50 (59) Placebo (64) NTX 50 + DIS 250 (65)	Axis I disorders	12	U.S.; outpatient VA	47	26	3	Psychiatric treatment as usual 100%	High for DIS vs. NTX

Table 31. Characteristics of head-to-head medication studies that evaluated subgroups (continued)

Author, Year Trial Name Design	Arm Dose, mg/day (N)	Subgroup(s)	Rx Dura- tion, Weeks	Setting	Age, Years	% Non- white	% Fe- male	Co-intervention	Risk of Bias
Petrakis, 2012 ¹⁷⁰ NA DBRCT	DES 200 + placebo (24) PAR 40 + placebo (20) PAR 40 + NTX 50 (22) DES 200 + NTX 50 (22)	PTSD, depression	12	U.S.; outpatient; multiple mental illness centers, most subjects from VAs	47	25	9	Clinical management/ compliance enhancement therapy 100%	High
Pettinati, 2010 ⁸⁹ NA DBRCT	SER 200 (40) NTX 100 (49) Placebo (39) SER 200 + NTX 100 (42)	Depression	14	U.S.; outpatient	43	35	38	CBT 100%	Medium
Baltieri, 2008 ¹⁰⁵ ; Baltieri, 2009 ¹¹⁴ NA DBRCT	TOP target 200, maximum 400 (52) NTX 50 (49) Placebo (54)	Smokers	12	Brazil; outpatient	44 to 45	29	0	Psychosocial 100%	High
De Sousa, 2008 ¹⁸⁷ NA OLRCT	TOP 150 (50) DIS 250 (50)	Men	39	India; center with facilities for both in- and outpatient treatment of alcohol dependence and substance abuse	43	100	0	Offered weekly supporting group psychotherapy – % NR	High

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: AA = Alcoholics Anonymous; ACA = acamprosate; CBI = combined behavioral intervention; CBT = cognitive behavioral therapy; DBRCT = double-blind randomized controlled trial; DES = desipramine; DIS = disulfiram; mg = milligram; MIRECC = Mental Illness Research, Education and Clinical Center; MM = medical management; N = number; NR = not reported; NTX = naltrexone; OLRCT = open-label randomized controlled trial; PAR = paroxetine; PTSD = post-traumatic stress disorder; SER = sertraline; TOP = topiramate; U.S. = United States; VA = Veterans Affairs.

Sex

Five studies—4 trials and 1 prospective cohort—provided evidence about the effectiveness of medications by sex.^{164,166,187,191,194}

Subgroup analyses from the COMBINE study,¹⁹⁴ the only study among this group rated as low risk of bias, found no significant association between sex and the impact of acamprosate or naltrexone treatment on percentage of days abstinent, time to heavy drinking, or percentage of heavy drinking days.

Three trials, all open-label and from the same group of investigators, and all rated as high risk of bias, found that naltrexone and topiramate have a greater impact than disulfiram and disulfiram has a greater impact than acamprosate on reducing drinking for men.^{164,166,187}

The prospective cohort study, rated as high risk of bias, found that treatment with topiramate had a greater impact than acamprosate or naltrexone on any drinking for men.¹⁹¹

Smokers

Two studies provided evidence about the effectiveness of medications by smoking status. Subgroup analyses from the COMBINE study⁵⁵ found that smokers who received naltrexone had more days abstinent (78 percent versus 72 percent, $p=0.004$) and fewer heavy drinking days (14 percent versus 20 percent, $p=0.003$) than smokers who received placebo. No data were reported on the effectiveness of acamprosate among smokers—only that smokers did not benefit differentially from acamprosate. Subgroup analyses from a trial comparing naltrexone, topiramate, and placebo found no association between the number of cigarettes smoked per day at the start of the trial and the effect of naltrexone or topiramate on any drinking outcomes.^{105,114}

People with Co-occurring Disorders

Six studies provided evidence about the effectiveness of medications on individuals with co-occurring psychiatric disorders or other substance use disorders. Five studies addressed co-occurring psychiatric disorders, including depression or anxiety,^{49,66,89} Axis I disorders,^{76,80-82} PTSD and depression,¹⁷⁰ and somatic distress/depression/anxiety;⁶³ 1 addressed co-occurring cocaine dependence.¹⁹³ Two were rated low risk of bias, 1 as medium risk of bias, and 3 as high risk of bias. Four were conducted in the United States, 1 in Germany,⁶³ and 1 in Australia.^{49,66}

The German study addressing patients with co-occurring somatic distress/depression/anxiety⁶³ evaluated the effects of naltrexone and acamprosate in patients with scores above and below the median on the Symptom Checklist-90 (SCL-90) and its subscales.¹⁹⁵ In patients with total SCL-90 scores above the median, naltrexone was associated with a longer time to lapse compared with acamprosate (51.3 versus 30.1 days, p NR). Similar differences between naltrexone and acamprosate were found for the above-median scores for somatic distress (45.5 versus 20.3), depression (53.4 versus 28.1), and anxiety (47.3 versus 24.4), though none reached statistical significance. Results for time to relapse were similar, and were not statistically significantly different.

In 1 U.S.-based study of patients with co-occurring alcohol dependence and depression,⁸⁹ patients treated with naltrexone reported numerically longer time to relapse than patients treated with sertraline (45.2 versus 39.9 days, p NR). A slightly higher percent of patients treated with sertraline (27.5 percent) remained abstinent during treatment compared with naltrexone (21.3 percent, p NR).

Another U.S.-based study compared disulfiram (plus placebo) with naltrexone in a population of veterans with comorbid Axis I disorders.⁷⁶ There were no significant differences between disulfiram and naltrexone in percentage of days abstinent (97 percent versus 95 percent, respectively, $p=0.55$), percentage of heavy drinking days (3.2 percent versus 4.0 percent, $p=0.65$), or percentage remaining abstinent (77.3 percent versus 64.4 percent, $p=0.11$). This study was rated high risk of bias. When subgroups of the Axis I disorders were examined, results were similar, with no significant differences in alcohol use outcomes by treatment for patients diagnosed with depression,⁸¹ borderline personality disorder,⁸⁰ antisocial personality disorder,⁸⁰ or post-traumatic stress disorder.⁸²

The Australian study examined acamprosate and naltrexone in patients with and without depression or anxiety.^{49,66} It did not find a naltrexone or acamprosate by depression interaction when assessing predictors of abstinence (no lapse)—odds ratios (ORs) were 0.78 (95% CI, 0.60

to 1.01) and 0.95 (95% CI, 0.81 to 1.12), respectively. It also reported no anxiety by naltrexone or acamprosate interaction when assessing predictors of abstinence (OR for acamprosate by anxiety interaction, 0.92; 95% CI, 0.74 to 1.15; OR for naltrexone by anxiety, 1.06; 95% CI, 0.83 to 1.35). When assessing predictors of no relapse (at least 4 drinks for females and at least 6 drinks for males), the study found a significant naltrexone by depression interaction—OR, 0.77 (95% CI, 0.63 to 0.95) but no significant interactions for acamprosate by depression, naltrexone by anxiety, or acamprosate by anxiety.

In the U.S.-based study of patients with PTSD and alcohol dependence,¹⁷⁰ desipramine was associated with a lower percentage of heavy drinking days ($p=0.009$) and fewer drinks per drinking day ($p=0.027$) compared with paroxetine, but specific alcohol use data were not reported and the study was rated high risk of bias.

In the U.S.-based study of patients with both alcohol and cocaine dependence,¹⁹³ disulfiram was associated with a significantly lower percentage of drinking days compared with naltrexone (4.0 percent versus 26.3 percent, respectively, $p<0.01$). This study was rated high risk of bias.

Key Question 6. Genetic Polymorphisms

For this KQ, we describe the characteristics of included studies and then evidence on whether any of the medications are more or less effective for adults with certain genetic polymorphisms compared with adults without such polymorphisms. The most commonly evaluated polymorphisms were those of the mu-opioid receptor gene. For most polymorphism-medication pairs, we found just 1 eligible study, and we graded the SOE as insufficient (because evidence was imprecise, unknown consistency, and medium or high risk of bias)—information on the study characteristics and results for polymorphism-medication pairs with just 1 eligible study is provided in Appendix G. These included 1 study each for the following: nalmefene and opioid receptor gene polymorphisms; topiramate or naltrexone and *DRD*, *HTR2A*, or *SLC6A* gene polymorphisms; olanzapine and *DRD* gene polymorphisms; acamprosate or naltrexone and *GATA4* polymorphisms; sertraline and *5-HTTLPR* polymorphisms; and disulfiram and *DBH* polymorphisms.

Characteristics of Included Studies

We found no studies that assessed the clinical utility of genotype-guided dosing strategies or genotype-guided medication selection, and none randomized by genotype; all included studies assessed the association between genotype and response to medication. We found 7 eligible studies assessing variation in naltrexone response related to polymorphisms of the opioid receptor gene (Table 32). Four studies were secondary or subgroup analyses of U.S.-based randomized controlled trials; 3 were prospective cohort studies conducted in Australia,¹⁹⁶ Korea,¹⁹⁷ or Spain.¹⁹⁸ All 7 studies assessed mu-opioid receptor gene (*OPRM1*) polymorphisms; 1 also assessed polymorphisms of the genes that encode for the delta- and kappa-opioid receptors (*OPRD1* and *OPRK1*, respectively). The main polymorphism tested is in exon 1 of the *OPRM1* gene (118A>G), resulting in an asparaginase to aspartate substitution at position 40 of the amino acid sequence (Asn40Asp) of the mu-opioid receptor.

Most of the studies used naltrexone 50 mg; 1 used 100 mg.¹⁹⁹ Duration of treatment ranged from 12 to 16 weeks. Mean age was very similar across studies, from 40 to 50 years. All subjects met criteria for alcohol dependence in 6 of the studies; 1 study reported that 95 percent of subjects met criteria for alcohol dependence.²⁰⁰ Three studies enrolled all males^{197,198,201}; the others enrolled between 30 and 43 percent females.

One additional study (Oslin et al., 2003) assessing the association between opioid receptor gene polymorphisms and naltrexone response was identified that did not meet inclusion criteria.²⁰² It pooled data for a subset of subjects from 3 separate trials, 1 of which was less than 12 weeks in treatment duration. Because this study may include useful information, and has been included in previous reviews, we conducted sensitivity analyses that include this study (see below in the Overview of Results section).

Table 32. Characteristics of included studies that assessed the association between opioid receptor gene polymorphisms and naltrexone response

Author, Year Study Design	Arm Dose, mg/day (N)	Genotypes Assessed	Rx Duration, Weeks	Setting	Age, Years	% Non- white	% Fe- male	Co-inter- vention	Risk of Bias
Anton, 2008 ¹⁹⁹ SSGA	Naltrexone 100 (301) Placebo (303)	OPRM1	16	U.S.; Outpatient 11 sites	45 to 46	0	30	MM 100% CBI 49% ACA % NR	Medium
Collier, 2001 ¹⁹⁶ Prospective Cohort	Naltrexone 50 (100)	OPRM1	12	Australia; substance abuse treatment, outpatient	43	NR	43	CBI 100%	Medium
Gelernter, 2007 ²⁰¹ SSGA	Naltrexone 50 (149) Placebo (64)	OPRM1 OPRD1 OPRK1	13	U.S.; Multisite VAMCs	50	26	0	NR	High
Kim, 2009 ¹⁹⁷ Prospective Cohort	Naltrexone 50 (32)	OPRM1	12	Korea; Multiple hospitals	46 to 49	100	0	CBT 100%	High
Kranzler, 2013 ²⁰⁰ SSGA	Naltrexone 50— daily or targeted (81) Placebo (77)	OPRM1	12	U.S.; Outpatient; university health center	49	3	42	Coping skills therapy 100%	Medium
O'Malley, 2008 ⁹⁹ SSGA	Naltrexone 50 (34) Placebo (34) Naltrexone 50 + Sertraline 100 (33) ^a	OPRM1	16	U.S.; Native and non- native Alaskans, outpatient	40	70	34	MM 100%	Medium
Rubio, 2002 ¹⁹⁸ Prospective Cohort	Naltrexone 50 (45)	OPRM1	12	Spain; outpatient	NR	NR	0	NR	Unclear

^a For the SSGA of OPRM1, usable DNA was available for 92 of the 101 participants in the randomized controlled trial. Of those, 17 had one or more copies of the Asp40 allele (9 placebo, 3 NTX only, and 5 NTX + sertraline), so the authors restricted statistical analyses to the participants who were homozygous for the Asn40 allele.

Note: Age, Years is the mean age in years, unless otherwise stated.

Abbreviations: mg = milligram; MM = medical management; N = number; NR = not reported; OPRM1 = the mu-opioid receptor gene; SSGA = secondary or subgroup analysis; U.S. = United States; VAMC = Veterans Administration Medical Center.

Overview of Results

Three of the studies reported some positive associations between polymorphisms and response to naltrexone (Table 33).^{197,199,200} Specifically, they reported an association between having a G allele (i.e., at least one Asp40) and better alcohol consumption outcomes.

Our meta-analyses for return to any drinking found no significant difference between AA homozygotes and those with at least one G allele among patients treated with naltrexone, both without (RD, -0.03; 95% CI, -0.6 to 0.5) and with inclusion of the studies rated as high or unclear risk of bias (RD, 0.01; 95% CI, -0.2 to 0.2). Sensitivity analyses including the Oslin 2003 study also found no difference for return to any drinking.²⁰²

Table 33. Results of included studies that assessed the association between mu-opioid receptor gene polymorphisms and naltrexone response

Author, year	Reported a Significant Positive Association?	AA, N	AA, Return to Any Drinking	AA, Return to Heavy Drinking—Relapse	AG/GG, N	AG/GG, Return to Any Drinking	AG/GG, Return to Heavy Drinking—Relapse
Anton, 2008 ¹⁹⁹	Yes ^a	115 ^b	NR	52	31 ^b	NR	4
Coller, 2001 ¹⁹⁶	No	NR	NR	NR	NR	NR	NR
Gelernter, 2007 ²⁰¹	No	98	NR	35	33	NR	12
Kim, 2009 ¹⁹⁷	Mixed ^c	16	8	6	16	9	3
Kranzler, 2013 ²⁰⁰	Yes	59	NR	NR	22	NR	NR
O'Malley, 2008 ⁹⁹	No ^d	25	16	16	3	2	2
Rubio, 2002 ¹⁹⁸	No	29	9	9	16	4	4

^a Statistically significant difference between groups for return to heavy drinking.

^b Data are for those who received naltrexone and medical management, and do not include those who received naltrexone + medical management + CBI. The study found no gene by medication by time interactions for the latter group for percentage of days abstinent or heavy drinking days, and did not report specific numbers by genotype for the outcomes.

^c Yes for time to first relapse ($p=0.014$); no for abstinent rate ($p=0.656$) and relapse rate ($p=0.072$).

^d Study authors restricted analyses to AA homozygotes because they had only 17 of 92 genotyped participants with at least one G allele. The results for the 75 AA homozygotes were similar to the results for the total sample, indicating that treatment efficacy was not dependent on the presence of the G allele.

Note: Table only includes data for subjects who received naltrexone; it does not include data for those who received placebo or who received naltrexone plus sertraline.

Abbreviations: CBI = combined behavioral intervention; N = number; NR = not reported.

Similarly, our meta-analyses for return to heavy drinking found no statistically significant difference between AA homozygotes and those with at least one G allele among patients treated with naltrexone, both without (RD, 0.26; 95% CI, -0.01 to 0.53) and with inclusion of the studies rated as high or unclear risk of bias (RD, 0.14; 95% CI, -0.03 to 0.3). Sensitivity analyses including the Oslin 2003 study,²⁰² along with all other studies regardless of risk of bias rating, found that a lower percentage of patients with a G allele returned to heavy drinking than AA homozygotes (RD, 0.16; 95% CI, 0.02 to 0.29).

Detailed Results of Individual Studies

Subgroup analysis from the COMBINE study found no gene by medication by time interactions for patients treated with medical management plus CBI, but reported an interaction between treatment and genotype for the time trend of percentage of days abstinent and for percentage of heavy drinking days for patients who received medical management (with no CBI).¹⁹⁹ Among those who received medical management, patients with at least one Asp40 allele and treated with naltrexone had a higher proportion of good clinical outcomes (87.1 percent) than patients homozygous for Asn40 treated with naltrexone (54.8 percent) and those who received placebo who did and did not have an Asp40 allele (48.6 percent and 54 percent, respectively).

The study conducted in Australia reported a significant decrease in alcohol use over time, but no genotype by time interaction and no difference between the two genotypic groups (median grams per week: AA, 48.0 versus AG or GG, 37.5, $p=0.78$).¹⁹⁶ It also reported no difference

between genotypic groups (AA versus AG or GG) for time to first relapse (11 versus 10 days, $p=0.40$) and for mean number of drinking days (17.6 versus 21.9, $p=0.56$).

One U.S.-based study, a secondary analysis of data from a trial conducted in Veteran's Affairs Medical Centers, reported no association between the *OPRM1* genotype and naltrexone response.²⁰¹ Patients who were homozygous AA had about the same rate of relapse as those who carried a G allele (35.7 versus 36.0). The study had several limitations, including drawing the study sample from a trial that did not show a positive effect of naltrexone.

The cohort study conducted in Korea provided outcome information only for the subjects who were adherent to naltrexone for 12 weeks (32 of 63 subjects who initiated treatment).¹⁹⁷ Among those, it reported longer time to relapse for patients with a G allele than for AA homozygotes (73.3 versus 59.9 days, $p=0.014$), but no statistically significant difference between groups for abstinence rate (43.8 percent versus 50 percent, $p=0.656$) or for relapse rate (18.8 percent versus 37.5 percent, $p=0.072$).

Another U.S.-based study, a secondary analysis of data from a trial conducted in a university-based center, reported that neither genotype nor medication significantly predicted mean daily drinking levels.²⁰⁰ However, it found a positive desire by genotype by medication condition interaction, with a significant desire by genotype interaction for the placebo group ($p=0.001$) but not for the naltrexone group ($p=0.74$). In other words, when the evening desire to drink was high, G allele carriers were at greater risk than AA homozygotes to drink more.²⁰⁰

One U.S.-based study, a secondary analysis of data from a trial conducted with Alaskans, restricted its analyses to AA homozygotes.⁹⁹ The authors reported that this was because they had only 17 of 92 genotyped participants with at least one G allele. They found that the results for the 75 AA homozygotes were similar to the results for the total sample (for percentage abstinent and percentage relapsed to a heavy drinking day), indicating that treatment efficacy was not dependent on the presence of the G allele.

The cohort study conducted in Spain ($N=45$) was reported as an abstract only, with very little details about the methods.¹⁹⁸ We assessed the risk of bias of this study as unclear due to very limited reporting of information. The study did not find a significant difference in consumption outcomes (abstinence/return to any drinking, or relapse) between patients who were AA homozygotes and those with a G allele.

Discussion

Below, we summarize the main findings and strength of evidence (SOE). We then discuss the findings in relation to what is already known, applicability of the findings, implications for decisionmaking, limitations, research gaps, and conclusions. When we have graded evidence as insufficient, it indicates that evidence is either unavailable, does not permit estimation of an effect, or does not permit us to draw a conclusion with at least a low level of confidence. It does not indicate that a treatment has been proven to lack efficacy.

Key Findings and Strength of Evidence

Efficacy and Comparative Effectiveness

We found moderate SOE that both acamprosate and naltrexone are effective for improving alcohol consumption outcomes (

Table 34). Numbers needed to treat (NNT) to prevent 1 person from returning to any drinking were 10 and 25, respectively. For return to heavy drinking, evidence did not support the efficacy of acamprosate, whereas naltrexone was efficacious with an NNT of 13. Relatively limited evidence from well-controlled trials does not adequately support the efficacy of disulfiram compared with placebo for preventing return to any drinking or for other alcohol consumption outcomes. Some disulfiram trials reported fewer drinking days for subjects who returned to any drinking and who had a complete set of assessment interviews, and suggest that disulfiram may have a role in the treatment of alcohol dependence for some individuals.

We found insufficient direct evidence to conclude that treatment with acamprosate or naltrexone leads to improvement in health outcomes—i.e., accidents, injuries, quality of life (QoL), function, or mortality. Very few trials reported any health outcomes, and the included trials were not designed or powered to assess impact on health outcomes—they typically focused on alcohol consumption outcomes. It is noteworthy that the largest pharmacotherapy trial in alcohol dependence, COMBINE, did report some evidence of improvement in QoL with naltrexone plus behavioral intervention (on the SF-12v2 physical health scale), but the difference between groups did not reach a clinically meaningful threshold.¹⁷⁶ Evidence from epidemiologic literature consistently relates high average alcohol consumption and heavy per-occasion use to an increased risk of health problems, such as cancers of the oral cavity, esophagus, larynx, colon, rectum, liver, and breast; liver cirrhosis; chronic pancreatitis; coronary heart disease; stroke; depression; preterm birth complications; fetal alcohol syndrome; and injuries and violence.^{21,203-206} Such epidemiologic evidence would suggest that improving alcohol consumption outcomes is likely to result in improved health outcomes.

Our meta-analyses of 3 head-to-head randomized controlled trials (RCTs) comparing acamprosate with naltrexone,^{49,53,61} all rated as low risk of bias, found no statistically significant difference between the two medications for improvement in alcohol consumption outcomes (Table 35). The COMBINE study was one of the 3 RCTs.⁵³ It found that patients receiving medical management with naltrexone, combined behavioral intervention (CBI), or both fared better on drinking outcomes than those who received placebo, but acamprosate showed no evidence of efficacy, with or without CBI.

Table 34. Summary of findings and strength of evidence for efficacy of FDA-approved medications for alcohol dependence

Intervention	Outcome	N studies; N subjects ^a	Results Effect Size (95% CI) ^b	Strength of Evidence
Acamprosate	Return to any drinking	15; 4,747	RD: -0.10 (-0.15 to -0.05); NNT 10	Moderate
	Return to heavy drinking	6; 2,239	RD: -0.01 (-0.05 to 0.03)	Moderate
	Percentage drinking days	12; 4,385	WMD: -9.4 (-13.8 to -5.0)	Moderate
	Percentage heavy drinking days	0; 0	NA	Insufficient
	Drinks per drinking day	1; 116	WMD: 0.4 (-1.8 to 2.6)	Insufficient
	Accidents or injuries	0; ^c 0	NA	Insufficient
	Quality of life or function	1; 612	NSD	Insufficient
	Mortality	7; 2,477	7 events (ACA) vs. 5 events (placebo)	Insufficient
Disulfiram	Return to any drinking	2; 492	RD: 0.04 (-0.03 to 0.11) ^d	Low
	Return to heavy drinking	0; 0	NA	Insufficient
	Percentage drinking days	2; 290	NSD ^e	Insufficient
	Percentage heavy drinking days	0; 0	NA	Insufficient
	Drinks per drinking day	0; 0	NA	Insufficient
	Accidents or injuries	0; 0	NA	Insufficient
	Quality of life or function	0; 0	NA	Insufficient
	Mortality	0; 0	NA	Insufficient
Naltrexone	Return to any drinking	21; 4,232	RD: -0.04 (-0.07 to -0.01); NNT 25	Moderate
	Return to heavy drinking	21; 3,794	RD: -0.08 (-0.12 to -0.04); NNT 13	Moderate
	Percentage drinking days	19; 3,329	WMD: -4.6 (-6.6 to -2.5)	Moderate
	Percentage heavy drinking days	10; 1,423	WMD: -3.6 (-5.9 to -1.4)	Moderate
	Drinks per drinking day	11; 1,422	WMD: -0.5 (-1.0 to -0.07)	Low
	Accidents or injuries	0; 0	NA	Insufficient
	Quality of life or function	4; 1,513	Some conflicting results ^f	Insufficient
	Mortality	6; 1,738	1 event (NTX) vs. 2 events (placebo)	Insufficient

^a Includes only studies rated as low or medium risk of bias included in the main analyses; these numbers do not include studies rated as high or unclear risk of bias that were included in sensitivity analyses.

^b Negative effect sizes favor intervention over placebo/control.

^c One study rated as unclear risk of bias reported that one patient in the placebo group died by “accident.” No other details on the cause or nature of the accident were provided.⁶⁵

^d From meta-analysis of disulfiram 250 mg vs. control (which was disulfiram 1mg).^{78,79} Meta-analysis including studies rated as high risk of bias also found no significant difference (RD -0.00; 95% CI, -0.10 to 0.09). Similarly, our meta-analysis found no statistically significant difference between disulfiram 250 mg per day and riboflavin (i.e., no disulfiram) (RD -0.04; 95% CI, -0.11 to 0.03).

^e One study (N=128) reported similar percentages and no significant difference;⁷⁹ the other reported that disulfiram was favored among the subset of subjects (N=162 of 605 subjects) who drank and had a complete set of assessment interviews, but it did not report this outcome for the full randomized sample.⁷⁸ Overall, evidence was insufficient due to imprecision, inconsistency, and indirectness.

^f Unable to pool data. Two studies found no significant difference between naltrexone- and placebo-treated subjects.^{87,176} One study reported that patients receiving injectable naltrexone 380 mg per day had greater improvement on the mental health summary score than those receiving placebo at 24 weeks (8.2 versus 6.2, $p=0.044$).^{117,210} One study measured alcohol-related consequences (with the DrInC) and reported that more subjects who received placebo (N=34) had at least 1 alcohol-related consequence than those who received naltrexone (N=34): 76% versus 45%, $p=0.02$.⁹⁹

Abbreviations: ACA = acamprosate; CI = confidence interval; FDA = U.S. Food and Drug Administration; N = number; NA = not applicable; NNT = number needed to treat; NSD = no statistically significant difference; NTX = naltrexone; RD = risk difference; WMD = weighted mean difference.

Table 35. Summary of findings and strength of evidence for comparative effectiveness of acamprosate and naltrexone

Intervention	Outcome	N studies; N subjects ^a	Results Effect Size (95% CI) ^b	Strength of Evidence
ACA vs. NTX	Return to any drinking	3; 800	RD: 0.02 (-0.03 to 0.08)	Moderate
	Return to heavy drinking	3; 800	RD: 0.01 (-0.06 to 0.07)	Moderate
	Percentage drinking days	2; 720	WMD: -2.98 (-13.4 to 7.5)	Low

^a Includes only studies rated as low or medium risk of bias included in the main analyses; these numbers do not include studies rated as high or unclear risk of bias that were included in sensitivity analyses.

^b Negative effect sizes favor acamprosate over naltrexone.

Note: Table only includes comparisons of medications with evidence of efficacy (as determined in KQ 1) and with sufficient data for synthesis. We did not include rows in this table for outcomes that we graded as having insufficient SOE (percentage heavy drinking days, drinks per drinking day, accidents or injuries, quality of life or function, and mortality).

Abbreviations: ACA = acamprosate; CI = confidence interval; N = number; NTX = naltrexone; RD = risk difference; WMD = weighted mean difference.

For the vast majority of medications used off-label, and those under investigation, the evidence either was insufficient to determine whether they are efficacious for reducing alcohol consumption or the evidence suggested that they are not efficacious for people with alcohol dependence. We found two exceptions. First, for topiramate, we found moderate SOE supporting efficacy for reducing drinking days, heavy drinking days, and drinks per drinking day—based on the results of 2 RCTs (total N=521).^{156,160} No included RCTs reported data for return to any drinking or return to heavy drinking. Second, for nalmefene, we found moderate SOE supporting efficacy for one alcohol consumption outcome—reduction in drinks per drinking day (weighted mean difference [WMD] -1.0; 95% CI, -1. to -0.3). However, the magnitude of benefit (reduction of 1 drink per drinking day) is not likely clinically significant, and we found insufficient evidence of efficacy for nalmefene for other consumption outcomes (return to any drinking, return to heavy drinking, and heavy drinking days) and low SOE that nalmefene is not efficacious for reducing drinking days (WMD -1.1; 95% CI, -7.6 to 5.4). [Note: we are aware that new evidence on nalmefene has been published after our literature search and that nalmefene has since been approved in other countries; this new evidence will be included in our update search while the report is being reviewed and any necessary changes will be made for our final report].

Harms

Adverse events were often not collected using standardized measures, and methods for systematically capturing adverse events were often not reported. Studies were generally not designed primarily to assess adverse events; the vast majority focused on alcohol consumption outcomes. Evidence for many potential adverse events was insufficient to determine whether the risk was increased or not, often primarily because of lack of precision. For most of the specific adverse events, point estimates favored placebo (i.e., there were more adverse events with medications), but the differences were not statistically significant.

In head-to-head studies, the risk of withdrawal due to adverse events was not significantly different between acamprosate and naltrexone, whereas the risk of headache was higher for those

treated with naltrexone. Compared with placebo, patients treated with acamprosate had a higher risk of anxiety, diarrhea, and vomiting, and those treated with naltrexone had a higher risk of dizziness, nausea, and vomiting.

According to the package insert,²⁰⁷ acamprosate is contraindicated for people with severe renal impairment (creatinine clearance 30 mL per minute or less) and requires dose adjustments for moderate renal impairment (creatinine clearance between 30 and 50 mL per minute). Precautions are listed to monitor for depression and suicidal ideation. Common side effects include diarrhea and somnolence.

Naltrexone is contraindicated for patients with acute hepatitis or liver failure, and for those currently using opioids or with anticipated need for opioids.^{208,209} It can precipitate severe withdrawal for patients dependent on opioids.^{208,209} Precautions are listed in the package insert for other hepatic disease, renal impairment, and history of suicide attempts or depression. Patients should be advised to carry a wallet card to alert medical personnel because larger doses may be required and respiratory depression may be deeper and more prolonged if opioid analgesia is needed. Common side effects include nausea, vomiting, decreased appetite, headache, dizziness, fatigue, somnolence, and anxiety. Injectable naltrexone can also cause injection site reactions. Serious adverse events include precipitation of severe withdrawal if the patient is dependent on opioids, and hepatotoxicity (although it is not believed to be a hepatotoxin at the recommended doses).

Primary Care Settings

We identified no eligible trials conducted completely in primary care settings, and no eligible trials assessing U.S. Food and Drug Administration (FDA)-approved medications that were conducted in primary care settings. The only included trial conducted partly in primary care settings compared nalmefene with placebo in 15 sites (about half were primary care settings) in Finland.¹⁴² One other trial (included in Key Question [KQ] 1 but not in KQ 4) that compared naltrexone with placebo for 12 weeks in the United States described the use of a “primary care model.”⁸⁶ Although the trial did not take place in a primary care setting (it was a treatment research center), and the investigators were from a department of psychiatry, the psychosocial co-intervention was delivered by a nurse practitioner with a primary care background, and the trial may have implications for how psychosocial co-interventions could be provided in primary care settings.

Barriers to prescribing medications for alcohol dependence in primary care may include lack of familiarity with the medications, lack of confidence in their effectiveness, or inability to provide suitable psychosocial co-interventions (e.g., due to competing demands or insufficient practice resources, personnel, or training). Further, primary care providers are typically trained to refer patients with alcohol dependence for specialized treatment. O’Malley and O’Connor recently reviewed the issues surrounding the use of medications for alcohol dependence in primary care settings.²¹⁰ They concluded that “the implementation and widespread use of medications to treat alcohol problems faces a unique set of barriers in primary care. Although primary care providers are proficient at prescribing a wide variety of medications, they generally are unfamiliar with medications for treating alcohol problems other than those used to treat alcohol withdrawal.” They referenced a growing body of research to support basic screening methods, brief interventions, and especially medication therapy that has yet to have a major impact on how primary care providers care for individuals at risk for or with alcohol problems.²¹¹

Like behavioral counseling interventions for risky drinking delivered in primary care, implementing the use of medications and psychosocial co-interventions for alcohol dependence in primary care might require development of support systems and additional provider and staff training.^{204,212} Two other publications that did not meet our inclusion criteria (due to the study design or comparators) may have important implications for the use of medications for alcohol dependence in primary care settings. First, a nested sequence of three U.S.-based RCTs compared naltrexone plus “primary care management” (PCM) with naltrexone plus cognitive behavioral therapy.²¹³ PCM was provided by nurse practitioners, physician assistants, and one internist in an initial 45-minute visit, followed by 15- to 20-minute sessions in weeks 1, 2, 3, 4, 6, 8, and 10. The study found no difference in response to treatment, as measured by avoiding persistent heavy drinking, between those who received PCM and those who received cognitive behavioral therapy (84.1 percent versus 86.5 percent). Among responders enrolled in a maintenance trial, it found higher response for those who received naltrexone and PCM than for those who received placebo and PCM (80.8 percent versus 51.9 percent, $p=0.03$). Second, a pragmatic trial with 149 general practitioners in France who were “used to managing alcohol-dependent patients in their daily practice” randomized patients ($N=422$) to acamprosate plus standard care or standard care alone.²¹⁴ Standard care in France was described as typically consisting of outpatient detoxification followed by a rehabilitation program (involving some type of psychotherapy). The trial reported better outcomes for the acamprosate group for the Alcohol-Related Problems Questionnaire score, the number of subjects with no alcohol-related problems, and for all secondary outcome measures, including QoL.

Subgroups and Genetic Polymorphisms

We did not find any convincing evidence that either naltrexone or acamprosate are more or less effective (compared with each other) for men or women, older adults, young adults, racial or ethnic minorities, smokers, or those with co-occurring disorders. For genetic polymorphisms, we found no studies that assessed the clinical utility of genotype-guided dosing strategies or genotype-guided medication selection, and none that randomized by genotype. All included studies were either secondary/subgroup analyses of trials or prospective cohort studies of people treated with a medication, and all assessed the association between genotype and response to medication (i.e., clinical validity). For most polymorphism-medication pairs, we found just 1 eligible study, and we graded the SOE as insufficient.

We found 7 eligible studies assessing variation in naltrexone response related to mu-opioid receptor gene (*OPRM1*) polymorphisms. Our meta-analyses for return to any drinking and return to heavy drinking found no significant difference between AA homozygotes and those with at least one G allele, both without and with inclusion of studies rated as high or unclear risk of bias. Of note, the total number of subjects contributing data to the analyses was relatively low, and firm conclusions are limited by the imprecision of the results. Point estimates for return to heavy drinking suggest it is possible that patients with at least one G allele might be more likely to respond to naltrexone, but confidence intervals were wide; additional studies are needed to improve confidence in the estimate of the effect.

Findings in Relation to What Is Already Known

Existing guidelines and systematic reviews support our main findings.^{13,34-36,171,173} As described in the introduction, the U.S. Department of Veterans Affairs (VA), National Institute on Alcohol Abuse and Alcoholism (NIAAA), and Substance Abuse and Mental Health Services

Administration (SAMHSA) all have guidelines addressing the use of pharmacotherapy for alcohol dependence.³⁴⁻³⁶ The various guidelines recommend that naltrexone and/or acamprosate routinely be considered for patients with alcohol dependence in combination with addiction-focused counseling.

Whereas we did not find statistically significant effects on alcohol consumption outcomes for injectable naltrexone, effect sizes for injectable naltrexone were similar to those found for 50 mg per day of oral naltrexone for return to any drinking and return to heavy drinking. Fewer studies and subjects were available for injectable naltrexone; thus, analyses have less precision.

Applicability

Most studies reported that 100 percent of subjects met criteria for alcohol dependence. We did not identify any studies that evaluated medications and reported them to be efficacious for people with alcohol use disorders who did not meet criteria for alcohol dependence (i.e., people with alcohol abuse or harmful alcohol use). The mean age of subjects was generally in the 40s, with very few studies enrolling slightly younger or older populations. Thus, it is uncertain whether the medications have similar efficacy for older (e.g., those 65 and older) or younger (e.g., in the 20s) subgroups as they have for patients enrolled in the trials. We did not find evidence to confirm or refute whether treatments are more or less efficacious for many other subgroups, including gender groups, racial or ethnic minorities, smokers or nonsmokers, and those with certain coexisting conditions.

Although the majority of included trials assessing the efficacy of acamprosate were conducted in Europe (15 of 20) and a minority were conducted in the United States (3 of 20), the opposite was true for naltrexone (27 of 42 in the United States and 6 of 42 in Europe). Further, the few studies of acamprosate conducted in the United States did not find it to be efficacious. It is unclear whether the different results were due to population differences or other factors. The European trials of acamprosate typically identified patients from inpatient settings or treatment programs, whereas the U.S.-based trials of acamprosate relied on advertisements and referrals. It is possible that this resulted in populations with differing alcoholism severity and differing potential for benefit. For example, studies of subjects recruited via advertisements may enroll people who have less severe disorders, and may be less applicable to patients with more severe forms of alcohol-use disorders.

Most studies required patients to abstain for at least a few days prior to initiating medication, and the medications are generally recommended for maintenance of abstinence. Acamprosate and injectable naltrexone are only approved for use in patients who have established abstinence, though the duration of required abstinence is not set. However, some studies enrolling patients who were not yet abstinent have reported reduction in heavy drinking with naltrexone^{83,215} or acamprosate.⁶⁰

Implications for Clinical and Policy Decisionmaking

Evidence supports the efficacy of more than one pharmacological treatment for alcohol dependence, and clinical uncertainty exists about what treatment to select for individual patients. Acamprosate and naltrexone have the best evidence supporting their efficacy, but head-to-head trials have not consistently established superiority of either medication. Thus, other factors may contribute to medication choices, such as heterogeneity of alcohol dependence, coexisting symptoms such as anxiety or insomnia, frequency of administration, cost, potential type of

benefits, potential adverse events, and availability of treatments (e.g., acamprosate is currently a nonformulary medication for the VA).

For example, acamprosate is typically dosed as two 333 mg tablets given three times daily, whereas oral naltrexone is one tablet once daily, and injectable naltrexone is given once monthly. Acamprosate is contraindicated for people with severe renal impairment and requires dose adjustments for moderate renal impairment. Naltrexone is contraindicated for patients with acute hepatitis or liver failure, and for those currently using opioids or with anticipated need for opioids, and it can precipitate severe withdrawal for patients dependent on opioids. Trials of topiramate have reported a significantly increased risk of many adverse events, including paresthesias, taste perversion, anorexia, difficulty with concentration/attention, nervousness, dizziness, pruritis, psychomotor slowing, and weight loss.^{156,160}

Given that medications for alcohol dependence have been underutilized, entities providing health care for people with alcohol dependence may need to develop systems to optimize dissemination and implementation. For example, these could include campaigns to educate providers about the use of medications for alcohol dependence; systems to screen for unhealthy alcohol use and to provide appropriate interventions for people with unhealthy alcohol use; systems to ensure that people with alcohol dependence have access to knowledgeable providers who can prescribe medications for alcohol dependence; or systems to remind or incentivize providers to use effective medications for alcohol dependence when appropriate.

Limitations of the Comparative Effectiveness Review Process

The scope of this review was focused on medications. We did not evaluate the effectiveness or comparative effectiveness of other interventions for alcohol use disorders (e.g., cognitive behavioral therapy, motivational enhancement therapy, 12-step programs). We required that trials have at least 12 weeks of follow-up from the time of medication initiation, excluding trials of shorter duration. Some might consider this approach to omit potentially important information from shorter trials. However, longitudinal studies have found that treatment periods of less than 6 months' duration may yield misleading conclusions about treatment efficacy, due to fluctuations in drinking behavior that are typical of the course of alcoholism^{216,217}—suggesting that a longer duration of follow-up (6 months or more) might more accurately reflect the outcomes of greatest interest and importance.

Finally, publication bias and selective reporting are potential limitations. Although we searched for unpublished studies and unpublished outcomes, and did not find direct evidence of either of these biases, many of the included trials were published prior to the availability of trial registries (e.g., clinicaltrials.gov) that would allow for greater certainty in determining the potential for either type of bias.

Limitations of the Evidence Base

The evidence base was inadequate to draw conclusions for some of our questions or subquestions of interest. In particular, as described above, we found insufficient direct evidence on health outcomes, limited and varying reporting on harms, no trials conducted completely in primary care settings, and scant head-to-head evidence on differences for population subgroups.

We found insufficient direct evidence to determine whether medications are efficacious for improving health outcomes. Although evidence from epidemiologic literature consistently relates

high average and heavy per-occasion alcohol use to an increased risk of health problems, it is challenging to estimate the magnitude of reduction in the risk of health problems that is derived from a reduction in consumption. For example, it is unclear how much benefit (for health outcomes) is derived from 10 percent fewer patients returning to any drinking, or from 8 percent fewer patients returning to heavy drinking.

Many of the included trials had methodological limitations introducing some risk of bias. Some trials had high proportions of subjects lost to follow up. High attrition rates are not uncommon in studies of psychiatric conditions. Methods of handling missing data varied, and some trials did nothing to address missing data (i.e., only analyzing completers). However, many trials conducted true intention-to-treat analyses and used appropriate methods of handling missing data, such as imputing return to heavy drinking for subjects lost to follow-up or multiple imputation.

Reporting of previous treatments and ongoing treatments (i.e., co-interventions) was variable across the included studies. We were often unable to determine whether subjects had received any previous treatments for alcohol dependence.

Research Gaps

We identified numerous gaps in the evidence that future research could address. Many of these gaps are highlighted in the previous sections of this Discussion. Of note, these gaps relate only to the KQs addressed by this report, and they should not eliminate a wide range of potentially important research that falls outside of our scope.

Table 36 summarizes the key gaps and potential future research that could address the gaps.

Conclusions

Acamprosate and naltrexone are effective for improving alcohol consumption outcomes for patients with alcohol dependence (moderate SOE). Numbers needed to treat (NNT) to prevent one person from returning to any drinking were 10 and 25, respectively; NNT to prevent one person from returning to heavy drinking was 13 for naltrexone. Our meta-analyses of 3 head-to-head trials found no statistically significant difference between the two medications for improvement in alcohol consumption outcomes (moderate SOE). With the exception of topiramate, for which we found moderate SOE supporting efficacy for improving some consumption outcomes, current evidence does not establish the efficacy of medications used off-label and those under investigation for people with alcohol dependence. We found insufficient direct evidence to conclude whether medications for alcohol dependence are effective for improving health outcomes. No eligible trials assessing FDA-approved medications were conducted in primary care settings. Evidence was generally insufficient to determine comparative effectiveness of acamprosate and naltrexone for subgroups.

Table 36. Evidence gaps for future research, by key question

KQ	Evidence Gap	Potential Future Research
1	Evidence was insufficient to determine efficacy of some medications.	Future studies could evaluate medications that have some evidence (often from 1 or 2 small trials) suggesting possible efficacy (e.g., baclofen) or medications that have not yet been studied with some theoretical basis to support their potential efficacy.
1	We found no head-to-head studies of oral naltrexone and injectable naltrexone.	Future studies could compare the benefits of harms of oral and injectable naltrexone.
1	We found insufficient evidence evaluating medications for people with alcohol use disorders who do not meet criteria for alcohol dependence (i.e., those with alcohol abuse or harmful alcohol use).	Future studies could evaluate the efficacy of acamprosate or naltrexone in such populations.
2	We found insufficient direct evidence to conclude that treatment with acamprosate or naltrexone leads to improvement in health outcomes.	Future studies could focus on health outcomes, such as accidents, injuries, QoL, function, or mortality.
3	Relatively few studies reported information about suicide, suicidal ideation, or self-harmful behaviors.	Additional studies could be conducted to determine whether precautions about suicide, suicidal thoughts, or self-harmful behaviors are warranted.
3	Little evidence was available to determine whether naltrexone can be used for people with various liver conditions. ^a	Future studies could evaluate the use of naltrexone for people with various chronic liver conditions.
4	No eligible trials assessed the use of FDA-approved medications in primary care settings.	Future studies could evaluate the use of acamprosate and naltrexone in primary care settings.
5	Evidence on whether any medications are more or less effective than other medications for population subgroups was scant.	Future studies could compare the use of acamprosate and naltrexone for subgroups of patients (e.g., enrolling subjects who all have depression or other psychiatric conditions; comparing effectiveness for men or women or among older or younger patients)
6	Relatively few subjects contributed data to our analyses of variation in naltrexone response and <i>OPRM1</i> polymorphisms. Patients with at least one G allele may be more likely to respond to naltrexone, but confidence intervals were wide and the effect was not statistically significant.	Additional studies are likely to change our confidence in the estimate of the effect and to change the estimate.
6	No studies assessed the clinical utility of genotype-guided dosing strategies or genotype-guided medication selection, and none randomized by genotype.	If variation in naltrexone response by <i>OPRM1</i> polymorphisms becomes established, then future studies could assess the clinical utility of using genotype-guided dosing strategies. For example, studies might compare the use of genotype-guided dosing strategies (e.g., use naltrexone for patients with at least one G allele, but use acamprosate for AA homozygotes) with using naltrexone or acamprosate for all subjects.
6	Only 1 study was available for most polymorphism-medication response associations.	Future studies could explore other genotypic associations (i.e., not limiting future studies to <i>OPRM1</i> polymorphisms).

^a The FDA removed the black box warning for hepatotoxicity for injectable naltrexone, but it is unclear whether naltrexone should be used in people with various chronic liver conditions.

Abbreviations: FDA = U.S. Food and Drug Administration; *OPRM1* = mu-opioid receptor gene; QoL = quality of life.

References

1. Saitz R. Clinical practice. Unhealthy alcohol use. *N Engl J Med*. 2005 Feb 10;352(6):596-607. PMID: 15703424.
2. Whitlock EP, Green CA, Polen MR. Behavioral Counseling Interventions in Primary Care to Reduce Risky/Harmful Alcohol Use. Systematic Evidence Review No. 30. AHRQ Publication No. 04-0533B. (Prepared by Oregon Evidence-based Practice Center under Contract No. 290-97-0018, Task Order No. 2). Rockville, MD: Agency for Healthcare Research and Quality; April 2004.
3. U.S. Department of Veterans Affairs. AUDIT-C Frequently Asked Questions. Updated April 28, 2010.
<http://www.queri.research.va.gov/tools/alcohol-misuse/alcohol-faqs.cfm>. Accessed June 27, 2011.
4. Mokdad AH, Marks JS, Stroup DF, et al. Actual causes of death in the United States, 2000. *JAMA*. 2004 Mar 10;291(10):1238-45. PMID: 15010446.
5. Whitlock EP, Polen MR, Green CA, et al. Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2004 Apr 6;140(7):557-68. PMID: 15068985.
6. Bouchery EE, Harwood HJ, Sacks JJ, et al. Economic costs of excessive alcohol consumption in the U.S., 2006. *Am J Prev Med*. 2011 Nov;41(5):516-24. PMID: 22011424.
7. Harwood HJ, Fountain D, Fountain G. Economic cost of alcohol and drug abuse in the United States, 1992: a report. *Addiction*. 1999 May;94(5):631-5. PMID: 10563025.
8. Centers for Disease Control and Prevention. FastStats: Alcohol Use. Updated January 27, 2012.
<http://www.cdc.gov/nchs/fastats/alcohol.htm>. Accessed May 21, 2012.
9. Isaac M, Janca A, Sartorius N. ICD-10 symptom glossary for mental disorders. Geneva: Division of Mental Health, World Health Organization; 1994.
10. Janca A, Ustun TB, van Drimmelen J, et al. ICD-10 symptom checklist for mental disorders, version 1.1. Geneva: Division of Mental Health, World Health Organization; 1994.
11. American Psychiatric Association. Diagnostic and statistical manual of mental disorders, 4th ed. Text rev. Washington, DC: American Psychiatric Publishing, Inc.; 2000.
12. Schuckit MA. Alcohol-use disorders. *Lancet*. 2009 Feb 7;373(9662):492-501. PMID: 19168210.
13. National Collaborating Centre for Mental Health, National Institute for Health & Clinical Excellence. Alcohol-use Disorders: The NICE Guidelines on Diagnosis, Assessment and Management of Harmful Drinking and Alcohol Dependence. The British Psychological Society and The Royal College of Psychiatrists 2011.
<http://www.nice.org.uk/nicemedia/live/13337/53190/53190.pdf>.
14. Hasin DS, Stinson FS, Ogburn E, et al. Prevalence, correlates, disability, and comorbidity of DSM-IV alcohol abuse and dependence in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch Gen Psychiatry*. 2007 Jul;64(7):830-42. PMID: 17606817.
15. Mertens JR, Weisner C, Ray GT, et al. Hazardous drinkers and drug users in HMO primary care: prevalence, medical conditions, and costs. *Alcohol Clin Exp Res*. 2005 Jun;29(6):989-98. PMID: 15976525.
16. Teesson M, Baillie A, Lynskey M, et al. Substance use, dependence and treatment seeking in the United States and Australia: a cross-national comparison. *Drug Alcohol Depend*. 2006 Feb 1;81(2):149-55. PMID: 16043307.
17. Hasin DS, Grant BF. The co-occurrence of DSM-IV alcohol abuse in DSM-IV alcohol dependence: results of the National Epidemiologic Survey on Alcohol and Related Conditions on heterogeneity that differ by population subgroup. *Arch Gen Psychiatry*. 2004 Sep;61(9):891-6. PMID: 15351767.
18. Mann K, Schafer DR, Langle G, et al. The long-term course of alcoholism, 5, 10 and 16 years after treatment. *Addiction*. 2005 Jun;100(6):797-805. PMID: 15918810.

19. Norstrom T. Per capita alcohol consumption and all-cause mortality in Canada, 1950-98. *Addiction*. 2004 Oct;99(10):1274-8. PMID: 15369565.
20. Rivara FP, Garrison MM, Ebel B, et al. Mortality attributable to harmful drinking in the United States, 2000. *J Stud Alcohol*. 2004 Jul;65(4):530-6. PMID: 15376828.
21. Corrao G, Bagnardi V, Zambon A, et al. A meta-analysis of alcohol consumption and the risk of 15 diseases. *Prev Med*. 2004 May;38(5):613-9. PMID: 15066364.
22. Cherpitel CJ, Ye Y. Alcohol-attributable fraction for injury in the U.S. general population: data from the 2005 National Alcohol Survey. *J Stud Alcohol Drugs*. 2008 Jul;69(4):535-8. PMID: 18612569.
23. . Alcohol-attributable deaths and years of potential life lost--United States, 2001. *MMWR Morb Mortal Wkly Rep*. 2004 Sep 24;53(37):866-70. PMID: 15385917.
24. Moos RH, Moos BS. Rates and predictors of relapse after natural and treated remission from alcohol use disorders. *Addiction*. 2006 Feb;101(2):212-22. PMID: 16445550.
25. Rumpf HJ, Bischof G, Hapke U, et al. Stability of remission from alcohol dependence without formal help. *Alcohol Alcohol*. 2006 May-Jun;41(3):311-4. PMID: 16490790.
26. Bodin MC, Romelsjo A. Predictors of abstinence and nonproblem drinking after 12-step treatment in Sweden. *J Stud Alcohol*. 2006 Jan;67(1):139-46. PMID: 16536138.
27. Cox WM, Rosenberg H, Hodgins CH, et al. United Kingdom and United States healthcare providers' recommendations of abstinence versus controlled drinking. *Alcohol Alcohol*. 2004 Mar-Apr;39(2):130-4. PMID: 14998830.
28. Dawson DA, Grant BF, Stinson FS, et al. Recovery from DSM-IV alcohol dependence: United States, 2001-2002. *Addiction*. 2005 Mar;100(3):281-92. PMID: 15733237.
29. Maisto SA, Clifford PR, Stout RL, et al. Drinking in the year after treatment as a predictor of three-year drinking outcomes. *J Stud Alcohol*. 2006 Nov;67(6):823-32. PMID: 17060998.
30. Schuckit MA. *Drug and Alcohol Abuse: A Clinical Guide to Diagnosis and Treatment*. 6th ed., New York: Springer; 2005.
31. Miller WR, Walters ST, Bennett ME. How effective is alcoholism treatment in the United States? *J Stud Alcohol*. 2001 Mar;62(2):211-20. PMID: 11327187.
32. O'Brien CP, McLellan AT. Myths about the treatment of addiction. *Lancet*. 1996 Jan 27;347(8996):237-40. PMID: 8551886.
33. Kranzler HR, Van Kirk J. Efficacy of naltrexone and acamprosate for alcoholism treatment: a meta-analysis. *Alcohol Clin Exp Res*. 2001 Sep;25(9):1335-41. PMID: 11584154.
34. U.S. Department of Veterans Affairs, U.S. Department of Defense. VA/DoD Clinical Practice Guideline for Management of Substance Use Disorders (SUD). 2009. http://www.healthquality.va.gov/sud/sud_full_601f.pdf.
35. Pettinati HM, Weiss RD, Miller WR, et al. COMBINE Monograph Series, Volume 2. Medical Management Treatment Manual: A Clinical Research Guide for Medically Trained Clinicians Providing Pharmacotherapy as Part of the Treatment for Alcohol Dependence. DHHS Publication No. (NIH) 04-5289. Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; 2004.
36. Center for Substance Abuse and Treatment. Incorporating alcohol pharmacotherapies into medical practice. (Treatment improvement protocol (TIP); no. 49). Substance Abuse and Mental Health Services Administration (SAMHSA) Rockville, MD: 2009.
37. West SL, Garbutt JC, Carey TS, et al. Pharmacotherapy for Alcohol Dependence. Evidence report number 3. (Contract 290-97-0011 to Research Triangle Institute, University of North Carolina, Chapel Hill) AHCPR publication no. 99-E004. Rockville, MD: Agency for Health Care Policy and Research; January 1999. <http://www.ncbi.nlm.nih.gov/books/NBK32930/>.
38. Garbutt JC, West SL, Carey TS, et al. Pharmacological treatment of alcohol dependence: a review of the evidence. *JAMA*. 1999 Apr 14;281(14):1318-25. PMID: 10208148.
39. *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*. Agency for Healthcare Research and Quality Rockville (MD). <http://www.ncbi.nlm.nih.gov/books/NBK47095/>.

40. Viswanathan M, Ansari MT, Berkman ND, et al. Assessing the Risk of Bias of Individual Studies in Systematic Reviews of Health Care Interventions. . Agency for Healthcare Research and Quality Methods Guide for Comparative Effectiveness Reviews AHRQ Publication No. 12-EHC047-EF. Rockville, MD: Agency for Healthcare Research and Quality; March 2012. www.effectivehealthcare.ahrq.gov/.
41. West SL, Gartlehner G, Mansfield AJ, et al. Comparative Effectiveness Review Methods: Clinical Heterogeneity. Methods Research Report. Prepared by RTI International -- University of North Carolina Evidence-based Practice Center under Contract No. 290-2007-10056-I AHRQ Publication No. 10-EHC070-EF. Rockville, MD: Agency for Healthcare Research and Quality; September 2010.
42. Sutton AJ, Abrams KR, Jones DR, et al. Methods for Meta-Analysis in Medical Research (Wiley Series in Probability and Statistics - Applied Probability and Statistics Section). London: Wiley; 2000.
43. Centers for Disease Control and Prevention. What is a standard drink in the United States? ; 2013. <http://www.cdc.gov/alcohol/faqs.htm#standDrink>. Accessed August 20, 2013.
44. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002 Jun 15;21(11):1539-58. PMID: 12111919.
45. Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ*. 2003 Sep 6;327(7414):557-60. PMID: 12958120.
46. Higgins JP, Green ST, eds. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]: The Cochrane Collaboration; 2011.
47. Owens DK, Lohr KN, Atkins D, et al. AHRQ series paper 5: grading the strength of a body of evidence when comparing medical interventions--Agency for Healthcare Research and Quality and the Effective Health-Care Program. *J Clin Epidemiol*. 2010 May;63(5):513-23. PMID: 19595577.
48. Atkins DC, S.; Gartlehner G.; et al. Chapter 6: Assessing the applicability of studies when comparing medical interventions. Agency for Healthcare Research and Quality AHRQ Publication No. 11-EHC019-EF. Rockville, MD: 2011.
49. Morley KC, Teesson M, Reid SC, et al. Naltrexone versus acamprosate in the treatment of alcohol dependence: A multi-centre, randomized, double-blind, placebo-controlled trial. *Addiction*. 2006 Oct;101(10):1451-62. PMID: 16968347.
50. Ralevski E, O'Brien E, Jane JS, et al. Effects of acamprosate on cognition in a treatment study of patients with schizophrenia spectrum disorders and comorbid alcohol dependence. *J Nerv Ment Dis*. 2011 Jul;199(7):499-505. PMID: 21716064.
51. Baltieri DA, De Andrade AG. Acamprosate in alcohol dependence: a randomized controlled efficacy study in a standard clinical setting. *J Stud Alcohol*. 2004 Jan;65(1):136-9. PMID: 15000513.
52. Pelc I, Verbanck P, Le Bon O, et al. Efficacy and safety of acamprosate in the treatment of detoxified alcohol-dependent patients. A 90-day placebo-controlled dose-finding study. *Br J Psychiatry*. 1997 Jul;171:73-7. PMID: 9328500.
53. Anton RF, O'Malley SS, Ciraulo DA, et al. Combined pharmacotherapies and behavioral interventions for alcohol dependence: the COMBINE study: a randomized controlled trial. *JAMA*. 2006 May 3;295(17):2003-17. PMID: 16670409.
54. Mason BJ, Goodman AM, Chabac S, et al. Effect of oral acamprosate on abstinence in patients with alcohol dependence in a double-blind, placebo-controlled trial: the role of patient motivation. *J Psychiatr Res*. 2006 Aug;40(5):383-93. PMID: 16546214.
55. Fucito LM, Park A, Gulliver SB, et al. Cigarette smoking predicts differential benefit from naltrexone for alcohol dependence. *Biol Psychiatry*. 2012;72(10):832-8.
56. Donovan DM, Anton RF, Miller WR, et al. Combined pharmacotherapies and behavioral interventions for alcohol dependence (The COMBINE Study): examination of posttreatment drinking outcomes. *J Stud Alcohol Drugs*. 2008 Jan;69(1):5-13. PMID: 18080059.
57. Besson J, Aebly F, Kasas A, et al. Combined efficacy of acamprosate and disulfiram in the treatment of alcoholism: a controlled study. *Alcohol Clin Exp Res*. 1998 May;22(3):573-9. PMID: 9622434.
58. Chick J, Howlett H, Morgan MY, et al. United Kingdom Multicentre Acamprosate Study (UKMAS): a 6-month prospective study of

- acamprosate versus placebo in preventing relapse after withdrawal from alcohol. *Alcohol Alcohol*. 2000 Mar-Apr;35(2):176-87. PMID: 10787394.
59. Geerlings PJ, Ansoms C, Van Den Brink W. Acamprosate and prevention of relapse in alcoholics. Results of a randomized, placebo-controlled, double-blind study in out-patient alcoholics in the Netherlands, Belgium and Luxembourg. *Eur Addict Res*. 1997;3(3):129-37.
 60. Gual A, Leheret P. Acamprosate during and after acute alcohol withdrawal: a double-blind placebo-controlled study in Spain. *Alcohol Alcohol*. 2001;36(5):413-8. PMID: CN-00367117.
 61. Kiefer F, Jahn H, Tarnaske T, et al. Comparing and combining naltrexone and acamprosate in relapse prevention of alcoholism: a double-blind, placebo-controlled study. *Arch Gen Psychiatry*. 2003 Jan;60(1):92-9. PMID: 12511176.
 62. Kiefer F, Andersohn F, Otte C, et al. Long-term effects of pharmacotherapy on relapse prevention in alcohol dependence. *Acta Neuropsychiatrica*. 2004;18:233-8.
 63. Kiefer F, Helwig H, Tarnaske T, et al. Pharmacological relapse prevention of alcoholism: clinical predictors of outcome. *Eur Addict Res*. 2005;11(2):83-91. PMID: 15785069.
 64. Lhuintre JP, Daoust M, Moore ND, et al. Ability of calcium bis acetyl homotaurine, a GABA agonist, to prevent relapse in weaned alcoholics. *Lancet*. 1985 May 4;1(8436):1014-6. PMID: 2859465.
 65. Lhuintre JP, Moore N, Tran G, et al. Acamprosate appears to decrease alcohol intake in weaned alcoholics. *Alcohol Alcohol*. 1990;25(6):613-22. PMID: 2085344.
 66. Morley KC, Teesson M, Sannibale C, et al. Clinical predictors of outcome from an Australian pharmacological relapse prevention trial. *Alcohol Alcohol*. 2010 Nov-Dec;45(6):520-6. PMID: 20952764.
 67. Paille FM, Guelfi JD, Perkins AC, et al. Double-blind randomized multicentre trial of acamprosate in maintaining abstinence from alcohol. *Alcohol Alcohol*. 1995 Mar;30(2):239-47. PMID: 7662044.
 68. Pelc I, Le Bon O, Leheret P, et al. Acamprosate in the Treatment of Alcohol Dependence: A 6-Month Postdetoxification Study. In: Soyka M, ed. *Acamprosate in Relapse Prevention of Alcoholism*. Springer Berlin Heidelberg; 1996:133-42.
 69. Pelc I, Le Bon O, Verbanck P, et al. Calciumacetylhomotaurinate for maintaining abstinence in weaned alcoholic patients: a placebo-controlled double-blind multi-centre study. In: Naranjo CA, Sellers EM, eds. *Novel Pharmacological Interventions for Alcoholism*. New York: Springer-Verlag; 1992.
 70. Poldrugo F. Acamprosate treatment in a long-term community-based alcohol rehabilitation programme. *Addiction*. 1997 Nov;92(11):1537-46. PMID: 9519495.
 71. Ralevski E, O'Brien E, Jane JS, et al. Treatment with acamprosate in patients with schizophrenia spectrum disorders and comorbid alcohol dependence. *J Dual Diagn*. 2011;7(1-2):64-73.
 72. Sass H, Soyka M, Mann K, et al. Relapse prevention by acamprosate. Results from a placebo-controlled study on alcohol dependence. *Arch Gen Psychiatry*. 1996 Aug;53(8):673-80. PMID: 8694680.
 73. Tempesta E, Janiri L, Bignamini A, et al. Acamprosate and relapse prevention in the treatment of alcohol dependence: a placebo-controlled study. *Alcohol Alcohol*. 2000 Mar-Apr;35(2):202-9. PMID: 10787398.
 74. Whitworth AB, Fischer F, Lesch OM, et al. Comparison of acamprosate and placebo in long-term treatment of alcohol dependence. *Lancet*. 1996 May 25;347(9013):1438-42. PMID: 8676626.
 75. Wolwer W, Frommann N, Janner M, et al. The effects of combined acamprosate and integrative behaviour therapy in the outpatient treatment of alcohol dependence: a randomized controlled trial. *Drug Alcohol Depend*. 2011 Nov 1;118(2-3):417-22. PMID: 21621929.
 76. Petrakis IL, Poling J, Levinson C, et al. Naltrexone and disulfiram in patients with alcohol dependence and comorbid psychiatric disorders. *Biol Psychiatry*. 2005 May 15;57(10):1128-37. PMID: 15866552.
 77. Ling W, Weiss DG, Charuvastra VC, et al. Use of disulfiram for alcoholics in methadone maintenance programs. A Veterans Administration Cooperative Study. *Arch Gen Psychiatry*. 1983 Aug;40(8):851-4. PMID: 6347118.
 78. Fuller RK, Branchey L, Brightwell DR, et al. Disulfiram treatment of alcoholism. A Veterans

- Administration cooperative study. *JAMA*. 1986 Sep 19;256(11):1449-55. PMID: 3528541.
79. Fuller RK, Roth HP. Disulfiram for the treatment of alcoholism. An evaluation in 128 men. *Ann Intern Med*. 1979 Jun;90(6):901-4. PMID: 389121.
 80. Ralevski E, Ball S, Nich C, et al. The impact of personality disorders on alcohol-use outcomes in a pharmacotherapy trial for alcohol dependence and comorbid Axis I disorders. *Am J Addict*. 2007 Nov-Dec;16(6):443-9. PMID: 18058408.
 81. Petrakis I, Ralevski E, Nich C, et al. Naltrexone and disulfiram in patients with alcohol dependence and current depression. *J Clin Psychopharmacol*. 2007 Apr;27(2):160-5. PMID: 17414239.
 82. Petrakis IL, Poling J, Levinson C, et al. Naltrexone and disulfiram in patients with alcohol dependence and comorbid post-traumatic stress disorder. *Biol Psychiatry*. 2006 Oct 1;60(7):777-83. PMID: 17008146.
 83. Garbutt JC, Kranzler HR, O'Malley SS, et al. Efficacy and tolerability of long-acting injectable naltrexone for alcohol dependence: a randomized controlled trial. *JAMA*. 2005 Apr 6;293(13):1617-25. PMID: 15811981.
 84. Johnson BA, Ait-Daoud N, Aubin HJ, et al. A pilot evaluation of the safety and tolerability of repeat dose administration of long-acting injectable naltrexone (Vivitrex) in patients with alcohol dependence. *Alcohol Clin Exp Res*. 2004 Sep;28(9):1356-61. PMID: 15365306.
 85. Kranzler HR, Wesson DR, Billot L. Naltrexone depot for treatment of alcohol dependence: a multicenter, randomized, placebo-controlled clinical trial. *Alcohol Clin Exp Res*. 2004 Jul;28(7):1051-9. PMID: 15252291.
 86. Monterosso JR, Flannery BA, Pettinati HM, et al. Predicting treatment response to naltrexone: the influence of craving and family history. *Am J Addict*. 2001 Summer;10(3):258-68. PMID: 11579624.
 87. Morgenstern J, Kuerbis AN, Chen AC, et al. A randomized clinical trial of naltrexone and behavioral therapy for problem drinking men who have sex with men. *J Consult Clin Psychol*. 2012;80(5):863-75. PMID: 22612306.
 88. Oslin DW, Lynch KG, Pettinati HM, et al. A placebo-controlled randomized clinical trial of naltrexone in the context of different levels of psychosocial intervention. *Alcohol Clin Exp Res*. 2008 Jul;32(7):1299-308. PMID: 18540910.
 89. Pettinati HM, Oslin DW, Kampman KM, et al. A double-blind, placebo-controlled trial combining sertraline and naltrexone for treating co-occurring depression and alcohol dependence. *Am J Psychiatry*. 2010 Jun;167(6):668-75. PMID: 20231324.
 90. Schmitz JM, Lindsay JA, Green CE, et al. High-dose naltrexone therapy for cocaine-alcohol dependence. *Am J Addict*. 2009 Sep-Oct;18(5):356-62. PMID: 19874153.
 91. Pettinati HM, Kampman KM, Lynch KG, et al. Gender differences with high-dose naltrexone in patients with co-occurring cocaine and alcohol dependence. *J Subst Abuse Treat*. 2008 Jun;34(4):378-90. PMID: 17664051.
 92. Oslin D, Liberto JG, O'Brien J, et al. Naltrexone as an adjunctive treatment for older patients with alcohol dependence. *Am J Geriatr Psychiatry*. 1997 Fall;5(4):324-32. PMID: 9363289.
 93. Balldin J, Berglund M, Borg S, et al. A 6-month controlled naltrexone study: combined effect with cognitive behavioral therapy in outpatient treatment of alcohol dependence. *Alcohol Clin Exp Res*. 2003 Jul;27(7):1142-9. PMID: 12878920.
 94. Heinala P, Alho H, Kiianmaa K, et al. Targeted use of naltrexone without prior detoxification in the treatment of alcohol dependence: a factorial double-blind, placebo-controlled trial. *J Clin Psychopharmacol*. 2001 Jun;21(3):287-92. PMID: 11386491.
 95. Krystal JH, Cramer JA, Krol WF, et al. Naltrexone in the treatment of alcohol dependence. *N Engl J Med*. 2001 Dec 13;345(24):1734-9. PMID: 11742047.
 96. Longabaugh R, Wirtz PW, Gulliver SB, et al. Extended naltrexone and broad spectrum treatment or motivational enhancement therapy. *Psychopharmacology (Berl)*. 2009 Oct;206(3):367-76. PMID: 19639303.
 97. Huang MC, Chen CH, Yu JM, et al. A double-blind, placebo-controlled study of naltrexone in the treatment of alcohol dependence in Taiwan. *Addict Biol*. 2005 Sep;10(3):289-92. PMID: 16109592.
 98. Lee A, Tan S, Lim D, et al. Naltrexone in the treatment of male alcoholics—An effectiveness study In Singapore. *Drug and Alcohol Review*. 2001;20(2):193-9.

99. O'Malley SS, Robin RW, Levenson AL, et al. Naltrexone alone and with sertraline for the treatment of alcohol dependence in Alaska natives and non-natives residing in rural settings: a randomized controlled trial. *Alcohol Clin Exp Res*. 2008 Jul;32(7):1271-83. PMID: 18482155.
100. Schmitz JM, Stotts AL, Sayre SL, et al. Treatment of cocaine-alcohol dependence with naltrexone and relapse prevention therapy. *Am J Addict*. 2004 Jul-Sep;13(4):333-41. PMID: 15370932.
101. Volpicelli JR, Clay KL, Watson NT, et al. Naltrexone in the treatment of alcoholism: predicting response to naltrexone. *J Clin Psychiatry*. 1995;56 Suppl 7:39-44. PMID: 7673104.
102. Volpicelli JR, Rhines KC, Rhines JS, et al. Naltrexone and alcohol dependence. Role of subject compliance. *Arch Gen Psychiatry*. 1997 Aug;54(8):737-42. PMID: 9283509.
103. O'Malley SS, Sinha R, Grilo CM, et al. Naltrexone and cognitive behavioral coping skills therapy for the treatment of alcohol drinking and eating disorder features in alcohol-dependent women: a randomized controlled trial. *Alcohol Clin Exp Res*. 2007 Apr;31(4):625-34. PMID: 17374042.
104. Anton RF, Myrick H, Wright TM, et al. Gabapentin combined with naltrexone for the treatment of alcohol dependence. *Am J Psychiatry*. 2011 Jul;168(7):709-17. PMID: 21454917.
105. Baltieri DA, Daro FR, Ribeiro PL, et al. Comparing topiramate with naltrexone in the treatment of alcohol dependence. *Addiction*. 2008 Dec;103(12):2035-44. PMID: 18855810.
106. Brown ES, Carmody TJ, Schmitz JM, et al. A randomized, double-blind, placebo-controlled pilot study of naltrexone in outpatients with bipolar disorder and alcohol dependence. *Alcohol Clin Exp Res*. 2009 Nov;33(11):1863-9. PMID: 19673746.
107. Petrakis IL, O'Malley S, Rounsaville B, et al. Naltrexone augmentation of neuroleptic treatment in alcohol abusing patients with schizophrenia. *Psychopharmacology (Berl)*. 2004 Mar;172(3):291-7. PMID: 14634716.
108. Killeen TK, Brady KT, Gold PB, et al. Effectiveness of naltrexone in a community treatment program. *Alcohol Clin Exp Res*. 2004 Nov;28(11):1710-7. PMID: 15547458.
109. Ahmadi J, Ahmadi N. A double blind, placebo-controlled study of naltrexone in the treatment of alcohol dependence. *German Journal of Psychiatry*. 2002;5(4):85-9.
110. Ahmadi J, Babaebeigi M, Maany I, et al. Naltrexone for alcohol-dependent patients. *Ir J Med Sci*. 2004 Jan-Mar;173(1):34-7. PMID: 15732235.
111. Anton RF, Moak DH, Waid LR, et al. Naltrexone and cognitive behavioral therapy for the treatment of outpatient alcoholics: results of a placebo-controlled trial. *Am J Psychiatry*. 1999 Nov;156(11):1758-64. PMID: 10553740.
112. Anton RF, Moak DH, Latham PK, et al. Posttreatment results of combining naltrexone with cognitive-behavior therapy for the treatment of alcoholism. *J Clin Psychopharmacol*. 2001 Feb;21(1):72-7. PMID: 11199951.
113. Anton RF, Moak DH, Latham P, et al. Naltrexone combined with either cognitive behavioral or motivational enhancement therapy for alcohol dependence. *J Clin Psychopharmacol*. 2005 Aug;25(4):349-57. PMID: 16012278.
114. Baltieri DA, Daro FR, Ribeiro PL, et al. Effects of topiramate or naltrexone on tobacco use among male alcohol-dependent outpatients. *Drug Alcohol Depend*. 2009 Nov 1;105(1-2):33-41. PMID: 19595518.
115. Chick J, Anton R, Chęcinski K, et al. A multicentre, randomized, double-blind, placebo-controlled trial of naltrexone in the treatment of alcohol dependence or abuse. *Alcohol Alcohol*. 2000 Nov-Dec;35(6):587-93. PMID: 11093966.
116. Fogaca MN, Santos-Galduroz RF, Eserian JK, et al. The effects of polyunsaturated fatty acids in alcohol dependence treatment--a double-blind, placebo-controlled pilot study. *BMC Clin Pharmacol*. 2011;11:10. PMID: 21787433.
117. Pettinati HM, Gastfriend DR, Dong Q, et al. Effect of extended-release naltrexone (XR-NTX) on quality of life in alcohol-dependent patients. *Alcohol Clin Exp Res*. 2009 Feb;33(2):350-6. PMID: 19053979.
118. Gastpar M, Bonnet U, Boning J, et al. Lack of efficacy of naltrexone in the prevention of alcohol relapse: results from a German multicenter study. *J Clin Psychopharmacol*. 2002 Dec;22(6):592-8. PMID: 12454559.
119. Guardia J, Caso C, Arias F, et al. A double-blind, placebo-controlled study of naltrexone in the

- treatment of alcohol-dependence disorder: results from a multicenter clinical trial. *Alcohol Clin Exp Res*. 2002 Sep;26(9):1381-7. PMID: 12351933.
120. Kranzler HR, Tennen H, Armeli S, et al. Targeted naltrexone for problem drinkers. *J Clin Psychopharmacol*. 2009 Aug;29(4):350-7. PMID: 19593174.
 121. Latt NC, Jurd S, Houseman J, et al. Naltrexone in alcohol dependence: a randomised controlled trial of effectiveness in a standard clinical setting. *Med J Aust*. 2002 Jun 3;176(11):530-4. PMID: 12064984.
 122. Monti PM, Rohsenow DJ, Swift RM, et al. Naltrexone and cue exposure with coping and communication skills training for alcoholics: treatment process and 1-year outcomes. *Alcohol Clin Exp Res*. 2001 Nov;25(11):1634-47. PMID: 11707638.
 123. Rohsenow DJ, Miranda R, Jr., McGeary JE, et al. Family history and antisocial traits moderate naltrexone's effects on heavy drinking in alcoholics. *Exp Clin Psychopharmacol*. 2007 Jun;15(3):272-81. PMID: 17563214.
 124. Rohsenow DJ, Colby SM, Monti PM, et al. Predictors of compliance with naltrexone among alcoholics. *Alcohol*. 2000;24(10):1542-9.
 125. Morris PL, Hopwood M, Whelan G, et al. Naltrexone for alcohol dependence: a randomized controlled trial. *Addiction*. 2001 Nov;96(11):1565-73. PMID: 11784454.
 126. O'Malley SS, Jaffe AJ, Chang G, et al. Naltrexone and coping skills therapy for alcohol dependence. A controlled study. *Arch Gen Psychiatry*. 1992 Nov;49(11):881-7. PMID: 1444726.
 127. O'Malley SS, Jaffe AJ, Chang G, et al. Six-month follow-up of naltrexone and psychotherapy for alcohol dependence. *Arch Gen Psychiatry*. 1996 Mar;53(3):217-24. PMID: 8611058.
 128. Ralevski E, Balachandra K, Gueorguieva R, et al. Effects of naltrexone on cognition in a treatment study of patients with schizophrenia and comorbid alcohol dependence. *J Dual Diagn*. 2006;2(4):53-69.
 129. Addolorato G, Leggio L, Ferrulli A, et al. Effectiveness and safety of baclofen for maintenance of alcohol abstinence in alcohol-dependent patients with liver cirrhosis: randomised, double-blind controlled study. *Lancet*. 2007 Dec 8;370(9603):1915-22. PMID: 18068515.
 130. Garbutt JC, Kampov-Polevoy AB, Gallop R, et al. Efficacy and safety of baclofen for alcohol dependence: a randomized, double-blind, placebo-controlled trial. *Alcohol Clin Exp Res*. 2010 Nov;34(11):1849-57. PMID: 20662805.
 131. Kranzler HR, Burleson JA, Del Boca FK, et al. Buspirone treatment of anxious alcoholics. A placebo-controlled trial. *Arch Gen Psychiatry*. 1994 Sep;51(9):720-31. PMID: 8080349.
 132. Malcolm R, Anton RF, Randall CL, et al. A placebo-controlled trial of buspirone in anxious inpatient alcoholics. *Alcohol Clin Exp Res*. 1992 Dec;16(6):1007-13. PMID: 1335217.
 133. Fawcett J, Kravitz HM, McGuire M, et al. Pharmacological treatments for alcoholism: revisiting lithium and considering buspirone. *Alcohol Clin Exp Res*. 2000 May;24(5):666-74. PMID: 10832908.
 134. Malec E, Malec T, Gagne MA, et al. Buspirone in the treatment of alcohol dependence: a placebo-controlled trial. *Alcohol Clin Exp Res*. 1996 Apr;20(2):307-12. PMID: 8730222.
 135. George DT, Rawlings R, Eckardt MJ, et al. Buspirone treatment of alcoholism: age of onset, and cerebrospinal fluid 5-hydroxyindolacetic acid and homovanillic acid concentrations, but not medication treatment, predict return to drinking. *Alcohol Clin Exp Res*. 1999 Feb;23(2):272-8. PMID: 10069556.
 136. Tiihonen J, Ryyanen OP, Kauhanen J, et al. Citalopram in the treatment of alcoholism: a double-blind placebo-controlled study. *Pharmacopsychiatry*. 1996 Jan;29(1):27-9. PMID: 8852531.
 137. Naranjo CA, Bremner KE, Lanctot KL. Effects of citalopram and a brief psycho-social intervention on alcohol intake, dependence and problems. *Addiction*. 1995 Jan;90(1):87-99. PMID: 7888983.
 138. Cornelius JR, Salloum IM, Ehler JG, et al. Fluoxetine in depressed alcoholics. A double-blind, placebo-controlled trial. *Arch Gen Psychiatry*. 1997 Aug;54(8):700-5. PMID: 9283504.
 139. Kabel DI, Petty F. A placebo-controlled, double-blind study of fluoxetine in severe alcohol dependence: adjunctive pharmacotherapy during and after inpatient treatment. *Alcohol Clin Exp Res*. 1996 Jun;20(4):780-4. PMID: 8800399.

140. Kranzler HR, Burleson JA, Korner P, et al. Placebo-controlled trial of fluoxetine as an adjunct to relapse prevention in alcoholics. *Am J Psychiatry*. 1995 Mar;152(3):391-7. PMID: 7864265.
141. Cornelius JR, Salloum IM, Ehler JG, et al. Double-blind fluoxetine in depressed alcoholic smokers. *Psychopharmacol Bull*. 1997;33(1):165-70. PMID: 9133770.
142. Karhuvaara S, Simojoki K, Virta A, et al. Targeted nalmefene with simple medical management in the treatment of heavy drinkers: a randomized double-blind placebo-controlled multicenter study. *Alcohol Clin Exp Res*. 2007 Jul;31(7):1179-87. PMID: 17451401.
143. Mason BJ, Salvato FR, Williams LD, et al. A double-blind, placebo-controlled study of oral nalmefene for alcohol dependence. *Arch Gen Psychiatry*. 1999;56(8):719-24.
144. Mason BJ, Ritvo EC, Morgan RO, et al. A double-blind, placebo-controlled pilot study to evaluate the efficacy and safety of oral nalmefene HCl for alcohol dependence. *Alcohol*. 1994;18(5):1162-7.
145. Anton RF, Pettinati H, Zweben A, et al. A multi-site dose ranging study of nalmefene in the treatment of alcohol dependence. *J Clin Psychopharmacol*. 2004 Aug;24(4):421-8. PMID: 15232334.
146. Brown ES, Garza M, Carmody TJ. A randomized, double-blind, placebo-controlled add-on trial of quetiapine in outpatients with bipolar disorder and alcohol use disorders. *J Clin Psychiatry*. 2008 May;69(5):701-5. PMID: 18312058.
147. Stedman M, Pettinati HM, Brown ES, et al. A double-blind, placebo-controlled study with quetiapine as adjunct therapy with lithium or divalproex in bipolar I patients with coexisting alcohol dependence. *Alcohol Clin Exp Res*. 2010 Oct;34(10):1822-31. PMID: 20626727.
148. Kampman KM, Pettinati HM, Lynch KG, et al. A double-blind, placebo-controlled pilot trial of quetiapine for the treatment of Type A and Type B alcoholism. *J Clin Psychopharmacol*. 2007 Aug;27(4):344-51. PMID: 17632217.
149. Brady KT, Sonne S, Anton RF, et al. Sertraline in the treatment of co-occurring alcohol dependence and posttraumatic stress disorder. *Alcohol Clin Exp Res*. 2005 Mar;29(3):395-401. PMID: 15770115.
150. Coskunol H, Gökden O, Ercan ES, et al. Long-term efficacy of sertraline in the prevention of alcoholic relapses in alcohol-dependent patients: a single-center, double-blind, randomized, placebo-controlled, parallel-group study. *Current Therapeutic Research*. 2002;63(11):759-71. PMID: 2003140542.
151. Gual A, Balcells M, Torres M, et al. Sertraline for the prevention of relapse in detoxicated alcohol dependent patients with a comorbid depressive disorder: a randomized controlled trial. *Alcohol Alcohol*. 2003 Nov-Dec;38(6):619-25. PMID: 14633652.
152. Kranzler HR, Armeli S, Tennen H, et al. A double-blind, randomized trial of sertraline for alcohol dependence: moderation by age of onset [corrected] and 5-hydroxytryptamine transporter-linked promoter region genotype. *J Clin Psychopharmacol*. 2011 Feb;31(1):22-30. PMID: 21192139.
153. Kranzler HR, Armeli S, Tennen H. Post-treatment outcomes in a double-blind, randomized trial of sertraline for alcohol dependence. *Alcohol Clin Exp Res*. 2012 Apr;36(4):739-44. PMID: 21981418.
154. Moak DH, Anton RF, Latham PK, et al. Sertraline and cognitive behavioral therapy for depressed alcoholics: results of a placebo-controlled trial. *J Clin Psychopharmacol*. 2003 Dec;23(6):553-62. PMID: 14624185.
155. Pettinati HM, Volpicelli JR, Luck G, et al. Double-blind clinical trial of sertraline treatment for alcohol dependence. *J Clin Psychopharmacol*. 2001 Apr;21(2):143-53. PMID: 11270910.
156. Johnson BA, Ait-Daoud N, Bowden CL, et al. Oral topiramate for treatment of alcohol dependence: a randomised controlled trial. *Lancet*. 2003 May 17;361(9370):1677-85. PMID: 12767733.
157. Rubio G, Martinez-Gras I, Manzanares J. Modulation of impulsivity by topiramate: implications for the treatment of alcohol dependence. *J Clin Psychopharmacol*. 2009 Dec;29(6):584-9. PMID: 19910725.
158. Ma JZ, Ait-Daoud N, Johnson BA. Topiramate reduces the harm of excessive drinking: implications for public health and primary care. *Addiction*. 2006 Nov;101(11):1561-8. PMID: 17034435.

159. Johnson BA, Ait-Daoud N, Akhtar FZ, et al. Oral topiramate reduces the consequences of drinking and improves the quality of life of alcohol-dependent individuals: a randomized controlled trial. *Arch Gen Psychiatry*. 2004 Sep;61(9):905-12. PMID: 15351769.
160. Johnson BA, Rosenthal N, Capece JA, et al. Topiramate for treating alcohol dependence: a randomized controlled trial. *JAMA*. 2007 Oct 10;298(14):1641-51. PMID: 17925516.
161. Johnson BA, Rosenthal N, Capece JA, et al. Improvement of physical health and quality of life of alcohol-dependent individuals with topiramate treatment: US multisite randomized controlled trial. *Arch Intern Med*. 2008 Jun 9;168(11):1188-99. PMID: 18541827.
162. Salloum IM, Cornelius JR, Daley DC, et al. Efficacy of valproate maintenance in patients with bipolar disorder and alcoholism: a double-blind placebo-controlled study. *Arch Gen Psychiatry*. 2005 Jan;62(1):37-45. PMID: 15630071.
163. Brady KT, Myrick H, Henderson S, et al. The use of divalproex in alcohol relapse prevention: a pilot study. *Drug Alcohol Depend*. 2002 Aug 1;67(3):323-30. PMID: 12127203.
164. de Sousa A, de Sousa A. An open randomized study comparing disulfiram and acamprosate in the treatment of alcohol dependence. *Alcohol Alcohol*. 2005 Nov-Dec;40(6):545-8. PMID: 16043433.
165. Laaksonen E, Koski-Jannes A, Salaspuro M, et al. A randomized, multicentre, open-label, comparative trial of disulfiram, naltrexone and acamprosate in the treatment of alcohol dependence. *Alcohol Alcohol*. 2008 Jan-Feb;43(1):53-61. PMID: 17965444.
166. De Sousa A, De Sousa A. A one-year pragmatic trial of naltrexone vs disulfiram in the treatment of alcohol dependence. *Alcohol Alcohol*. 2004 Nov-Dec;39(6):528-31. PMID: 15525790.
167. Nava F, Premi S, Manzato E, et al. Comparing treatments of alcoholism on craving and biochemical measures of alcohol consumption. *J Psychoactive Drugs*. 2006 Sep;38(3):211-7. PMID: 17165363.
168. De Sousa A, De Sousa A. An open randomized trial comparing disulfiram and naltrexone in adolescents with alcohol dependence. *J Subst Use*. 2008;13(6):382-8.
169. Martinotti G, Di Nicola M, Di Giannantonio M, et al. Aripiprazole in the treatment of patients with alcohol dependence: a double-blind, comparison trial vs. naltrexone. *J Psychopharmacol*. 2009 Mar;23(2):123-9. PMID: 18515460.
170. Petrakis IL, Ralevski E, Desai N, et al. Noradrenergic vs serotonergic antidepressant with or without naltrexone for veterans with PTSD and comorbid alcohol dependence. *Neuropsychopharmacology*. 2012 Mar;37(4):996-1004. PMID: 22089316.
171. Rosner S, Hackl-Herrwerth A, Leucht S, et al. Acamprosate for alcohol dependence. *Cochrane Database Syst Rev*. 2010(9):CD004332. PMID: 20824837.
172. Jorgensen CH, Pedersen B, Tonnesen H. The efficacy of disulfiram for the treatment of alcohol use disorder. *Alcohol Clin Exp Res*. 2011 Oct;35(10):1749-58. PMID: 21615426.
173. Rösner S, Hackl-Herrwerth A, Leucht S, et al. Opioid antagonists for alcohol dependence. *Cochrane Database Syst Rev*: John Wiley & Sons, Ltd; 2010.
174. Mason BJ, Leher P. Acamprosate for alcohol dependence: a sex-specific meta-analysis based on individual patient data. *Alcohol Clin Exp Res*. 2012 Mar;36(3):497-508. PMID: 21895717.
175. National Collaborating Centre for Mental Health. Alcohol-Use Disorders. Diagnosis, Assessment and Management of Harmful Drinking and Alcohol Dependence. National Clinical Practice Guideline 115. National Institute for Health & Clinical Excellence. London: Psychiatrists TBPSaTRCo; 2011. <http://guidance.nice.org.uk/CG115/Guidance/pdf/English>.
176. LoCastro JS, Youngblood M, Cisler RA, et al. Alcohol treatment effects on secondary nondrinking outcomes and quality of life: the COMBINE study. *J Stud Alcohol Drugs*. 2009 Mar;70(2):186-96. PMID: 19261230.
177. Miller WR, Tonigan JS, Longabaugh R. The Drinker Inventory of Consequences (DrInC): An instrument for assessing adverse consequences of alcohol abuse. Test manual. (Volume 4, Project MATCH Monograph Series). Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; 1995.

178. Ware J, Kosinski M, Dewey J. How to Score Version 2 of the SF-36 Health Survey. Lincoln, RI.: QualityMetric Incorporated; 2000.
179. Chick J, Aschauer H, Hornik K. Efficacy of fluvoxamine in preventing relapse in alcohol dependence: a one-year, double-blind, placebo-controlled multicentre study with analysis by typology. *Drug Alcohol Depend.* 2004 Apr 9;74(1):61-70. PMID: 15072808.
180. Endicott J, Nee J, Harrison W, et al. Quality of Life Enjoyment and Satisfaction Questionnaire: a new measure. *Psychopharmacol Bull.* 1993;29(2):321-6. PMID: 8290681.
181. Sheehan KH, Sheehan DV. Assessing treatment effects in clinical trials with the discan metric of the Sheehan Disability Scale. *Int Clin Psychopharmacol.* 2008 Mar;23(2):70-83. PMID: 18301121.
182. . EuroQol--a new facility for the measurement of health-related quality of life. The EuroQol Group. *Health Policy.* 1990 Dec;16(3):199-208. PMID: 10109801.
183. Koskenvuo M. The Finnish Twin Registry Baseline Characteristics. Helsinki: Kansanterveyslaitoksen laitokset: Helsingin yliopisto; 1979.
184. Scott J, Huskisson EC. Graphic representation of pain. *Pain.* 1976 Jun;2(2):175-84. PMID: 1026900.
185. Florez G, Garcia-Portilla P, Alvarez S, et al. Using topiramate or naltrexone for the treatment of alcohol-dependent patients. *Alcohol Clin Exp Res.* 2008 Jul;32(7):1251-9. PMID: 18482157.
186. Florez G, Saiz PA, Garcia-Portilla P, et al. Topiramate for the treatment of alcohol dependence: comparison with naltrexone. *Eur Addict Res.* 2011;17(1):29-36. PMID: 20975274.
187. De Sousa AA, De Sousa J, Kapoor H. An open randomized trial comparing disulfiram and topiramate in the treatment of alcohol dependence. *J Subst Abuse Treat.* 2008 Jun;34(4):460-3. PMID: 17629442.
188. Rubio G, Jimenez-Arriero MA, Ponce G, et al. Naltrexone versus acamprosate: one year follow-up of alcohol dependence treatment. *Alcohol Alcohol.* 2001 Sep-Oct;36(5):419-25. PMID: 11524308.
189. Tollefson GD, Lancaster SP, Montague-Clouse J. The association of buspirone and its metabolite 1-pyrimidinylpiperazine in the remission of comorbid anxiety with depressive features and alcohol dependency. *Psychopharmacol Bull.* 1991;27(2):163-70.
190. Tollefson GD, Montague-Clouse J, Tollefson SL. Treatment of comorbid generalized anxiety in a recently detoxified alcoholic population with a selective serotonergic drug (buspirone). *J Clin Psychopharmacol.* 1992 Feb;12(1):19-26. PMID: 1552035.
191. Narayana PL, Gupta AK, Sharma PK. Use of anti-craving agents in soldiers with alcohol dependence syndrome. *Medical Journal Armed Forces India.* 2008;64(4):320-4.
192. Volpicelli JR, Pettinati HM, McLellan AT, et al. Combining medication and psychosocial Treatments for Addiction: The BRENDA Approach. New York, NY: The Guilford Press; 2001.
193. Carroll K, Ziedonis D, O'Malley SS, et al. Pharmacologic interventions for alcohol- and cocaine-abusing individuals: A pilot study of disulfiram vs. naltrexone. *Am J Addict.* 1993;2(1):77-9.
194. Greenfield SF, Pettinati HM, O'Malley S, et al. Gender differences in alcohol treatment: an analysis of outcome from the COMBINE study. *Alcohol Clin Exp Res.* 2010 Oct;34(10):1803-12. PMID: 20645934.
195. Derogatis LR, Lipman RS, Covi L. SCL-90: an outpatient psychiatric rating scale--preliminary report. *Psychopharmacol Bull.* 1973 Jan;9(1):13-28. PMID: 4682398.
196. Collier JK, Cahill S, Edmonds C, et al. OPRM1 A118G genotype fails to predict the effectiveness of naltrexone treatment for alcohol dependence. *Pharmacogenet Genomics.* 2011 Dec;21(12):902-5. PMID: 21946895.
197. Kim SG, Kim CM, Choi SW, et al. A mu opioid receptor gene polymorphism (A118G) and naltrexone treatment response in adherent Korean alcohol-dependent patients. *Psychopharmacology (Berl).* 2009;201(4):611-8.
198. Rubio G, Ponce G, Jiménez-Arriero MA, et al. Polymorphism for m-opioid receptor (+118) as a prognostic variable of naltrexone in alcohol dependence treatment: Preliminary results. *Eur Neuropsychopharmacol.* 2002;12:397.
199. Anton RF, Oroszi G, O'Malley S, et al. An evaluation of mu-opioid receptor (OPRM1) as a predictor of naltrexone response in the treatment

- of alcohol dependence: results from the Combined Pharmacotherapies and Behavioral Interventions for Alcohol Dependence (COMBINE) study. *Arch Gen Psychiatry*. 2008 Feb;65(2):135-44. PMID: 18250251.
200. Kranzler HR, Armeli S, Covault J, et al. Variation in OPRM1 moderates the effect of desire to drink on subsequent drinking and its attenuation by naltrexone treatment. *Addiction Biology*. 2013;18(1):193-201.
 201. Gelernter J, Gueorguieva R, Kranzler HR, et al. Opioid receptor gene (OPRM1, OPRK1, and OPRD1) variants and response to naltrexone treatment for alcohol dependence: results from the VA Cooperative Study. *Alcohol Clin Exp Res*. 2007 Apr;31(4):555-63. PMID: 17374034.
 202. Oslin DW, Berrettini W, Kranzler HR, et al. A functional polymorphism of the mu-opioid receptor gene is associated with naltrexone response in alcohol-dependent patients. *Neuropsychopharmacology*. 2003 Aug;28(8):1546-52. PMID: 12813472.
 203. Rehm J, Baliunas D, Borges GL, et al. The relation between different dimensions of alcohol consumption and burden of disease: an overview. *Addiction*. 2010 May;105(5):817-43. PMID: 20331573.
 204. Jonas DE, Garbutt JC, Amick HR, et al. Behavioral counseling after screening for alcohol misuse in primary care: a systematic review and meta-analysis for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2012 Nov 6;157(9):645-54. PMID: 23007881.
 205. Bondy SJ, Rehm J, Ashley MJ, et al. Low-risk drinking guidelines: the scientific evidence. *Can J Public Health*. 1999 Jul-Aug;90(4):264-70. PMID: 10489725.
 206. Shalala DE. 10th Special Report to the U.S. Congress on Alcohol and Health: Highlights From Current Research: From the Secretary of Health and Human Services. Washington, DC: U.S. Department of Health and Human Services; 2000.
<http://pubs.niaaa.nih.gov/publications/10report/intro>. Accessed 8 June 2012.
 207. CAMPRAL® (acamprosate calcium) [package insert]. . St. Louis, MO: Forest Pharmaceuticals, Inc.; 2005.
 208. REVIA® (naltrexone hydrochloride) [package insert]. . Duramed Pharmaceuticals, Inc. Pomona, NY; 2009.
 209. VIVITROL® (naltrexone for extended-release injectable suspension) [package insert]. . Waltham, MA: Alkermes, Inc.; 2010.
 210. O'Malley SS, O'Connor PG. Medications for unhealthy alcohol use: across the spectrum. *Alcohol Res Health*. 2011;33(4):300-12. PMID: 23580015.
 211. D'Amico EJ, Paddock SM, Burnam A, et al. Identification of and guidance for problem drinking by general medical providers: results from a national survey. *Med Care*. 2005 Mar;43(3):229-36. PMID: 15725979.
 212. Jonas DE, Garbutt JC, Brown JM, et al. Screening, Behavioral Counseling, and Referral in Primary Care to Reduce Alcohol Misuse. Comparative Effectiveness Review No. 64. (Prepared by the RTI International-University of North Carolina Evidence-based Practice Center under Contract No. 290-2007-10056-I.) AHRQ Publication No. 12-EHC055-EF. Rockville, MD: Agency for Healthcare Research and Quality; July 2012.
www.effectivehealthcare.ahrq.gov/reports/final.cfm.
 213. O'Malley SS, Rounsaville BJ, Farren C, et al. Initial and maintenance naltrexone treatment for alcohol dependence using primary care vs specialty care: a nested sequence of 3 randomized trials. *Arch Intern Med*. 2003 Jul 28;163(14):1695-704. PMID: 12885685.
 214. Kiritze-Topor P, Huas D, Rosenzweig C, et al. A pragmatic trial of acamprosate in the treatment of alcohol dependence in primary care. *Alcohol Alcohol*. 2004 Nov-Dec;39(6):520-7. PMID: 15304381.
 215. Kranzler HR, Armeli S, Tennen H, et al. Targeted naltrexone for early problem drinkers. *J Clin Psychopharmacol*. 2003 Jun;23(3):294-304. PMID: 12826991.
 216. Polich JM, Armor DJ, Braiker HB. Stability and change in drinking patterns. *The Course of Alcoholism: Four Years After Treatment*. New York, NY: John Wiley & Sons; 1981:159-200.
 217. Kissin B, Charnoff SM, Rosenblatt SM. Drug and placebo responses in chronic alcoholics. *Psychiatr Res Rep Am Psychiatr Assoc*. 1968 Mar;24:44-60. PMID: 4889329.

Appendix A. Search Strategy

PubMed

Search Query	Items found
#1 Search "Alcohol-Related Disorders" [MeSH]	92008
#2 Search "Alcoholism" [MeSH]	64059
#3 Search "Alcohol Drinking" [MeSH]	46842
#4 Search alcohol depend*	8221
#5 Search "alcohol misuse"	1331
#6 Search alcohol addiction*	724
#7 Search "alcohol abuse"	12291
#8 Search problem drink*	2220
#9 Search alcohol problem*	2955
#10 Search "alcohol consumption"	25255
#11 Search harmful alcohol*	223
#12 Search harmful drink*	244
#13 Search ((drinking[tiab] OR drinker[tiab] OR drinkers[tiab]) AND alcohol[tiab])	24901
#14 Search (#1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13)	146095
#15 Search "Alcohol Deterrents"[MeSH]	1053
#16 Search ("Naltrexone"[Mesh] OR naltrexone)	7566
#17 Search ReVia	7567
#18 Search Vivitrol	12
#19 Search ("acamprosate" [Supplementary Concept] OR acamprosate)	603
#20 Search Campral	605
#21 Search ("Disulfiram"[Mesh] OR Disulfiram)	3661
#22 Search Antabuse	3730
#23 Search ("Amitriptyline"[Mesh] OR Amitriptyline)	7828
#24 Search ("aripiprazole" [Supplementary Concept] OR aripiprazole)	2079
#25 Search ("atomoxetine" [Supplementary Concept] OR atomoxetine)	964
#26 Search ("Baclofen"[Mesh] OR Baclofen)	6326
#27 Search ("Buspirone"[Mesh] OR Buspirone)	2546
#28 Search ("Citalopram"[Mesh] OR citalopram)	4661
#29 Search ("Desipramine"[Mesh] OR Desipramine)	7383
#30 Search escitalopram	4916
#31 Search ("Fluoxetine"[Mesh] OR Fluoxetine)	10276
#32 Search ("Fluvoxamine"[Mesh] OR Fluvoxamine)	2470
#33 Search ("gabapentin" [Supplementary Concept] OR gabapentin)	4127
#34 Search ("Imipramine"[Mesh] OR Imipramine)	12137
#35 Search ("nalmefene" [Supplementary Concept] OR nalmefene)	245
#36 Search ("olanzapine" [Supplementary Concept] OR olanzapine)	6265
#37 Search ("Ondansetron"[Mesh] OR Ondansetron)	3576
#38 Search ("Paroxetine"[Mesh] OR paroxetine)	4965
#39 Search ("Prazosin"[Mesh] OR Prazosin)	12613
#40 Search ("quetiapine" [Supplementary Concept] OR quetiapine)	3081
#41 Search ("Sertraline"[Mesh] OR Sertraline)	3462
#42 Search ("topiramate"[Supplementary Concept] OR topiramate)	3130
#43 Search ("Valproic Acid"[Mesh] OR Valproate)	13895
#44 Search ("varenicline"[Supplementary Concept] OR varenicline)	820
#45 Search ("Viloxazine"[Mesh] OR Viloxazine)	318
#46 Search (#15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28 or #29 or #30 or #31 or #32 or #33 or #34 or #35 or #36 or #37 or #38 or #39 or #40 or #41 or #42 or #43 or #44 or #45)	105016
#47 Search (#14 and #46)	3828
#48 Search (#14 and #46) Filters: Humans	2856
#49 Search (#14 and #46) Filters: Humans; English	2406

Search Query	Items found
#50 Search (#14 and #46) Filters: Humans; English; Adult: 19+ years	1296
#51 Search (#14 and #46) Filters: Publication date from 1970/01/01; Humans; English; Adult: 19+ years	1270
#52 Search (comment[pt] OR editorial[pt] OR letter[pt] OR news[pt])	1353911
#53 Search (#51 not #52)	1165

PSYCINFO 2-11-13

Search ID#	Search Terms	Search Options	Actions
S49	S48	Limiters - Publication Year from: 1970-2013; English; Language: English; Age Groups: Adulthood (18 yrs & older); Population Group: Human Search modes - Boolean/Phrase	View Results (957) View Details Edit
S48	S14 AND S46	Narrow by SubjectAge: - adulthood (18 yrs & older) Search modes - Boolean/Phrase	View Results (997) View Details Edit
S47	S14 AND S46	Search modes - Boolean/Phrase	View Results (1,938) View Details Edit
S46	S15 OR S16 OR S17 OR S18 OR S19 OR S20 OR S21 OR S22 OR S23 OR S24 OR S25 OR S26 OR S27 OR S28 OR S29 OR S30 OR S31 OR S32 OR S33 OR S34 OR S35 OR S36 OR S37 OR S38 OR S39 OR S40 OR S41 OR S42 OR S43 OR S44 OR S45	Search modes - Boolean/Phrase	View Results (33,948) View Details Edit
S45	Viloxazine	Search modes - Boolean/Phrase	View Results (108) View Details Edit
S44	varenicline	Search modes - Boolean/Phrase	View Results (314) View Details Edit
S43	"Valproic Acid" OR Valproate	Search modes - Boolean/Phrase	View Results (3,170) View Details Edit
S42	topiramate	Search modes - Boolean/Phrase	View Results (1,147) View Details Edit
S41	Sertraline	Search modes - Boolean/Phrase	View Results (2,064) View Details Edit
S40	quetiapine	Search modes - Boolean/Phrase	View Results (2,387) View Details Edit
S39	Prazosin	Search modes - Boolean/Phrase	View Results (486) View Details Edit
S38	Paroxetine	Search modes - Boolean/Phrase	View Results (2,731) View Details Edit
S37	Ondansetron	Search modes - Boolean/Phrase	View Results (367) View Details Edit
S36	olanzapine	Search modes - Boolean/Phrase	View Results (4,620) View Details Edit
S35	nalmeferine	Search modes - Boolean/Phrase	View Results (68) View Details Edit
S34	Imipramine	Search modes - Boolean/Phrase	View Results (3,866) View Details Edit

Search ID#	Search Terms	Search Options	Actions
S33	gabapentin	Search modes - Boolean/Phrase	View Results (954) View Details Edit
S32	Fluvoxamine	Search modes - Boolean/Phrase	View Results (1,413) View Details Edit
S31	Fluoxetine	Search modes - Boolean/Phrase	View Results (5,313) View Details Edit
S30	escitalopram	Search modes - Boolean/Phrase	View Results (759) View Details Edit
S29	Desipramine	Search modes - Boolean/Phrase	View Results (1,996) View Details Edit
S28	Citalopram	Search modes - Boolean/Phrase	View Results (1,977) View Details Edit
S27	Buspirone	Search modes - Boolean/Phrase	View Results (1,303) View Details Edit
S26	Baclofen	Search modes - Boolean/Phrase	View Results (936) View Details Edit
S25	atomoxetine	Search modes - Boolean/Phrase	View Results (495) View Details Edit
S24	aripiprazole	Search modes - Boolean/Phrase	View Results (1,410) View Details Edit
S23	Amitriptyline	Search modes - Boolean/Phrase	View Results (2,183) View Details Edit
S22	Antabuse	Search modes - Boolean/Phrase	View Results (154) View Details Edit
S21	Disulfiram	Search modes - Boolean/Phrase	View Results (573) View Details Edit
S20	Campral	Search modes - Boolean/Phrase	View Results (13) View Details Edit
S19	acamprosate	Search modes - Boolean/Phrase	View Results (329) View Details Edit
S18	Vivitrol	Search modes - Boolean/Phrase	View Results (12) View Details Edit
S17	ReVia	Search modes - Boolean/Phrase	View Results (18) View Details Edit
S16	naltrexone	Search modes - Boolean/Phrase	View Results (2,556) View Details Edit
S15	"Alcohol Deterrents"	Search modes - Boolean/Phrase	View Results (1) View Details Edit

Search ID#	Search Terms	Search Options	Actions
S14	S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13	Search modes - Boolean/Phrase	View Results (69,149) View Details Edit
S13	TI ((drinking OR drinker OR drinkers) AND alcohol) OR AB ((drinking OR drinker OR drinkers) AND alcohol)	Search modes - Boolean/Phrase	View Results (19,034) View Details Edit
S12	harmful drink*	Search modes - Boolean/Phrase	View Results (355) View Details Edit
S11	harmful alcohol*	Search modes - Boolean/Phrase	View Results (479) View Details Edit
S10	"alcohol consumption"	Search modes - Boolean/Phrase	View Results (11,811) View Details Edit
S9	alcohol problem*	Search modes - Boolean/Phrase	View Results (10,184) View Details Edit
S8	problem drink*	Search modes - Boolean/Phrase	View Results (4,978) View Details Edit
S7	"alcohol abuse"	Search modes - Boolean/Phrase	View Results (20,553) View Details Edit
S6	alcohol addiction*	Search modes - Boolean/Phrase	View Results (2,985) View Details Edit
S5	"alcohol misuse"	Search modes - Boolean/Phrase	View Results (1,159) View Details Edit
S4	alcohol depend*	Search modes - Boolean/Phrase	View Results (14,899) View Details Edit
S3	(DE "Alcohol Drinking Attitudes" OR DE "Alcohol Drinking Patterns") OR (DE "Alcohol Intoxication")	Search modes - Boolean/Phrase	View Results (19,320) View Details Edit
S2	DE "Alcoholism"	Search modes - Boolean/Phrase	View Results (23,596) View Details Edit
S1	"Alcohol-Related Disorders"	Search modes - Boolean/Phrase	View Results (203) View Details Edit

CINAHL 2-11-13

#	Query	Limiters/Expanders	Last Run Via	Results
S50 S48 NOT S49		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	6
S49 PT comment OR editorial OR letter OR news		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	65,069
S48 S47		Limiters - Published Date from: 19700101-20131231; English Language; Exclude MEDLINE records; Human; Language: English; Age Groups: All Adult Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	6
S47 S14 AND S46		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	676
S46 S15 OR S16 OR S17 OR S18 OR S19 OR S20 OR S21 OR S22 OR S23 OR S24 OR S25 OR S26 OR S27 OR S28 OR S29 OR S30 OR S31 OR S32 OR S33 OR S34 OR S35 OR S36 OR S37 OR S38 OR S39 OR S40 OR S41 OR S42 OR S43 OR S44 OR S45		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	9,728
S45 Viloxazine		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	4
S44 varenicline		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	233
S43 "Valproic Acid" OR Valproate		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	1,159
S42 topiramate		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	676
S41 Sertraline		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	602
S40 quetiapine		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	555
S39 Prazosin		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	0
S38 Paroxetine		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	695
S37 Ondansetron		Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	491

#	Query	Limiters/Expanders	Last Run Via	Results
S36	olanzapine	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	1,060
S35	nalmeferene	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	20
S34	Imipramine	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	211
S33	gabapentin	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	909
S32	Fluvoxamine	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	139
S31	Fluoxetine	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	1,095
S30	escitalopram	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	173
S29	Desipramine	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	111
S28	Citalopram	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	566
S27	Buspirone	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	147
S26	Baclofen	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	608
S25	atomoxetine	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	245
S24	aripiprazole	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	391
S23	Amitriptyline	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	512
S22	Antabuse	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	16
S21	Disulfiram	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	153
S20	Campral	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	5
S19	acamprosate	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	115
S18	Vivitrol	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	35

#	Query	Limiters/Expanders	Last Run Via	Results
S17	ReVia	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	11
S16	Naltrexone	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	800
S15	(MH "Alcohol Deterrents")	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	150
S14	S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	25,084
S13	T1 ((drinking OR drinker OR drinkers) AND alcohol) OR AB ((drinking OR drinker OR drinkers) AND alcohol)	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	5,120
S12	harmful drink*	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	145
S11	harmful alcohol*	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	199
S10	"alcohol consumption"	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	4,342
S9	alcohol problem*	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	2,512
S8	problem drink*	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	744
S7	"alcohol abuse"	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	6,285
S6	alcohol addiction*	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	291
S5	"alcohol misuse"	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	490
S4	alcohol depend*	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	2,448
S3	(MH "Alcohol Drinking")	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	10,939
S2	(MH "Alcoholism")	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	7,614
S1	(MH "Alcohol-Related Disorders")	Search modes - Boolean/Phrase	Interface - EBSCOhost Search Screen - Advanced Search Database - CINAHL with Full Text	297

EMBASE minus PubMed– 2-6-13

ID	Search	Hits
#50	#48 NOT #49 AND [1970-2013]/py	1,730
#49	editorial:it OR letter:it OR note:it AND [1970-2013]/py	1,757,884
#48	#47 AND ([adult]/lim OR [aged]/lim) AND [humans]/lim AND [english]/lim AND [embase]/lim AND [1970-2013]/py	1,929
#47	#14 AND #46	10,860
#46	#15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34 OR #35 OR #36 OR #37 OR #38 OR #39 OR #40 OR #41 OR #42 OR #43 OR #44 OR #45	249,162
#45	'viloxazine'/exp OR viloxazine	1,500
#44	'varenicline'/exp OR varenicline	2,131
#43	'valproic acid'/exp OR 'valproic acid' OR 'valproate'/exp OR valproate	47,334
#42	'topiramate'/exp OR topiramate	13,558
#41	'sertraline'/exp OR sertraline	18,228
#40	'quetiapine'/exp OR quetiapine	14,084
#39	'prazosin'/exp OR prazosin	23,503
#38	'paroxetine'/exp OR paroxetine	21,767
#37	'ondansetron'/exp OR ondansetron	12,066
#36	'olanzapine'/exp OR olanzapine	22,547
#35	'nalmefene'/exp OR nalmefene	851
#34	'imipramine'/exp OR imipramine	33,844
#33	'gabapentin'/exp OR gabapentin	18,926
#32	'fluvoxamine'/exp OR fluvoxamine	11,524
#31	'fluoxetine'/exp OR fluoxetine	35,680
#30	'escitalopram'/exp OR escitalopram	5,709
#29	'desipramine'/exp OR desipramine	20,984
#28	'citalopram'/exp OR citalopram	16,194
#27	'buspirone'/exp OR buspirone	7,963
#26	'baclofen'/exp OR baclofen	14,053
#25	'atomoxetine'/exp OR atomoxetine	2,961
#24	'aripiprazole'/exp OR aripiprazole	7,609
#23	'amitriptyline'/exp OR amitriptyline	32,939
#22	'antabuse'/exp OR antabuse	7,397
#21	'disulfiram'/exp OR disulfiram	7,707
#20	'campral'/exp OR campral	1,631
#19	'acamprosate'/exp OR acamprosate	1,672
#18	'vivitrol'/exp OR vivitrol	10,702
#17	'revia'/exp OR revia	10,713
#16	'naltrexone'/exp OR naltrexone	11,537
#15	'alcohol deterrents'	14
#14	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13	218,034
#13	drinking:ti OR drinker:ti OR drinkers:ti AND alcohol:ti OR (drinking:ab OR drinker:ab OR drinkers:ab AND alcohol:ab)	31,024
#12	harmful AND drink*	1,545
#11	harmful AND alcohol*	2,907
#10	'alcohol consumption'/exp	64,311
#9	'alcohol'/exp AND problem*	50,149
#8	problem AND drink*	41,548
#7	'alcohol abuse'/exp	20,002
#6	'alcohol'/exp AND addiction*	36,999
#5	'alcohol misuse'	1,697
#4	'alcohol'/exp AND depend*	30,931
#3	'drinking behavior'/exp	32,528
#2	'alcoholism'/exp	95,795
#1	'alcohol-related disorders'/exp	95,795

Cochrane Library

Cochrane Reviews – 209

Other reviews – 12

Trials – 587

Technology Assessments (1) Economic Evaluations (9) Cochrane Groups (3)

ID	Search	Hits
#1	[mh "Alcohol-Related Disorders"]	3159
#2	[mh Alcoholism]	2169
#3	[mh "Alcohol Drinking"]	2082
#4	alcohol depend*	3909
#5	"alcohol misuse"	170
#6	alcohol addiction*	1223
#7	"alcohol abuse"	1013
#8	problem drink*	1172
#9	alcohol problem*	2315
#10	"alcohol consumption"	2443
#11	harmful alcohol*	426
#12	harmful drink*	195
#13	(drinking:ti or drinking:ab or drinker:ti or drinker:ab or drinkers:ti or drinkers:ab) and (alcohol:ti or alcohol:ab)	2424
#14	#1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13	9573
#15	[mh "Alcohol Deterrents"]	146
#16	[mh Naltrexone] or naltrexone	1073
#17	ReVia	8
#18	Vivitrol	3
#19	acamprosate	182
#20	Campral	8
#21	[mh Disulfiram] or Disulfiram	224
#22	Antabuse	24
#23	[mh Amitriptyline] or Amitriptyline	2241
#24	aripiprazole	431
#25	atomoxetine	217
#26	[mh Baclofen] or Baclofen	346
#27	[mh Buspirone] or Buspirone	488
#28	[mh Citalopram] or Citalopram	1232
#29	[mh Desipramine] or Desipramine	797
#30	escitalopram	507
#31	[mh Fluoxetine] or Fluoxetine	2595
#32	[mh Fluvoxamine] or Fluvoxamine	846
#33	gabapentin	770
#34	[mh Imipramine] or Imipramine	2152
#35	nalmefene	79
#36	olanzapine	1881
#37	[mh Ondansetron] or Ondansetron	1664
#38	[mh Paroxetine] or Paroxetine	1915
#39	[mh Prazosin] or Prazosin	1010
#40	quetiapine	773
#41	[mh Sertraline] or Sertraline	1450
#42	topiramate	604
#43	[mh "Valproic Acid"] or Valproate	1172
#44	varenicline	215
#45	[mh Viloxazine] or Viloxazine	142
#46	#15 or #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27 or #28 or #29 or #30 or #31 or #32 or #33 or #34 or #35 or #36 or #37 or #38 or #39 or #40 or #41 or #42 or #43 or #44 or #45	19565

ID	Search	Hits
#47	#14 and #46	1238
#48	comment:pt or editorial:pt or letter:pt or news:pt	6337
#49	#47 not #48 from 1970 to 2013	1223
#50	adult or adults or [mh adult]	292412
#51	#49 and #50	821

Appendix B. Excluded Studies

Not published in English

1. Barrias JA, Chabac S, Ferreira L, et al. Acamprosate: multicenter Portuguese efficacy and tolerance evaluation study. *Psiquiatria Clinica*. 1997;18:149-60.
2. Castro LA, Laranjeira R. [A double blind, randomized and placebo-controlled clinical trial with naltrexone and brief intervention in outpatient treatment of alcohol dependence]. *Jornal Brasileiro de Psiquiatria*. 2009;58(2):79-85. PMID: CN-00754994.
3. Geerlings P, Ansoms C, Van DBW. Acamprosate and relapse prevention in outpatient alcoholics; results from a randomized, placebo-controlled double-blind study in the Benelux. *Tijdschrift Voor Alcohol, Drugs En Andere Psychotrope Stoffen*. 1995;21(3):129-41. PMID: CN-00170357.
4. Ivanets NN, Anokhina IP, Kogan BM, et al. [The efficacy and mechanisms of action of lerivon in alcoholism]. *Zhurnal neurologii i psikiatrii imeni S.S. Korsakova / Ministerstvo zdravookhraneniia i meditsinsko? promyshlennosti Rossi?sko? Federatsii, Vserossi?skoe obshchestvo nevrologov [i] Vserossi?skoe obshchestvo psikiatrov*. 1996;96(5):52-8. PMID: CN-00134237.
5. Kiefer F, Jahn H, Holzbach R, et al. The NALCAM-study: Efficacy, tolerability, outcome. *Sucht*. 2003;49(6):342-51. PMID: CN-00475102.
6. Krupitski EM, Burakov AM, Ivanov VB, et al. [The use of baclofen for treating affective disorders in alcoholism]. *Zhurnal neurologii i psikiatrii imeni S.S. Korsakova / Ministerstvo zdravookhraneniia i meditsinsko? promyshlennosti Rossi?sko? Federatsii, Vserossi?skoe obshchestvo nevrologov [i] Vserossi?skoe obshchestvo psikiatrov*. 1994;94(1):57-61. PMID: CN-00102296.
7. Ladewig D, Knecht T, Leher P, et al. [Acamprosate--a stabilizing factor in long-term withdrawal of alcoholic patients]. *Therapeutische Umschau. Revue thérapeutique*. 1993;50(3):182-8. PMID: 8475472.
8. Mao YM, Zeng MD, Li YM, et al. [Capsule metadoxine in the treatment of alcoholic liver disease: a randomized, double-blind, placebo-controlled, multicenter study]. *Zhonghua gan zang bing za zhi = Zhonghua ganzangbing zazhi = Chinese journal of hepatology*. 2009;17(3):213-6. PMID: 19335986.
9. Ponce G, Sánchez-García J, Rubio G, et al. [Efficacy of naltrexone in the treatment of alcohol dependence disorder in women]. *Actas españolas de psiquiatria*. 2005;33(1):13-8. PMID: 15704026.
10. Pondé de-Sena E, Santos-Jesus R, Almeida Sarmiento C, et al. Use of carbamazepine-buspirone combination in alcohol dependence. *Jornal Brasileiro de Psiquiatria*. 1997;46(12):645-9. PMID: 1998-11798-003.
11. Roussaux JP, Hers D, Ferauge M. Does acamprosate influence alcohol consumption of weaned alcoholics? *Journal De Pharmacie De Belgique*. 1996;51(2):65-8. PMID: CN-00173028.

Not original research

1. Anderson N, Oliver MN. Oral topiramate effective for alcoholism. *J Fam Pract*. 2003;52(9):682-7.
2. Anton RF. Naltrexone for the management of alcohol dependence. *N Engl J Med*. 2008 Aug 14;359(7):715-21. PMID: 18703474.
3. Dongier M. What treatment options exist for alcohol abuse? *J Psychiatry Neurosci*. 2003 Jan;28(1):80. PMID: 12587852.

4. Drake RE, Mercer-McFadden C, Mueser KT, et al. Review of integrated mental health and substance abuse treatment for patients with dual disorders (Structured abstract). *Schizophr Bull*; 1998. p. 589-608.
5. Drtil J. Placebo therapy of alcohol dependent persons. *Homeost Health Dis*. 1997;38(4).
6. Fuller RK, Williford WO, Lee KK, et al. Veterans Administration cooperative study of disulfiram in the treatment of alcoholism: study design and methodological considerations. *Control Clin Trials*. 1984 Sep;5(3):263-73. PMID: 6386330.
7. Gache P, Hadengue A. Baclofen improves abstinence in alcoholic cirrhosis: Still better to come? *J Hepatol*. 2008;49(6):1083-5.
8. Gallant D. Reduction of ethanol intake by pharmacologic agents--investigational problems. *Alcohol Clin Exp Res*. 1992 Aug;16(4):836-7. PMID: 1530148.
9. Goldstein MZ, Pataki A, Webb MT. Alcoholism among elderly persons. *Psychiatr Serv*. 1996 Sep;47(9):941-3. PMID: 8875656.
10. Hussar DA. New drugs: acamprosate calcium and solifenacin succinate. *J Am Pharm Assoc* (2003). 2005 Jan-Feb;45(1):109-11. PMID: 15730126.
11. Johnson BA, Ait-Daoud N, Bowden CL, et al. Oral topiramate was effective as an adjunct to standardised medication compliance management in alcohol dependence. *Evid-Based Med*. 2004;9(1):24.
12. Kingsbury SJ, Salzman C. Disulfiram in the treatment of alcoholic patients with schizophrenia. *Hosp Community Psychiatry*. 1990;41(2):133-4.
13. Lesch OM, Walter H. Subtypes of alcoholism and their role in therapy. *Alcohol Alcohol*. 1996;31(Suppl 1):63-7. PMID: 8737003.
14. Lesch OM, Riegler A, Gutierrez K, et al. The European acamprosate trials: Conclusions for research and therapy. *J Biomed Sci*. 2001;8(1):89-95.
15. Leslie WS, Hankey CR, Lean ME. Weight gain as an adverse effect of some commonly prescribed drugs: a systematic review (Structured abstract). *QJM*. 2007;100(7):395-404. PMID: DARE-12007002731.
16. Luggen AS. Alcohol and the older adult. *Adv Nurse Pract*. 2006 Jan;14(1):47-52. PMID: 16425516.
17. Malec TS, Malec EA, Dongier M. Efficacy of buspirone in alcohol dependence: A review. *Alcoholism*. 1996;20(5):853-8.
18. Monti PM, Rohsenow DJ. Coping-skills training and cue-exposure therapy in the treatment of alcoholism. *Alcohol Research & Health*. 1999;23(2):107-15. PMID: 10890804.
19. O'Malley SS. Current strategies for the treatment of alcohol dependence in the United States. *Drug Alcohol Depend*. 1995;39(Suppl 1):S3-S7. PMID: 8565795.
20. O'Malley SS. Opioid antagonists in the treatment of alcohol dependence: Clinical efficacy and prevention of relapse. *Alcohol Alcohol*. 1996;31(Suppl 1):77-81. PMID: 8737005.
21. Peterson AM. Improving adherence in patients with alcohol dependence: A new role for pharmacists. *Am J Health Syst Pharm*. 2007;64(5 SUPPL.):S23-S9.
22. Swift RM. The pharmacotherapy of alcohol dependence: Clinical and economic aspects. *Econ Neurosci*. 2001;3(12):62-6.
23. Swift R. Emerging approaches to managing alcohol dependence. *Am J Health Syst Pharm*. 2007;64(5 Suppl):S12-S22.

Wrong population

1. Abtahi MA, Abtahi SH, Fazel F, et al. Topiramate and the vision: A systematic review. *Clinical Ophthalmology*. 2012;6(1):117-31.
2. Balldin J, Berggren U, Engel J, et al. Effect of citalopram on alcohol intake in heavy drinkers. *Alcohol Clin Exp Res*. 1994 Oct;18(5):1133-6. PMID: 7847595.

3. Bonnet U, Specka M, Leweke FM, et al. Gabapentin's acute effect on mood profile -- a controlled study on patients with alcohol withdrawal. *Prog Neuropsychopharmacol Biol Psychiatry*. 2007 Mar 30;31(2):434-8. PMID: 17178181.
4. Brahen LS, Capone T, Heller RC, et al. Controlled clinical study of naltrexone side effects comparing first-day doses and maintenance regimens. *Am J Drug Alcohol Abuse*. 1978;5(2):235-45. PMID: CN-00019998.
5. Brahen LS, Henderson RK, Capone T, et al. Naltrexone treatment in a jail work-release program. *J Clin Psychiatry*. 1984;45(9, Sect 2):49-52.
6. Carter AC, Miranda R, Gwaltney C, et al. Naltrexone's impact on alcohol expectancies in heavy drinkers. *Alcoholism*. 2010;34(6):177A.
7. Davidson D, Saha C, Scifres S, et al. Naltrexone and brief counseling to reduce heavy drinking in hazardous drinkers. *Addict Behav*. 2004 Aug;29(6):1253-8. PMID: 15236831.
8. Doty P, De Wit H. Effects of naltrexone pretreatment on the subjective and performance effects of ethanol in social drinkers. *Behav Pharmacol*. 1995;6(4):386-94.
9. Farwell WR, Stump TE, Wang J, et al. Weight gain and new onset diabetes associated with olanzapine and risperidone. *J Gen Intern Med*. 2004;19(12):1200-5.
10. Goodwin FK. From the Alcohol, Drug Abuse, and Mental Health Administration. *JAMA*. 1990 Mar 23-30;263(12):1610. PMID: 2155326.
11. Howland RH, Rush AJ, Wisniewski SR, et al. Concurrent anxiety and substance use disorders among outpatients with major depression: Clinical features and effect on treatment outcome. *Drug Alcohol Depend*. 2009;99(1-3):248-60. PMID: 18986774.
12. Johnson JL, Wiechelt SA, Ahmed AU, et al. Outcomes for substance user treatment in women: Results from the Baltimore Drug and Alcohol Treatment Outcomes Study. *Subst Use Misuse*. 2003;38(11-13):1807-29+920-922.
13. King A, Cao D, Vanier C, et al. Naltrexone decreases heavy drinking rates in smoking cessation treatment: an exploratory study. *Alcohol Clin Exp Res*. 2009 Jun;33(6):1044-50. PMID: 19302083.
14. Krylov EN. Psychotropic activity of the antialcohol preparation Proproten-100. *Bull Exp Biol Med*. 2003 Jan;135 Suppl 7:176-80. PMID: 12949691.
15. Malcolm R, Myrick LH, Veatch LM, et al. Self-reported sleep, sleepiness, and repeated alcohol withdrawals: a randomized, double blind, controlled comparison of lorazepam vs gabapentin. *J Clin Sleep Med*. 2007 Feb 15;3(1):24-32. PMID: 17557449.
16. Malla A. An outcome study comparing refusers and acceptors of treatment for alcoholism. *Can J Psychiatry*. 1988 Apr;33(3):183-7. PMID: 3383091.
17. Marrazzi MA, Wroblewski JM, Kinzie J, et al. High-dose naltrexone and liver function safety. *Am J Addict*. 1997;6(1):21-9.
18. Martin PR, Adinoff B, Lane E, et al. Fluvoxamine treatment of alcoholic amnesic disorder. *Eur Neuropsychopharmacol*. 1995 Mar;5(1):27-33. PMID: 7542052.
19. McGeary JE, Monti PM, Rohsenow DJ, et al. Genetic moderators of naltrexone's effects on alcohol cue reactivity. *Alcohol Clin Exp Res*. 2006 Aug;30(8):1288-96. PMID: 16899031.
20. Miranda R, Jr., MacKillop J, Monti PM, et al. Effects of topiramate on urge to drink and the subjective effects of alcohol: a preliminary laboratory study. *Alcohol Clin Exp Res*. 2008 Mar;32(3):489-97. PMID: 18215213.
21. Miranda R, MacKillop J, Meehan J, et al. Biobehavioral effects of topiramate among heavy drinkers. *Alcoholism*. 2011;35:21A.
22. Miranda R, Ray L, Reynolds E, et al. Effects of naltrexone on adolescent drinking. *Alcoholism*. 2012;36:89A.
23. Mitchell JM, Fields HL, White RL, et al. The Asp40 μ -Opioid Receptor Allele Does Not Predict Naltrexone Treatment Efficacy in Heavy Drinkers. *J Clin Psychopharmacol*. 2007;27(1):112-5. PMID: 17224736.

24. Mitchell JM, Teague CH, Kayser AS, et al. Varenicline decreases alcohol consumption in heavy-drinking smokers. *Psychopharmacology (Berl)*. 2012 Oct;223(3):299-306. PMID: 22547331.
25. Myrick H, Malcolm R, Randall PK, et al. A double-blind trial of gabapentin versus lorazepam in the treatment of alcohol withdrawal. *Alcohol Clin Exp Res*. 2009 Sep;33(9):1582-8. PMID: 19485969.
26. Naranjo CA, Sellers EM, Sullivan JT, et al. The serotonin uptake inhibitor citalopram attenuates ethanol intake. *Clin Pharmacol Ther*. 1987 Mar;41(3):266-74. PMID: 3469057.
27. Naranjo CA, Poulos CX, Bremner KE, et al. Fluoxetine attenuates alcohol intake and desire to drink. *Int Clin Psychopharmacol*. 1994 Sep;9(3):163-72. PMID: 7814825.
28. Niederhofer H, Staffen W, Mair A. Comparison of naltrexone and placebo in treatment of alcohol dependence of adolescents. *Alcohol Treat Q*. 2003;21(2):87-95.
29. Nunes EV, Levin FR. Treatment of Depression in Patients with Alcohol or Other Drug Dependence: A Meta-analysis. *JAMA*. 2004;291(15):1887-96.
30. O'Carroll RE, Moffoot AP, Ebmeier KP, et al. Effects of fluvoxamine treatment on cognitive functioning in the alcoholic Korsakoff syndrome. *Psychopharmacology (Berl)*. 1994 Sep;116(1):85-8. PMID: 7862935.
31. O'Malley SS, Krishnan-Sarin S, McKee SA, et al. Dose-dependent reduction of hazardous alcohol use in a placebo-controlled trial of naltrexone for smoking cessation. *Int J Neuropsychopharmacol*. 2009 Jun;12(5):589-97. PMID: 18796184.
32. O'Malley S. Naltrexone for heavy drinking in young adults. *Alcohol*. 2012;36:334A.
33. Perugi G, Toni C, Frare F, et al. Effectiveness of adjunctive gabapentin in resistant bipolar disorder: is it due to anxious-alcohol abuse comorbidity? *J Clin Psychopharmacol*. 2002 Dec;22(6):584-91. PMID: 12454558.
34. Petrakis IL, Carroll KM, Nich C, et al. Disulfiram treatment for cocaine dependence in methadone-maintained opioid addicts. *Addiction*. 2000;95(2):219-28. PMID: 10723850.
35. Prochaska JJ, Delucchi K, Hall SM. A meta-analysis of smoking cessation interventions with individuals in substance abuse treatment or recovery (Structured abstract). *J Consult Clin Psychol*; 2004. p. 1144-56.
36. Simon GE, Heiligenstein J, Revicki D. More people with depression continued treatment with fluoxetine than with desipramine or imipramine. *Evid Based Med*. 2000;5(2):51.
37. Spies CD, Dubisz N, Neumann T, et al. Therapy of alcohol withdrawal syndrome in intensive care unit patients following trauma: results of a prospective, randomized trial. *Crit Care Med*. 1996 Mar;24(3):414-22. PMID: 8625628.
38. Tauscher-Wisniewski S, Disch D, Plewes J, et al. Evaluating suicide-related adverse events in clinical trials of fluoxetine treatment in adults for indications other than major depressive disorder. *Psychol Med*. 2007;37(11):1585-93.

Wrong intervention

1. Ait-Daoud N, Johnson BA, Javors M, et al. Combining ondansetron and naltrexone treats biological alcoholics: corroboration of self-reported drinking by serum carbohydrate deficient transferrin, a biomarker. *Alcohol Clin Exp Res*. 2001 Jun;25(6):847-9. PMID: 11410720.
2. Ait-Daoud N, Johnson BA, Prihoda TJ, et al. Combining ondansetron and naltrexone reduces craving among biologically predisposed alcoholics: preliminary clinical evidence. *Psychopharmacology (Berl)*. 2001 Feb;154(1):23-7. PMID: 11292002.

3. Allen JP, Sillamauke P, Anton R. Contribution of carbohydrate deficient transferrin to gamma glutamyl transpeptidase in evaluating progress of patients in treatment for alcoholism. *Alcohol Clin Exp Res*. 1999 Jan;23(1):115-20. PMID: 10029211.
4. Anton RF, Myrick H, Baros AM, et al. Efficacy of a combination of flumazenil and gabapentin in the treatment of alcohol dependence: relationship to alcohol withdrawal symptoms. *J Clin Psychopharmacol*. 2009 Aug;29(4):334-42. PMID: 19593171.
5. Collins GB, Janesz JW, Byerly-Thrope J. The Cleveland Clinic Alcohol Rehabilitation Program: A treatment outcome study. A preliminary report. *Cleve Clin Q*. 1985;52(2):245-51.
6. DiClemente CC, Doyle SR, Donovan D. Predicting treatment seekers' readiness to change their drinking behavior in the COMBINE study. *Alcohol*. 2009;33(5):879-92.
7. Dinh-Zarr Tho B, Goss Cynthia W, Heitman E, et al. Interventions for preventing injuries in problem drinkers. *Cochrane Database Syst Rev*. 2004(3)PMID: CD001857.
8. Epstein EE, Rhines KC, Cook S, et al. Changes in alcohol craving and consumption by phase of menstrual cycle in alcohol dependent women. *J Subst Use*. 2006;11(5):323-32. PMID: 2009347535.
9. Fantozzi R, Caramelli L, Ledda F, et al. Biological markers and therapeutic outcome in alcoholic disease: a twelve-year survey. *Klin Wochenschr*. 1987 Jan 5;65(1):27-33. PMID: 3560786.
10. Guerrini I, Gentili C, Nelli G, et al. A follow up study on the efficacy of metadoxine in the treatment of alcohol dependence. *Subst Abuse Treat Prev Policy*. 2006;1:35. PMID: 17176456.
11. Hasin DS, Endicott J, Keller MB. Alcohol problems in psychiatric patients: 5-year course. *Compr Psychiatry*. 1991 Jul-Aug;32(4):303-16. PMID: 1935019.
12. Jensen SB, Christoffersen CB, Noerregaard A. Apomorphine in outpatient treatment of alcohol intoxication and abstinence: a double-blind study. *Br J Addict Alcohol Other Drugs*. 1977 Dec;72(4):325-30. PMID: 341937.
13. Johnsen J, Morland J. Disulfiram implant: a double-blind placebo controlled follow-up on treatment outcome. *Alcohol Clin Exp Res*. 1991 Jun;15(3):532-6. PMID: 1877740.
14. Johnson BA, Ait-Daoud N, Prihoda TJ. Combining ondansetron and naltrexone effectively treats biologically predisposed alcoholics: from hypotheses to preliminary clinical evidence. *Alcohol Clin Exp Res*. 2000 May;24(5):737-42. PMID: 10832917.
15. Landabaso MA, Iraurgi I, Sanz J, et al. Naltrexone in the treatment of alcoholism. Two-year follow up results. *Eur J Psychiat*. 1999;13(2):97-105.
16. Maremmani AGI, Pani PP, Rovai L, et al. Long-term (gamma)-hydroxybutyric acid (GHB) and disulfiram combination therapy in GHB treatment-resistant chronic alcoholics. *Int J Environmen Res Public Health*. 2011;8(7):2816-27.
17. Moak DH, Anton RF, Malcolm R, et al. Alcoholic subjects with anxiety disorder: Characteristics of completers and noncompleters in a pharmacologic study. *Am J Addict*. 1993;2(1):39-47.
18. Oslin DW. Treatment of late-life depression complicated by alcohol dependence. *Am J Geriatr Psychiatry*. 2005 Jun;13(6):491-500. PMID: 15956269.
19. Witte J, Bentley K, Evins AE, et al. A randomized, controlled, pilot study of acamprosate added to escitalopram in adults with major depressive disorder and alcohol use disorder. *J Clin Psychopharmacol*. 2012;32(6):787-96. PMID: 23131884.

Wrong comparator

1. Agosti V, Nunes EV, O'Shea D. Do manualized psychosocial interventions help reduce relapse among alcohol-dependent adults treated with naltrexone or placebo? A meta-analysis. *Am J Addict*. 2012;21(6):501-7.
2. Aguiar P, Neto D, Lambaz R, et al. Prognostic factors during outpatient treatment for alcohol dependence: Cohort study with 6 months of treatment follow-up. *Alcohol Alcohol*. 2012;47(6):702-10.
3. Aliyev NN. Trial of interferon in chronic alcoholism. *Psychiatry Res*. 1994;54(3):307-8. PMID: 7792335.
4. Altintoprak AE, Zorlu N, Coskunol H, et al. Effectiveness and tolerability of mirtazapine and amitriptyline in alcoholic patients with co-morbid depressive disorder: a randomized, double-blind study. *Hum Psychopharmacol*. 2008 Jun;23(4):313-9. PMID: 18327889.
5. Bergstrom B, Ohlin H, Lindblom PE, et al. Is disulfiram implantation effective? *Lancet*. 1982 Jan 2;1(8262):49-50. PMID: 6119442.
6. Brady KT, Sonne SC, Roberts JM. Sertraline treatment of comorbid posttraumatic stress disorder and alcohol dependence. *J Clin Psychiatry*. 1995 Nov;56(11):502-5. PMID: 7592501.
7. Capone C, Kahler CW, Swift RM, et al. Does family history of alcoholism moderate naltrexone's effects on alcohol use? *J Stud Alcohol Drugs*. 2011 Jan;72(1):135-40. PMID: 21138720.
8. Caputo F, Addolorato G, Stoppo M, et al. Comparing and combining gamma-hydroxybutyric acid (GHB) and naltrexone in maintaining abstinence from alcohol: an open randomised comparative study. *Eur Neuropsychopharmacol*. 2007 Dec;17(12):781-9. PMID: 17611081.
9. Carroll KM, Nich C, Ball SA, et al. Treatment of cocaine and alcohol dependence with psychotherapy and disulfiram. *Addiction*. 1998 May;93(5):713-27. PMID: 9692270.
10. Croissant B, Klein O, Gehrlein L, et al. Quetiapine in relapse prevention in alcoholics suffering from craving and affective symptoms: a case series. *Eur Psychiatry*. 2006 Dec;21(8):570-3. PMID: 17161284.
11. Croissant B, Diehl A, Klein O, et al. A pilot study of oxcarbazepine versus acamprosate in alcohol-dependent patients. *Alcohol Clin Exp Res*. 2006 Apr;30(4):630-5. PMID: 16573580.
12. Croop RS, Faulkner EB, Labriola DF. The safety profile of naltrexone in the treatment of alcoholism. Results from a multicenter usage study. The Naltrexone Usage Study Group. *Arch Gen Psychiatry*. 1997 Dec;54(12):1130-5. PMID: 9400350.
13. Donovan D, Mattson ME, Cisler RA, et al. Quality of life as an outcome measure in alcoholism treatment research. *J Stud Alcohol*. 2005;66(SUPPL. 15):119-39.
14. Feeney GF, Connor JP, Mc DYR, et al. Is acamprosate use in alcohol dependence treatment reflected in improved subjective health status outcomes beyond cognitive behavioural therapy alone? *J Addict Dis*. 2006;25(4):49-58. PMID: 17088225.
15. Fernandez Miranda JJ, Marina Gonzalez PA, Montes Perez M, et al. Topiramate as add-on therapy in non-respondent alcohol dependant patients: a 12 month follow-up study. *Actas Esp Psiquiatr*. 2007 Jul-Aug;35(4):236-42. PMID: 17592785.
16. Flannery BA, Poole SA, Gallop RJ, et al. Alcohol craving predicts drinking during treatment: an analysis of three assessment instruments. *J Stud Alcohol*. 2003 Jan;64(1):120-6. PMID: 12608492.
17. Fucito LM, Toll BA, Wu R, et al. A preliminary investigation of varenicline for heavy drinking smokers. *Psychopharmacology (Berl)*. 2011 Jun;215(4):655-63. PMID: 21221531.
18. Gopalakrishnan R, Ross J, O'Brien C, et al. Course of late-life depression with alcoholism following combination therapy. *J Stud Alcohol Drugs*. 2009 Mar;70(2):237-41. PMID: 19261235.

19. Guardia J, Roncero C, Galan J, et al. A double-blind, placebo-controlled, randomized pilot study comparing quetiapine with placebo, associated to naltrexone, in the treatment of alcohol-dependent patients. *Addict Behav.* 2011 Mar;36(3):265-9. PMID: 21146937.
20. Janiri L, Martinotti G, Di Nicola M. Aripiprazole for relapse prevention and craving in alcohol-dependent subjects: Results from a pilot study. *J Clin Psychopharmacol.* 2007;27(5):519-20. PMID: 17873691.
21. Kampman KM, Pettinati HM, Lynch KG, et al. Initiating acamprosate within-detoxification versus post-detoxification in the treatment of alcohol dependence. *Addict Behav.* 2009 Jun-Jul;34(6-7):581-6. PMID: 19345510.
22. Kiritze-Topor P, Huas D, Rosenzweig C, et al. A pragmatic trial of acamprosate in the treatment of alcohol dependence in primary care. *Alcohol Alcohol.* 2004 Nov-Dec;39(6):520-7. PMID: 15304381.
23. Knox WJ. Four-year follow-up of veterans treated on a small alcoholism treatment ward. *Q J Stud Alcohol.* 1972 Mar;33(1):105-10. PMID: 4551019.
24. Kranzler HR, Tennen H, Penta C, et al. Targeted naltrexone treatment of early problem drinkers. *Addict Behav.* 1997 May-Jun;22(3):431-6. PMID: 9183513.
25. Kranzler HR, Pierucci-Lagha A, Feinn R, et al. Effects of ondansetron in early- versus late-onset alcoholics: a prospective, open-label study. *Alcohol Clin Exp Res.* 2003 Jul;27(7):1150-5. PMID: 12878921.
26. Lewis DC. The clinical usefulness of narcotic antagonists: preliminary findings on the use of naltrexone. *Am J Drug Alcohol Abuse.* 1975;2(3-4):403-15. PMID: 1227300.
27. Liebson I, Bigelow G. A behavioural-pharmacological treatment of dually addicted patients. *Behav Res Ther.* 1972 Nov;10(4):403-5. PMID: 4637497.
28. Likhitsathian S, Saengcharnchai P, Uttawichai K, et al. Cognitive changes in topiramate-treated patients with alcoholism: a 12-week prospective study in patients recently detoxified. *Psychiatry Clin Neurosci.* 2012 Apr;66(3):235-41. PMID: 22443246.
29. Martinotti G, Di Nicola M, Romanelli R, et al. High and low dosage oxcarbazepine versus naltrexone for the prevention of relapse in alcohol-dependent patients. *Hum Psychopharmacol.* 2007 Apr;22(3):149-56. PMID: 17397097.
30. Martinotti G, Di Nicola M, Tedeschi D, et al. Pregabalin versus naltrexone in alcohol dependence: a randomised, double-blind, comparison trial. *J Psychopharmacol.* 2010 Sep;24(9):1367-74. PMID: 19346279.
31. McRae-Clark AL, Verduin ML, Tolliver BK, et al. An open-label trial of aripiprazole treatment in dual diagnosis individuals: Safety and efficacy. *J Dual Diagn.* 2009;5(1):83-96.
32. Monnelly EP, Ciraulo DA, Knapp C, et al. Quetiapine for treatment of alcohol dependence. *J Clin Psychopharmacol.* 2004 Oct;24(5):532-5. PMID: 15349010.
33. Morgan MY, Landron F, Leher P. Improvement in quality of life after treatment for alcohol dependence with acamprosate and psychosocial support. *Alcohol Clin Exp Res.* 2004 Jan;28(1):64-77. PMID: 14745303.
34. Muhonen LH, Lahti J, Sinclair D, et al. Treatment of alcohol dependence in patients with co-morbid major depressive disorder--predictors for the outcomes with memantine and escitalopram medication. *Subst Abuse Treat Prev Policy.* 2008;3:20. PMID: 18834506.
35. Muhonen LH, Lonnqvist J, Juva K, et al. Double-blind, randomized comparison of memantine and escitalopram for the treatment of major depressive disorder comorbid with alcohol dependence. *J Clin Psychiatry.* 2008 Mar;69(3):392-9. PMID: 18348597.

36. Murthy KK, Praveenlal K. An experience with disulfiram in the management of Alcohol Dependence Syndrome: I. Side effects of disulfiram and symptoms of alcohol-disulfiram reaction. *Ind J Psychol Med.* 1988;11(2):145-8.
37. Ojehagen A, Skjaeris A, Berglund M. Long-term use of aversive drugs in outpatient alcoholism treatment. *Acta Psychiatr Scand.* 1991 Aug;84(2):185-90. PMID: 1950615.
38. O'Malley SS, Garbutt JC, Gastfriend DR, et al. Efficacy of extended-release naltrexone in alcohol-dependent patients who are abstinent before treatment. *J Clin Psychopharmacol.* 2007 Oct;27(5):507-12. PMID: 17873686.
39. Oroszi G, Anton RF, O'Malley S, et al. OPRM1 Asn40Asp predicts response to naltrexone treatment: a haplotype-based approach. *Alcohol Clin Exp Res.* 2009 Mar;33(3):383-93. PMID: 19053977.
40. Palatty PL, Saldanha E. Status of disulfiram in present day alcoholic deaddiction therapy. *Indian J Psychiatry.* 2011;53(1):25-9.
41. Powell BJ, Penick EC, Read MR, et al. Comparison of three outpatient treatment interventions: a twelve-month follow-up of men alcoholics. *J Stud Alcohol.* 1985 Jul;46(4):309-12. PMID: 2993750.
42. Ray LA, Oslin DW. Naltrexone for the treatment of alcohol dependence among African Americans: results from the COMBINE Study. *Drug Alcohol Depend.* 2009 Dec 1;105(3):256-8. PMID: 19717248.
43. Reid SC, Teesson M, Sannibale C, et al. The efficacy of compliance therapy in pharmacotherapy for alcohol dependence: a randomized controlled trial. *J Stud Alcohol.* 2005 Nov;66(6):833-41. PMID: 16459945.
44. Roy A. Treating depression among alcoholics. *Can J Psychiatry.* 1996;41(3):194-5.
45. Rubio G, Ponce G, Jimenez-Arriero MA, et al. Effects of topiramate in the treatment of alcohol dependence. *Pharmacopsychiatry.* 2004 Jan;37(1):37-40. PMID: 14750047.
46. Rubio G, Ponce G, Rodriguez-Jimenez R, et al. Clinical predictors of response to naltrexone in alcoholic patients: who benefits most from treatment with naltrexone? *Alcohol Alcohol.* 2005 May-Jun;40(3):227-33. PMID: 15797885.
47. Ulrichsen J, Nielsen MK, Ulrichsen M. Disulfiram in severe alcoholism: an open controlled study. *Nord J Psychiatry.* 2010;64(6):356-62.
48. Wilens TE, Adler LA, Tanaka Y, et al. Correlates of alcohol use in adults with ADHD and comorbid alcohol use disorders: exploratory analysis of a placebo-controlled trial of atomoxetine. *Curr Med Res Opin.* 2011 Dec;27(12):2309-20. PMID: 22029549.

Wrong outcome

1. Testing combined pharmacotherapies and behavioral interventions in alcohol dependence: Rationale and methods. *Alcohol.* 2003;27(7):1107-22.
2. Altamura AC, Mauri MC, Girardi T, et al. Alcoholism and depression: a placebo controlled study with viloxazine. *Int J Clin Pharmacol Res.* 1990;10(5):293-8. PMID: 2079386.
3. Anton RF, Moak DH, Latham PK. The obsessive compulsive drinking scale: A new method of assessing outcome in alcoholism treatment studies. *Arch Gen Psychiatry.* 1996 Mar;53(3):225-31. PMID: 8611059.
4. Armeli S, Feinn R, Tennen H, et al. The effects of naltrexone on alcohol consumption and affect reactivity to daily interpersonal events among heavy drinkers. *Exp Clin Psychopharmacol.* 2006 May;14(2):199-208. PMID: 16756424.
5. Back SE, Brady KT, Sonne SC, et al. Symptom improvement in co-occurring PTSD and alcohol dependence. *J Nerv Ment Dis.* 2006 Sep;194(9):690-6. PMID: 16971821.
6. Baltieri DA, de Andrade AG. Efficacy of acamprosate in the treatment of alcohol-dependent outpatients. *Rev Bras Psiquiatr.* 2003 Sep;25(3):156-9. PMID: 12975689.

7. Baltieri DA, Correa Filho JM. Role of two clusters of male alcoholics in treatment retention. *Eur Addict Res.* 2012;18(4):201-11.
8. Baros AM, Latham PK, Moak DH, et al. What role does measuring medication compliance play in evaluating the efficacy of naltrexone? *Alcohol Clin Exp Res.* 2007 Apr;31(4):596-603. PMID: 17374038.
9. Beresford TP, Arciniegas D, Clapp L, et al. Reduction of affective lability and alcohol use following traumatic brain injury: a clinical pilot study of anti-convulsant medications. *Brain Inj.* 2005 Apr;19(4):309-13. PMID: 15832875.
10. Bujarski S, O'Malley SS, Lunney K, et al. The effects of drinking goal on treatment outcome for alcoholism. *J Consult Clin Psychol.* 2013;81(1):13-22. PMID: 23231573.
11. Chen J, Johnson BA, Wang XQ, et al. Trajectory analyses in alcohol treatment research. *Alcohol Clin Exp Res.* 2012 Aug;36(8):1442-8. PMID: 22525000.
12. Corbin WR, Scott C, Leeman RF, et al. Early Subjective Response and Acquired Tolerance as Predictors of Alcohol Use and Related Problems in a Clinical Sample. *Alcohol.* 2013.
13. Cornelius JR, Salloum IM, Cornelius MD, et al. Preliminary report: double-blind, placebo-controlled study of fluoxetine in depressed alcoholics. *Psychopharmacol Bull.* 1995;31(2):297-303. PMID: 7491382.
14. Cramer J, Rosenheck R, Kirk G, et al. Medication compliance feedback and monitoring in a clinical trial: predictors and outcomes. *Value Health.* 2003 Sep-Oct;6(5):566-73. PMID: 14627063.
15. Cutler RB. Abatement of craving in recovering alcoholics: A descriptive analysis. *Addict Res Theory.* 2005;13(2):111-27.
16. Falk D, Wang XQ, Liu L, et al. Percentage of subjects with no heavy drinking days: Evaluation as an efficacy endpoint for alcoholclinical trials. *Alcohol.* 2010;34(12):2022-34.
17. Feinn R, Tennen H, Cramer J, et al. Measurement and prediction of medication compliance in problem drinkers. *Alcohol Clin Exp Res.* 2003 Aug;27(8):1286-92. PMID: 12966323.
18. Flannery BA, Volpicelli JR, Pettinati HM. Psychometric properties of the Penn Alcohol Craving Scale. *Alcohol Clin Exp Res.* 1999 Aug;23(8):1289-95. PMID: 10470970.
19. Fox HC, Anderson GM, Tuit K, et al. Prazosin effects on stress- and cue-induced craving and stress response in alcohol-dependent individuals: preliminary findings. *Alcohol Clin Exp Res.* 2012 Feb;36(2):351-60. PMID: 21919922.
20. Frick KM, Loessl B, Brueck RK, et al. What works for patients in outpatient treatment for alcohol addiction? an explorative study into clients' evaluation of subjective factors and therapy satisfaction. *Subst Abuse.* 2011;5(1):27-34.
21. Fuller R, Roth H, Long S. Compliance with disulfiram treatment of alcoholism. *J Chronic Dis.* 1983;36(2):161-70. PMID: 6337171.
22. George DT, Phillips MJ, Lifshitz M, et al. Fluoxetine treatment of alcoholic perpetrators of domestic violence: A 12-week, double-blind, randomized, placebo-controlled intervention study. *J Clin Psychiatry.* 2011;72(1):60-5.
23. Gueorguieva R, Wu R, Donovan D, et al. Naltrexone and combined behavioral intervention effects on trajectories of drinking in the COMBINE study. *Drug Alcohol Depend.* 2010 Mar 1;107(2-3):221-9. PMID: 19969427.
24. Hernandez-Avila CA, Burleson JA, Kranzler HR. Stage of change as a predictor of abstinence among alcohol-dependent subjects in pharmacotherapy trials. *Substance Abuse.* 1998;19(2):81-91.
25. Jaffe AJ, Rounsaville B, Chang G, et al. Naltrexone, relapse prevention, and supportive therapy with alcoholics: an analysis of patient treatment matching. *J Consult Clin Psychol.* 1996 Oct;64(5):1044-53. PMID: 8916634.

26. Johnson BA, Roache JD, Ait-Daoud N, et al. Ondansetron reduces the craving of biologically predisposed alcoholics. *Psychopharmacology (Berl)*. 2002 Apr;160(4):408-13. PMID: 11919668.
27. Johnson BA, Ait-Daoud N, Ma JZ, et al. Ondansetron reduces mood disturbance among biologically predisposed, alcohol-dependent individuals. *Alcohol Clin Exp Res*. 2003 Nov;27(11):1773-9. PMID: 14634493.
28. Kiefer F, Jahn H, Otte C, et al. Hypothalamic-pituitary-adrenocortical axis activity: a target of pharmacological anticraving treatment? *Biol Psychiatry*. 2006 Jul 1;60(1):74-6. PMID: 16483549.
29. Kranzler HR, Burleson JA, Brown J, et al. Fluoxetine treatment seems to reduce the beneficial effects of cognitive-behavioral therapy in type B alcoholics. *Alcohol Clin Exp Res*. 1996 Dec;20(9):1534-41. PMID: 8986200.
30. Kravitz HM, Fawcett J, McGuire M, et al. Treatment attrition among alcohol-dependent men: is it related to novelty seeking personality traits? *J Clin Psychopharmacol*. 1999 Feb;19(1):51-6. PMID: 9934943.
31. Krupitsky EM, Burakov AM, Ivanov VB, et al. Baclofen administration for the treatment of affective disorders in alcoholic patients. *Drug Alcohol Depend*. 1993 Sep;33(2):157-63. PMID: 8261880.
32. Labbate LA, Sonne SC, Randal CL, et al. Does comorbid anxiety or depression affect clinical outcomes in patients with post-traumatic stress disorder and alcohol use disorders? *Compr Psychiatry*. 2004 Jul-Aug;45(4):304-10. PMID: 15224273.
33. Lapham S, Forman R, Alexander M, et al. The effects of extended-release naltrexone on holiday drinking in alcohol-dependent patients. *J Subst Abuse Treat*. 2009 Jan;36(1):1-6. PMID: 18775624.
34. Martin PR, Loewy J, Liou S, et al. Correlation of Serum Gamma-Glutamyl Transferase With Alcohol Consumption. 158th Annual Meeting of the American Psychiatric Association; 2005 May 21-26; Atlanta, GA; 2005. p. Nr215.
35. Mason BJ, Kocsis JH. Desipramine treatment of alcoholism. *Psychopharmacol Bull*. 1991;27(2):155-61. PMID: 1924663.
36. Mason BJ, Light JM, Williams LD, et al. Proof-of-concept human laboratory study for protracted abstinence in alcohol dependence: effects of gabapentin. *Addict Biol*. 2009 Jan;14(1):73-83. PMID: 18855801.
37. Meszaros K, Willinger U, Fischer G, et al. The tridimensional personality model: influencing variables in a sample of detoxified alcohol dependents. *European Fluvoxamine in Alcoholism Study Group. Compr Psychiatry*. 1996 Mar-Apr;37(2):109-14. PMID: 8654059.
38. Muhonen LH, Lahti J, Alho H, et al. Serotonin transporter polymorphism as a predictor for escitalopram treatment of major depressive disorder comorbid with alcohol dependence. *Psychiatry Res*. 2011 Mar 30;186(1):53-7. PMID: 20800901.
39. O'Brien CP, Volpicelli LA, Volpicelli JR. Naltrexone in the treatment of alcoholism: a clinical review. *Alcohol*. 1996 Jan-Feb;13(1):35-9. PMID: 8837932.
40. O'Malley SS, Jaffe AJ, Rode S, et al. Experience of a 'slip' among alcoholics treated with naltrexone or placebo. *Am J Psychiatry*. 1996;153(2):281-3.
41. Ooteman W, Naassila M, Koeter MW, et al. Predicting the effect of naltrexone and acamprosate in alcohol-dependent patients using genetic indicators. *Addict Biol*. 2009 Jul;14(3):328-37. PMID: 19523047.
42. Oslin D, Liberto JG, O'Brien J, et al. Tolerability of naltrexone in treating older, alcohol-dependent patients. *Am J Addict*. 1997 Summer;6(3):266-70. PMID: 9256993.
43. Palliyath S, Schwartz BD. Disulfiram neuropathy: electrophysiological study. *Electromyogr Clin Neurophysiol*. 1988 Jun-Jul;28(5):245-7. PMID: 2847909.
44. Ray LA, Hutchison KE, Bryan A. Psychosocial predictors of treatment outcome, dropout, and change processes in a pharmacological clinical trial for alcohol dependence. *Addict Disord Their Treat*. 2006;5(4):179-90.

45. Richardson K, Baillie A, Reid S, et al. Do acamprosate or naltrexone have an effect on daily drinking by reducing craving for alcohol? *Addiction*. 2008 Jun;103(6):953-9. PMID: 18482418.
46. Rohsenow DJ, Colby SM, Monti PM, et al. Predictors of compliance with naltrexone among alcoholics. *Alcohol Clin Exp Res*. 2000 Oct;24(10):1542-9. PMID: 11045863.
47. Rosenthal RN, Perkel C, Singh P, et al. A pilot open randomized trial of valproate and phenobarbital in the treatment of acute alcohol withdrawal. *Am J Addict*. 1998 Summer;7(3):189-97. PMID: 9702286.
48. Roy A. Placebo-controlled study of sertraline in depressed recently abstinent alcoholics. *Biol Psychiatry*. 1998 Oct 1;44(7):633-7. PMID: 9787889.
49. Schaumberg K, Kuerbis A, Morgenstern J, et al. Attributions of Change and Self-Efficacy in a Randomized Controlled Trial of Medication and Psychotherapy for Problem Drinking. *Behav Ther*. 2013;44(1):88-99.
50. Scott C, Corbin WR, Leeman RF, et al. The influence of initial subjective response and acquired tolerance to alcohol on drinking behavior and problems in a clinical sample. *Alcohol*. 2011;35:165A.
51. Seneviratne C, Johnson BA. Serotonin transporter genomic biomarker for quantitative assessment of ondansetron treatment response in alcoholics. *Front Psychiatry*. 2012;3PMID: 22470354.
52. Smith Erica J, Lui S, Terplan M. Pharmacologic Interventions for Pregnant Women Enrolled in Alcohol Treatment. *Cochrane Database Syst Rev*: John Wiley & Sons, Ltd; 2009.
53. Snyder S, Karacan I, Salis PJ. Disulfiram and nocturnal penile tumescence in the chronic alcoholic. *Biol Psychiatry*. 1981 Apr;16(4):399-406. PMID: 7225493.
54. Soyka M, Sass H. Acamprosate: a new pharmacotherapeutic approach to relapse prevention in alcoholism--preliminary data. *Alcohol Alcohol Suppl*. 1994;2:531-6. PMID: 8974379.
55. Specka M, Lieb B, Kuhlmann T, et al. Marked reduction of heavy drinking did not reduce nicotine use over 1 year in a clinical sample of alcohol-dependent patients. *Pharmacopsychiatry*. 2011;44(3):120-1. PMID: 21298613.
56. Staner L, Boeijinga P, Danel T, et al. Effects of acamprosate on sleep during alcohol withdrawal: A double-blind placebo-controlled polysomnographic study in alcohol-dependent subjects. *Alcohol Clin Exp Res*. 2006 Sep;30(9):1492-9. PMID: 16930211.
57. Stapleton JM, Eckardt MJ, Martin P, et al. Treatment of alcoholic organic brain syndrome with the serotonin reuptake inhibitor fluvoxamine: a preliminary study. *Adv Alcohol Subst Abuse*. 1988;7(3-4):47-51. PMID: 2464912.
58. Suh JJ, Pettinati HM, Kampman KM, et al. Gender differences in predictors of treatment attrition with high dose naltrexone in cocaine and alcohol dependence. *Am J Addict*. 2008 Nov-Dec;17(6):463-8. PMID: 19034737.
59. Swearingen CE, Moyer A, Finney JW. Alcoholism treatment outcome studies, 1970-1998: An expanded look at the nature of the research. *Addict Behav*. 2003;28(3):415-36.
60. Thevos AK, Brown JM, Malcolm R, et al. Alcohol treatment: measurement of effectiveness by global outcome. *Soc Work Health Care*. 1996;23(3):57-71. PMID: 8865515.
61. Vaughan MD, Hook JN, Wagley JN, et al. Changes in affect and drinking outcomes in a pharmacobehavioral trial for alcohol dependence. *Addict Disord Their Treat*. 2012;11(1):14-25.
62. Volpicelli JR, Alterman AI, Hayashida M, et al. Naltrexone in the treatment of alcohol dependence. *Arch Gen Psychiatry*. 1992 Nov;49(11):876-80. PMID: 1345133.
63. Weinrieb RM, Van Horn DH, McLellan AT, et al. Alcoholism treatment after liver transplantation: lessons learned from a clinical trial that failed. *Psychosomatics*. 2001 Mar-Apr;42(2):110-6. PMID: 11239123.

64. Witkiewitz K. Predictors of heavy drinking during and following treatment. *Psychol Addict Behav*. 2011;25(3):426-38.
65. Zarkin GA, Bray JW, Aldridge A, et al. The effect of alcohol treatment on social costs of alcohol dependence: results from the COMBINE study. *Med Care*. 2010 May;48(5):396-401. PMID: 20393362.

Wrong setting

1. Christensen JK. Side-effects after Antabuse--myths or reality? *Br J Clin Pract Suppl*. 1984;36:21-8. PMID: 6437426.
2. Drobos DJ, Anton RF. Drinking in alcoholics following an alcohol challenge research protocol. *J Stud Alcohol*. 2000 Mar;61(2):220-4. PMID: 10757131.
3. Drobos DJ, Anton RF, Thomas SE, et al. A clinical laboratory paradigm for evaluating medication effects on alcohol consumption: naltrexone and nalmefene. *Neuropsychopharmacology*. 2003 Apr;28(4):755-64. PMID: 12655322.
4. Goyer PF, Major LF. Hepatotoxicity in disulfiram-treated patients. *J Stud Alcohol*. 1979;40(1):133-7. PMID: 449328.
5. Hameedi FA, Rosen MI, McCance-Katz EF, et al. Behavioral, physiological, and pharmacological interaction of cocaine and disulfiram in humans. *Biol Psychiatry*. 1995 Apr 15;37(8):560-3. PMID: 7619981.
6. Hutchison KE, Wooden A, Swift RM, et al. Olanzapine reduces craving for alcohol: a DRD4 VNTR polymorphism by pharmacotherapy interaction. *Neuropsychopharmacology*. 2003 Oct;28(10):1882-8. PMID: 12888781.
7. King AC, Volpicelli JR, Frazer A, et al. Effect of naltrexone on subjective alcohol response in subjects at high and low risk for future alcohol dependence. *Psychopharmacology (Berl)*. 1997 Jan;129(1):15-22. PMID: 9122358.
8. Knox PC, Donovan DM. Using naltrexone in inpatient alcoholism treatment. *J Psychoactive Drugs*. 1999 Oct-Dec;31(4):373-88. PMID: 10681104.
9. Lake CR, Major LF, Ziegler MG, et al. Increased sympathetic nervous system activity in alcoholic patients treated with disulfiram. *Am J Psychiatry*. 1977 Dec;134(12):1411-4. PMID: 920841.
10. McKee SA, Harrison EL, O'Malley SS, et al. Varenicline reduces alcohol self-administration in heavy-drinking smokers. *Biol Psychiatry*. 2009 Jul 15;66(2):185-90. PMID: 19249750.
11. Myrick H, Anton R, Voronin K, et al. A double-blind evaluation of gabapentin on alcohol effects and drinking in a clinical laboratory paradigm. *Alcohol Clin Exp Res*. 2007 Feb;31(2):221-7. PMID: 17250613.
12. Naranjo CA, Poulos CX, Bremner KE, et al. Citalopram decreases desirability, liking, and consumption of alcohol in alcohol-dependent drinkers. *Clin Pharmacol Ther*. 1992 Jun;51(6):729-39. PMID: 1535302.
13. Peachey JE, Zilm DH, Robinson GM, et al. A placebo-controlled double-blind comparative clinical study of the disulfiram- and calcium carbimide-acetaldehyde mediated ethanol reactions in social drinkers. *Alcohol Clin Exp Res*. 1983 Spring;7(2):180-7. PMID: 6346921.
14. Snyder S, Keeler M. Acute effects of disulfiram on anxiety levels of chronic alcoholics. *Int Pharmacopsychiatry*. 1981;16(1):49-56. PMID: 7028657.
15. Snyder S, Karacan I, Salis PJ. Effects of disulfiram on the sleep of chronic alcoholics. *Curr Alcohol*. 1981;8:159-66. PMID: 6282542.
16. Voronin K, Randall P, Myrick H, et al. Aripiprazole effects on alcohol consumption and subjective reports in a clinical laboratory paradigm--possible influence of self-control. *Alcohol Clin Exp Res*. 2008 Nov;32(11):1954-61. PMID: 18782344.

Wrong study design

1. Addolorato G, Leggio L, Ferrulli A, et al. Dose-response effect of baclofen in reducing daily alcohol intake in alcohol dependence: secondary analysis of a randomized, double-blind, placebo-controlled trial. *Alcohol Alcohol*. 2011 May-Jun;46(3):312-7. PMID: 21414953.
2. Adler L, Wilens T, Zhang S, et al. Retrospective safety analysis of atomoxetine in adult ADHD patients with or without comorbid alcohol abuse and dependence. *Am J Addict*. 2009 Sep-Oct;18(5):393-401. PMID: 19874159.
3. Angelone SM, Bellini L, Di Bella D, et al. Effects of fluvoxamine and citalopram in maintaining abstinence in a sample of Italian detoxified alcoholics. *Alcohol Alcohol*. 1998 Mar-Apr;33(2):151-6. PMID: 9566477.
4. Auriacombe M, Robinson M, Grabot D, et al. Naltrexone is ineffective to prevent relapse to alcohol in a realistic outpatient setting. A double blind one-year controlled study [abstract]. *College of Problems of Drug Dependence*; 2000 San Juan, Puerto Rico.
5. Barber TJ, Marett B, Waldron S, et al. Are disulfiram-like reactions associated with abacavir-containing antiretroviral regimens in clinical practice? *AIDS*. 2007 Aug 20;21(13):1823-4. PMID: 17690585.
6. Baros AM, Latham PK, Anton RF. Naltrexone and cognitive behavioral therapy for the treatment of alcohol dependence: do sex differences exist? *Alcohol Clin Exp Res*. 2008 May;32(5):771-6. PMID: 18336635.
7. Basu D, Jhirwal OP, Mattoo SK. Clinical characterization of use of acamprosate and naltrexone: data from an addiction center in India. *Am J Addict*. 2005 Jul-Sep;14(4):381-95. PMID: 16188718.
8. Blomqvist O, Hernandez-Avila CA, Burleson JA, et al. Self-Efficacy as a Predictor of Relapse during Treatment for Alcohol Dependence. *Addict Disord Their Treat*. 2003;2(4):135-45.
9. Brunette MF, O'Keefe C, Zimmet S, et al. Clozapine, olanzapine, or typical antipsychotics for alcohol use disorder in patients with schizophrenia. *J Dual Diagn*. 2008;4(4):344-54.
10. Buri C, Moggi F, Giovanoli A, et al. Prescription procedures in medication for relapse prevention after inpatient treatment for alcohol use disorders in Switzerland. *Alcohol Alcohol*. 2007 Jul-Aug;42(4):333-9. PMID: 17517820.
11. Burman WJ, Terra M, Breese P, et al. Lack of toxicity from concomitant directly observed disulfiram and isoniazid-containing therapy for active tuberculosis. *Int J Tuberc Lung Dis*. 2002;6(9):839-42.
12. Caputo F, Addolorato G, Lorenzini F, et al. Gamma-hydroxybutyric acid versus naltrexone in maintaining alcohol abstinence: an open randomized comparative study. *Drug Alcohol Depend*. 2003 May 1;70(1):85-91. PMID: 12681528.
13. Chick J, Gough K, Falkowski W, et al. Disulfiram treatment of alcoholism. *Br J Psychiatry*. 1992 Jul;161:84-9. PMID: 1638335.
14. Cornelius JR, Salloum IM, Ehler JG, et al. Double-blind fluoxetine in depressed alcoholic smokers. *Psychopharmacol Bull*. 1997;33(1):165-70. PMID: 9133770.
15. Cornelius JR, Salloum IM, Thase ME, et al. Fluoxetine versus placebo in depressed alcoholic cocaine abusers. *Psychopharmacol Bull*. 1998;34(1):117-21. PMID: 9564208.
16. Cornelius JR, Perkins KA, Salloum IM, et al. Fluoxetine versus placebo to decrease the smoking of depressed alcoholic patients. *J Clin Psychopharmacol*. 1999;19(2):183-4. PMID: 10211921.
17. Cornelius JR, Salloum IM, Haskett RF, et al. Fluoxetine versus placebo in depressed alcoholics: a 1-year follow-up study. *Addict Behav*. 2000 Mar-Apr;25(2):307-10. PMID: 10795957.
18. Critchfield GC, Eddy DM. A confidence profile analysis of the effectiveness of disulfiram in the treatment of chronic alcoholism. *Med Care*. 1987 Dec;25(12 Suppl):S66-75. PMID: 3323686.

19. Davidson D, Wirtz PW, Gulliver SB, et al. Naltrexone's suppressant effects on drinking are limited to the first 3 months of treatment. *Psychopharmacology (Berl)*. 2007 Sep;194(1):1-10. PMID: 17514344.
20. Diehl A, Ulmer L, Mutschler J, et al. Why is disulfiram superior to acamprosate in the routine clinical setting? A retrospective long-term study in 353 alcohol-dependent patients. *Alcohol Alcohol*. 2010 May-Jun;45(3):271-7. PMID: 20348436.
21. Duckert F, Johnsen J. Behavioral use of disulfiram in the treatment of problem drinking. *Int J Addict*. 1987 May;22(5):445-54. PMID: 3596857.
22. Dundon W, Lynch KG, Pettinati HM, et al. Treatment outcomes in type A and B alcohol dependence 6 months after serotonergic pharmacotherapy. *Alcohol Clin Exp Res*. 2004 Jul;28(7):1065-73. PMID: 15252293.
23. Farren CK, O'Malley SS. Occurrence and management of depression in the context of naltrexone treatment of alcoholism. *Am J Psychiatry*. 1999 Aug;156(8):1258-62. PMID: 10450269.
24. Feeney GF, Connor JP, Young RM, et al. Combined acamprosate and naltrexone, with cognitive behavioural therapy is superior to either medication alone for alcohol abstinence: a single centres' experience with pharmacotherapy. *Alcohol Alcohol*. 2006 May-Jun;41(3):321-7. PMID: 16467406.
25. Finigan MW, Perkins T, Zold-Kilbourn P, et al. Preliminary evaluation of extended-release naltrexone in Michigan and Missouri drug courts. *J Subst Abuse Treat*. 2011 Oct;41(3):288-93. PMID: 21696912.
26. Freeman SA. Focus on time-sensitive factors that may respond to treatment. *Curr Psychiatry*. 2012;11(1):53-7.
27. Gamburg AL, Aranovich AG, Rasnyuk VA, et al. Some aspects of contemporary treatment of chronic alcoholics. *Soviet Neurology & Psychiatry*. 1984;17(2):50-7.
28. Gerra G, Caccavari R, Delsignore R, et al. Effects of fluoxetine and Ca-acetyl-homotaurinate on alcohol intake in familial and nonfamilial alcoholic patients. *Curr Ther Res Clin Exp*. 1992;52(2):291-5.
29. Graf M. Pharmacological treatments at the East Kent Alcohol Service. *Neuropsychopharmacol Hung*. 2012;14:38.
30. Gueorguieva R, Wu R, Pittman B, et al. New insights into the efficacy of naltrexone based on trajectory-based reanalyses of two negative clinical trials. *Biol Psychiatry*. 2007 Jun 1;61(11):1290-5. PMID: 17224132.
31. Higgins ST, Budney AJ, Bickel WK, et al. Disulfiram therapy in patients abusing cocaine and alcohol. *Am J Psychiatry*. 1993;150(4):675-6. PMID: 8465895.
32. Hobbs JD, Kushner MG, Lee SS, et al. Meta-analysis of supplemental treatment for depressive and anxiety disorders in patients being treated for alcohol dependence (Structured abstract). *Am J Addict*; 2011. p. 319-29.
33. Ionescu R. Lithium salts in alcohol addiction therapy. *Revue Roumaine de Neurologie et Psychiatrie*. 1985;23(1):3-10.
34. Johnson BA, O'Malley SS, Ciraulo DA, et al. Dose-ranging kinetics and behavioral pharmacology of naltrexone and acamprosate, both alone and combined, in alcohol-dependent subjects. *J Clin Psychopharmacol*. 2003 Jun;23(3):281-93. PMID: 12826990.
35. Karam-Hage M, Brower KJ. Open pilot study of gabapentin versus trazodone to treat insomnia in alcoholic outpatients. *Psychiatry Clin Neurosci*. 2003;57(5):542-4.
36. Kiec-Swierczynska M, Krecisz B, Fabicka B. Systemic contact dermatitis from implanted disulfiram. *Contact Dermatitis*. 2000 Oct;43(4):246-7. PMID: 11011945.
37. Kiefer F, Jimenez-Arriero MA, Klein O, et al. Cloninger's typology and treatment outcome in alcohol-dependent subjects during pharmacotherapy with naltrexone. *Addict Biol*. 2008 Mar;13(1):124-9. PMID: 17573782.
38. Koeter MW, van den Brink W, Lehter P. Effect of early and late compliance on the effectiveness of acamprosate in the treatment of alcohol dependence. *J Subst Abuse Treat*. 2010 Oct;39(3):218-26. PMID: 20627222.

39. Koski A, Ojanpera I, Vuori E. Interaction of alcohol and drugs in fatal poisonings. *Hum Exp Toxicol*. 2003 May;22(5):281-7. PMID: 12774892.
40. Krampe H, Stawicki S, Wagner T, et al. Follow-up of 180 alcoholic patients for up to 7 years after outpatient treatment: impact of alcohol deterrents on outcome. *Alcohol Clin Exp Res*. 2006 Jan;30(1):86-95. PMID: 16433735.
41. Kranzler HR, Del Boca F, Korner P, et al. Adverse effects limit the usefulness of fluvoxamine for the treatment of alcoholism. *J Subst Abuse Treat*. 1993 May-Jun;10(3):283-7. PMID: 8315702.
42. Kranzler HR, Van Kirk J. Efficacy of naltrexone and acamprosate for alcoholism treatment: A meta-analysis. *Alcohol*. 2001;25(9):1335-41.
43. Kranzler HR, Gage A. Acamprosate efficacy in alcohol-dependent patients: summary of results from three pivotal trials. *Am J Addict*. 2008 Jan-Feb;17(1):70-6. PMID: 18214726.
44. Krystal JH, Gueorguieva R, Cramer J, et al. Naltrexone is associated with reduced drinking by alcohol dependent patients receiving antidepressants for mood and anxiety symptoms: Results from VA cooperative study no. 425, "naltrexone in the treatment of alcoholism". *Alcohol*. 2008;32(1):85-91.
45. Leggio L, Ferrulli A, Zambon A, et al. Baclofen promotes alcohol abstinence in alcohol dependent cirrhotic patients with hepatitis C virus (HCV) infection. *Addict Behav*. 2012 Apr;37(4):561-4. PMID: 22244707.
46. Mann K, Kiefer F, Smolka M, et al. Searching for responders to acamprosate and naltrexone in alcoholism treatment: rationale and design of the PREDICT study. *Alcohol Clin Exp Res*. 2009 Apr;33(4):674-83. PMID: 19170666.
47. Mason BJ, Leher P. Effects of nicotine and illicit substance use on alcoholism treatment outcomes and acamprosate efficacy. *J Addict Med*. 2009;3(3):164-71. PMID: 21769013.
48. Minuk GY, Rockman GE, German GB, et al. The use of sodium valproate in the treatment of alcoholism. *J Addict Dis*. 1995;14(2):67-74. PMID: 8541361.
49. Monnelly EP, LoCastro JS, Gagnon D, et al. Quetiapine versus trazodone in reducing rehospitalization for alcohol dependence: A large data-base study. *J Addict Med*. 2008;2(3):128-34. PMID: 21768982.
50. Mueser KT, Noordsy DL, Fox L, et al. Disulfiram treatment for alcoholism in severe mental illness. *Am J Addict*. 2003 May-Jun;12(3):242-52. PMID: 12851020.
51. Naranjo CA, Knoke DM, Bremner KE. Variations in response to citalopram in men and women with alcohol dependence. *J Psychiatry Neurosci*. 2000 May;25(3):269-75. PMID: 10863887.
52. Neto D, Lambaz R, Tavares JE. Compliance with aftercare treatment, including disulfiram, and effect on outcome in alcohol-dependent patients. *Alcohol Alcohol*. 2007 Nov-Dec;42(6):604-9. PMID: 17878216.
53. Niederhofer H, Staffen W. Acamprosate and its efficacy in treating alcohol dependent adolescents. *Eur Child Adolesc Psychiatry*. 2003 Jun;12(3):144-8. PMID: 12768462.
54. Nunes EV, McGrath PJ, Quitkin FM, et al. Imipramine treatment of alcoholism with comorbid depression. *Am J Psychiatry*. 1993 Jun;150(6):963-5. PMID: 8494079.
55. O'Malley SS, Croop RS, Wroblewski JM, et al. Naltrexone in the treatment of alcohol dependence: A combined analysis of two trials. *Psychiatr Ann*. 1995;25(11):681-8.
56. O'Malley SS, Rounsaville BJ, Farren C, et al. Initial and maintenance naltrexone treatment for alcohol dependence using primary care vs specialty care: a nested sequence of 3 randomized trials. *Arch Intern Med*. 2003 Jul 28;163(14):1695-704. PMID: 12885685.
57. Oncken C, Van Kirk J, Kranzler HR. Adverse effects of oral naltrexone: analysis of data from two clinical trials. *Psychopharmacology (Berl)*. 2001 Apr;154(4):397-402. PMID: 11349393.

58. Oslin DW, Pettinati H, Volpicelli JR. Alcoholism treatment adherence: older age predicts better adherence and drinking outcomes. *Am J Geriatr Psychiatry*. 2002 Nov-Dec;10(6):740-7. PMID: 12427583.
59. Oslin D. OPRM1 sequence variation and clinical studies of alcohol dependent adults. *Alcohol*. 2011;35:314A.
60. Paparrigopoulos T, Tzavellas E, Karaikos D, et al. Treatment of alcohol dependence with low-dose topiramate: an open-label controlled study. *BMC Psychiatry*. 2011;11:41. PMID: 21401921.
61. Pettinati HM, Volpicelli JR, Kranzler HR, et al. Sertraline treatment for alcohol dependence: interactive effects of medication and alcoholic subtype. *Alcohol Clin Exp Res*. 2000 Jul;24(7):1041-9. PMID: 10924008.
62. Pettinati HM, Volpicelli JR, Pierce JD, Jr., et al. Improving naltrexone response: an intervention for medical practitioners to enhance medication compliance in alcohol dependent patients. *J Addict Dis*. 2000;19(1):71-83. PMID: 10772604.
63. Pettinati HM, Dundon W, Lipkin C. Gender differences in response to sertraline pharmacotherapy in Type A alcohol dependence. *Am J Addict*. 2004 May-Jun;13(3):236-47. PMID: 15370943.
64. Pettinati HM, Silverman BL, Battisti JJ, et al. Efficacy of extended-release naltrexone in patients with relatively higher severity of alcohol dependence. *Alcohol Clin Exp Res*. 2011 Oct;35(10):1804-11. PMID: 21575016.
65. Prisciandaro JJ, Desantis SM, Bandyopadhyay D. Simultaneous Modeling of the Impact of Treatments on Alcohol Consumption and Quality of Life in the COMBINE Study: A Coupled Hidden Markov Analysis. *Alcohol*. 2012;36(12):2141-9.
66. Roosa BA. Alcoholism treatment and medical utilization. US: ProQuest Information & Learning; 1986.
67. Rosenberg CM. Drug maintenance in the outpatient treatment of chronic alcoholism. *Arch Gen Psychiatry*. 1974 Mar;30(3):373-7. PMID: 4813140.
68. Rosenthal RN, Gage A, Perhach JL, et al. Acamprosate: Safety and tolerability in the treatment of alcohol dependence. *J Addict Med*. 2008;2(1):40-50. PMID: 21768971.
69. Rubio G, Manzanares J, Lopez-Munoz F, et al. Naltrexone improves outcome of a controlled drinking program. *J Subst Abuse Treat*. 2002 Dec;23(4):361-6. PMID: 12495798.
70. Schmidt LG, Smolka MN. Results from two pharmacotherapy trials show alcoholic smokers were more severely alcohol dependent but less prone to relapse than alcoholic non-smokers. *Alcohol Alcohol*. 2007 May-Jun;42(3):241-6. PMID: 17526634.
71. Sellers EM, Naranjo CA, Kadlec K. Do serotonin uptake inhibitors decrease smoking? Observations in a group of heavy drinkers. *J Clin Psychopharmacol*. 1987 Dec;7(6):417-20. PMID: 2963034.
72. Tollefson GD. Serotonin and alcohol: Interrelationships. *Psychopathology*. 1989;22(SUPPL. 1):37-48.
73. Tolliver BK, McRae AL, Sonne SC, et al. Safety and tolerability of acamprosate in alcohol-dependent individuals with bipolar disorder: An open-label pilot study. *Addict Disord Their Treat*. 2009;8(1):33-8.
74. Ulmer A, Müller M, Frietsch B. Dihydrocodeine/agonists for alcohol dependents. *Front Psychiatry*. 2012;3PMID: 22470353.
75. Vaz de Lima Fabiana B, Andriolo Régis B, da Silveira Dartiu X. Dopaminergic antagonists for alcohol dependence. *Cochrane Database Syst Rev*. 2010(4)PMID: CD008460.
76. Verheul R, Leher P, Geerlings PJ, et al. Predictors of acamprosate efficacy: results from a pooled analysis of seven European trials including 1485 alcohol-dependent patients. *Psychopharmacology (Berl)*. 2005 Mar;178(2-3):167-73. PMID: 15322728.
77. Volpicelli JR, Watson NT, King AC, et al. Effect of naltrexone on alcohol 'high' in alcoholics. *Am J Psychiatry*. 1995;152(4):613-5. PMID: 7694913.
78. Weinrieb RM, O'Brien CP. Naltrexone in the treatment of alcoholism. *Annu Rev Med*. 1997;48:477-87. PMID: 9046978.

79. Wilens TE, Adler LA, Weiss MD, et al. Atomoxetine treatment of adults with ADHD and comorbid alcohol abuse. Proceedings of the 69th Annual Scientific Meeting of the College on Problems of Drug Dependence; 2007 June 16-21; Quebec City, Canada; 2007.
80. Workeneh B, Balakumaran A, Bichet DG, et al. The dilemma of diagnosing the cause of hypernatraemia: Drinking habits vs diabetes insipidus. *Nephrol Dial Transplant*. 2004;19(12):3165-7.
81. Yen MH, Ko HC, Tang FI, et al. Study of hepatotoxicity of naltrexone in the treatment of alcoholism. *Alcohol*. 2006 Feb;38(2):117-20. PMID: 16839858.

Treatment duration less than 12 weeks

1. Addolorato G, Caputo F, Capristo E, et al. Baclofen efficacy in reducing alcohol craving and intake: a preliminary double-blind randomized controlled study. *Alcohol Alcohol*. 2002 Sep-Oct;37(5):504-8. PMID: 12217947.
2. Brower KJ, Myra Kim H, Strobbe S, et al. A randomized double-blind pilot trial of gabapentin versus placebo to treat alcohol dependence and comorbid insomnia. *Alcohol Clin Exp Res*. 2008 Aug;32(8):1429-38. PMID: 18540923.
3. Bruno F. Buspirone in the treatment of alcoholic patients. *Psychopathology*. 1989;22 Suppl 1:49-59. PMID: 2657838.
4. Christensen JK, Ronsted P, Vaag UH. Side effects after disulfiram. Comparison of disulfiram and placebo in a double-blind multicentre study. *Acta Psychiatr Scand*. 1984 Apr;69(4):265-73. PMID: 6372372.
5. Ciraulo DA, Dong Q, Silverman BL, et al. Early treatment response in alcohol dependence with extended-release naltrexone. *J Clin Psychiatry*. 2008 Feb;69(2):190-5. PMID: 18348601.
6. Eriksson M, Berggren U, Blennow K, et al. Further investigation of citalopram on alcohol consumption in heavy drinkers: responsiveness possibly linked to the DRD2 A2/A2 genotype. *Alcohol*. 2001 May;24(1):15-23. PMID: 11524178.
7. Furieri FA, Nakamura-Palacios EM. Gabapentin reduces alcohol consumption and craving: a randomized, double-blind, placebo-controlled trial. *J Clin Psychiatry*. 2007 Nov;68(11):1691-700. PMID: 18052562.
8. Galarza NJ, Diaz Ramirez D, Guzman F, et al. The use of naltrexone to treat ambulatory patients with alcohol dependence. *Bol Asoc Med P R*. 1997 Oct-Dec;89(10-12):157-60. PMID: 9577049.
9. Grassi MC, Cioce AM, Giudici FD, et al. Short-term efficacy of Disulfiram or Naltrexone in reducing positive urinalysis for both cocaine and cocaethylene in cocaine abusers: A pilot study. *Pharmacol Res*. 2007;55(2):117-21.
10. Hammarberg A, Nylander I, Zhou Q, et al. The effect of acamprosate on alcohol craving and correlation with hypothalamic pituitary adrenal (HPA) axis hormones and beta-endorphin. *Brain Res*. 2009 Dec 11;1305 Suppl:S2-6. PMID: 19799882.
11. Hammarberg A, Jayaram-Lindstrom N, Beck O, et al. The effects of acamprosate on alcohol-cue reactivity and alcohol priming in dependent patients: a randomized controlled trial. *Psychopharmacology (Berl)*. 2009 Jul;205(1):53-62. PMID: 19319508.
12. Hernandez-Avila CA, Song C, Kuo L, et al. Targeted versus daily naltrexone: secondary analysis of effects on average daily drinking. *Alcohol Clin Exp Res*. 2006 May;30(5):860-5. PMID: 16634855.
13. Hersh D, Van Kirk JR, Kranzler HR. Naltrexone treatment of comorbid alcohol and cocaine use disorders. *Psychopharmacology (Berl)*. 1998 Sep;139(1-2):44-52. PMID: 9768541.
14. Janiri L, Gobbi G, Mannelli P, et al. Effects of fluoxetine and antidepressant doses on short-term outcome of detoxified alcoholics. *Int Clin Psychopharmacol*. 1996;11(2):109-17. PMID: 8803648.

15. Janiri L, Gobbi G, Mannelli P, et al. Effects of fluoxetine at antidepressant doses on short-term outcome of detoxified alcoholics. *Int Clin Psychopharmacol*. 1996 Jun;11(2):109-17. PMID: 8803648.
16. Johnson BA, Roache JD, Javors MA, et al. Ondansetron for reduction of drinking among biologically predisposed alcoholic patients: A randomized controlled trial. *JAMA*. 2000 Aug 23-30;284(8):963-71. PMID: 10944641.
17. Johnson BA, Ait-Daoud N, Seneviratne C, et al. Pharmacogenetic approach at the serotonin transporter gene as a method of reducing the severity of alcohol drinking. *Am J Psychiatry*. 2011 Mar;168(3):265-75. PMID: 21247998.
18. Kenna GA, Zywiak WH, McGeary JE, et al. A within-group design of nontreatment seeking 5-HTTLPR genotyped alcohol-dependent subjects receiving ondansetron and sertraline. *Alcohol Clin Exp Res*. 2009 Feb;33(2):315-23. PMID: 19032576.
19. Kranzler HR, Modesto-Lowe V, Nuwayser ES. Sustained-release naltrexone for alcoholism treatment: a preliminary study. *Alcohol Clin Exp Res*. 1998 Aug;22(5):1074-9. PMID: 9726277.
20. Kranzler HR, Modesto-Lowe V, Van Kirk J. Naltrexone vs. nefazodone for treatment of alcohol dependence. A placebo-controlled trial. *Neuropsychopharmacology*. 2000 May;22(5):493-503. PMID: 10731624.
21. Kranzler HR, Armeli S, Tennen H, et al. Targeted naltrexone for early problem drinkers. *J Clin Psychopharmacol*. 2003 Jun;23(3):294-304. PMID: 12826991.
22. Kranzler HR, Armeli S, Feinn R, et al. Targeted naltrexone treatment moderates the relations between mood and drinking behavior among problem drinkers. *J Consult Clin Psychol*. 2004 Apr;72(2):317-27. PMID: 15065964.
23. Kranzler HR, Mueller T, Cornelius J, et al. Sertraline treatment of co-occurring alcohol dependence and major depression. *J Clin Psychopharmacol*. 2006 Feb;26(1):13-20. PMID: 16415699.
24. Litten RZ, Fertig JB, Falk DE, et al. A double-blind, placebo-controlled trial to assess the efficacy of quetiapine fumarate XR in very heavy-drinking alcohol-dependent patients. *Alcohol Clin Exp Res*. 2012 Mar;36(3):406-16. PMID: 21950727.
25. Loo H, Malka R, Defrance R, et al. Tianeptine and amitriptyline. Controlled double-blind trial in depressed alcoholic patients. *Neuropsychobiology*. 1988;19(2):79-85. PMID: 3067116.
26. Myrick H, Li X, Randall PK, et al. The effect of aripiprazole on cue-induced brain activation and drinking parameters in alcoholics. *J Clin Psychopharmacol*. 2010 Aug;30(4):365-72. PMID: 20571434.
27. Namkoong K, Lee BO, Lee PG, et al. Acamprosate in Korean alcohol-dependent patients: a multi-centre, randomized, double-blind, placebo-controlled study. *Alcohol Alcohol*. 2003 Mar-Apr;38(2):135-41. PMID: 12634260.
28. Naranjo CA, Kadlec KE, Sanhueza P, et al. Fluoxetine differentially alters alcohol intake and other consummatory behaviors in problem drinkers. *Clin Pharmacol Ther*. 1990 Apr;47(4):490-8. PMID: 2328557.
29. Pettinati HM, Kampman KM, Lynch KG, et al. A double blind, placebo-controlled trial that combines disulfiram and naltrexone for treating co-occurring cocaine and alcohol dependence. *Addict Behav*. 2008 May;33(5):651-67. PMID: 18079068.
30. Prisciandaro JJ, Brown DG, Brady KT, et al. Comorbid anxiety disorders and baseline medication regimens predict clinical outcomes in individuals with co-occurring bipolar disorder and alcohol dependence: Results of a randomized controlled trial. *Psychiatry Res*. 2011 Aug 15;188(3):361-5. PMID: 21641663.
31. Randall CL, Johnson MR, Thevos AK, et al. Paroxetine for social anxiety and alcohol use in dual-diagnosed patients. *Depress Anxiety*. 2001;14(4):255-62. PMID: 11754136.
32. Ray LA, Miranda R, Jr., MacKillop J, et al. A preliminary pharmacogenetic investigation of adverse events from topiramate in heavy drinkers. *Exp Clin Psychopharmacol*. 2009 Apr;17(2):122-9. PMID: 19331489.

33. Roache JD. L/S serotonin transporter polymorphism predicts serotonergic treatment outcome in alcohol dependence. *Alcohol*. 2011;35:339A.
34. Sellers EM, Higgins GA, Tompkins DM, et al. Serotonin and alcohol drinking. *NIDA Res Monogr*. 1992;119:141-5. PMID: 1435969.
35. Sellers EM, Toneatto T, Romach MK, et al. Clinical efficacy of the 5-HT₃ antagonist ondansetron in alcohol abuse and dependence. *Alcohol Clin Exp Res*. 1994 Aug;18(4):879-85. PMID: 7978099.
36. Simpson TL, Saxon AJ, Meredith CW, et al. A pilot trial of the alpha-1 adrenergic antagonist, prazosin, for alcohol dependence. *Alcohol Clin Exp Res*. 2009 Feb;33(2):255-63. PMID: 18945226.
37. Sloan TB, Roache JD, Johnson BA. The role of anxiety in predicting drinking behaviour. *Alcohol Alcohol*. 2003 Jul-Aug;38(4):360-3. PMID: 12814905.
38. Su N, Zhang L, Fei F, et al. The brain-derived neurotrophic factor is associated with alcohol dependence-related depression and antidepressant response. *Brain Res*. 2011 Sep 30;1415:119-26. PMID: 21880305.
39. Tidey JW, Monti PM, Rohsenow DJ, et al. Moderators of naltrexone's effects on drinking, urge, and alcohol effects in non-treatment-seeking heavy drinkers in the natural environment. *Alcohol Clin Exp Res*. 2008 Jan;32(1):58-66. PMID: 18028530.
40. Tolliver BK, Brady KT. Glutamate neurotransmission as a potential therapeutic target in bipolar disorder: An overview of evidence and implications for treatment of co-occurring disorders. *Bipolar Disord*. 2011;13:100.
41. Tolliver BK, Desantis SM, Brown DG, et al. A randomized, double-blind, placebo-controlled clinical trial of acamprosate in alcohol-dependent individuals with bipolar disorder: a preliminary report. *Bipolar Disord*. 2012 Feb;14(1):54-63. PMID: 22329472.
42. Toneatto T, Brands B, Selby P. A randomized, double-blind, placebo-controlled trial of naltrexone in the treatment of concurrent alcohol use disorder and pathological gambling. *Am J Addict*. 2009 May-Jun;18(3):219-25. PMID: 19340640.
43. Trevisan LA, Ralevski E, Keegan K, et al. Alcohol detoxification and relapse prevention using valproic acid versus gabapentin in alcohol-dependent patients. *Addict Disord Their Treat*. 2008;7(3):119-28.

Systematic review with search that ended prior to 2007

1. Edwards JD, Jr., Dill JE. Alcoholism clinic in a military setting: a combined disulfiram and group therapy outpatient program. *Mil Med*. 1974 Mar;139(3):206-9. PMID: 4206169.
2. Garbutt JC, West SL, Carey TS, et al. Pharmacological treatment of alcohol dependence: a review of the evidence. *JAMA*. 1999 Apr 14;281(14):1318-25. PMID: 10208148.
3. Miller WR, Wilbourne PL. Mesa Grande: A methodological analysis of clinical trials of treatment for alcohol use disorders. *Addiction*. 2002;97(3):265-77. PMID: 11964100.
4. Monahan SC, Finney JW. Explaining abstinence rates following treatment for alcohol abuse: a quantitative synthesis of patient, research design and treatment effects (Structured abstract). *Addiction*. 1996. p. 787-805.
5. Pettinati HM, O'Brien CP, Rabinowitz AR, et al. The status of naltrexone in the treatment of alcohol dependence: specific effects on heavy drinking. *J Clin Psychopharmacol*. 2006 Dec;26(6):610-25. PMID: 17110818.
6. Rosner S, Leucht S, Leher P, et al. Acamprosate supports abstinence, naltrexone prevents excessive drinking: evidence from a meta-analysis with unreported outcomes. *J Psychopharmacol*. 2008 Jan;22(1):11-23. PMID: 18187529.

7. Snyder JL, Bowers TG. The efficacy of acamprosate and naltrexone in the treatment of alcohol dependence: a relative benefits analysis of randomized controlled trials. *Am J Drug Alcohol Abuse*. 2008;34(4):449-61. PMID: 18584575.
8. Srisurapanont M, Jarusuraisin N. Naltrexone for the treatment of alcoholism: a meta-analysis of randomized controlled trials. *Int J Neuropsychopharmacol*. 2005 Jun;8(2):267-80. PMID: 15850502.
9. Streeton C, Whelan G. Naltrexone, a relapse prevention maintenance treatment of alcohol dependence: a meta-analysis of randomized controlled trials. *Alcohol Alcohol*. 2001 Nov-Dec;36(6):544-52. PMID: 11704620.

Appendix C. Criteria Used for Evaluating Studies' Risk of Bias

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Petrakis, 2012 ¹ NA DBRCT	NR/CND	NR/CND	Yes	44.3	24 ^a	Yes	No	Yes
Kranzler, 2012 ² NA SSGA	Yes	NR/CND	Yes	38	12 (14 at 6 month follow up)	Yes	No	Yes
Fogaca, 2011 ³ NA DBRCT	NR/CND	NR/CND	NR/CND	46	15% (between PUFAs group and NTX+PUFAs); 0% (between NTX and placebo groups as both were 45% attrition)	Yes	No	NR/CND
Ralevski, 2011 ⁴ ; Ralevski, 2001 ⁵ NA DBRCT	NR/CND	NR/CND	Yes, except all 4 women were randomized to the placebo group	35	NR/CND	Yes	No	NR/CND
Wolwer, 2011 ⁶ NA DBRCT	NR/CND	Yes	Yes, except for fewer women in IBT+placebo group	~20 lost to follow-up; 55% did not complete (most due to relapse)	4	No	No	NR/CND
Anton, 2011 ⁷ NA DBRCT	NR/CND	NR/CND	Yes	3% had no drinking data; 35% did not complete treatment; 12 to 18% provided drinking data for all 16 weeks	1% for no drinking data; 10% for not completing treatment; 6% for providing drinking data for all 16 weeks	No	No	NR/CND
Kranzler, 2011 ⁸ DBRCT	Yes	NR/CND	Yes	38% did not complete	12	Yes	No	Yes
Florez, 2011 ⁹ NA OLRCT	NR/CND	NR/CND	Yes	9	5	No	No	Yes

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Garbutt, 2010 ¹⁰ NA DBRCT	Yes	NR/CND	Yes	24	8	No	No	NR/CND
Stedman, 2010 ¹¹ NA DBRCT	NR/CND	NR/CND	Yes	57	1	Yes	NR/CND	NR/CND
Kiefer, 2011 ¹² NA SSGA	Yes (for the PREDICT study)	NR/CND	NR/CND	NR/CND	NR/CND	NR/CND	NR/CND	NR/CND
Pettinati, 2010 ¹³ NA DBRCT	Yes	NR/CND	Yes	43 (did not complete study, but just 3/170 subjects had no data for drinking outcomes)	6.5	Yes	No	Yes
Rubio, 2009 ¹⁴ NA DBRCT	NR/CND	NR/CND	Yes	17	2	No	No	NR/CND
Schmitz, 2009 ¹⁵ NA DBRCT	Yes	Yes	No	76% completed 12 weeks; 60% completed 6 weeks; lost to follow- up/missing data NR	NR (but median survival times before dropout were similar)	Yes	NR/CND	NR/CND
Brown, 2009 ¹⁶ NA DBRCT	NR/CND	NR/CND	Mixed	48	17	Yes	Yes	NR/CND
Longabaugh, 2009 ¹⁷ NA DBRCT	Yes	NR/CND	No	18	NR/CND for the 4 groups; 0% for those receiving BST vs. MET	No	NR/CND	NR/CND
Kranzler, 2009 ¹⁸ NA DBRCT	NR/CND	NR/CND	NR/CND	15	NR/CND	No	NR/CND	Yes
Baltieri, 2008 ¹⁹ , Yes Baltieri, 2009 ²⁰ NA DBRCT	Yes	Yes	Yes	45	4.3, 16.6, and 20.9 differences between each pair of groups	Yes	Yes	NR/CND

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Florez, 2008 ²¹ NA SSGA	NA (NR/CND for the parent study)	NR/CND	NR/CND	NR/CND	NR/CND	NR/CND	NR/CND	NR/CND
Oslin, 2008 ²² NA DBRCT	NR/CND	NR/CND	No	23	5 (for all NTX vs. all placebo)	No	No	Yes
Arias, 2008 ²³ NA SSGA	NA (yes for the parent study)	Yes for the parent study	No	33	NR/CND	Yes	NR/CND	NR/CND
Martinotti, 2009 ²⁴ NA DBRCT	Yes	Yes	NR/CND	25	1	No	No	NR/CND
Florez, 2008 ²⁵ NA OLRCT	NR/CND	NR/CND	No	10	4	No	No	Yes
O'Malley, 2008 ²⁶ NA DBRCT	NR/CND	NR/CND	Yes	33% did not complete; 15 25% unable to contact or declined further contact or moved		Yes	No	Yes
Wilens, 2008 ²⁷ NA DBRCT	NRCND	NR/CND	Yes	54	20	Yes	No	Yes
Brown, 2008 ²⁸ NA DBRCT	NR/CND	NR/CND	Yes, for most characteristics; No, for race/ethnicity, and concomitant medications	NR/CND	NR/CND	NR/CND	Yes	NR/CND

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Anton, 2008 ²⁹ COMBINE SSGA	NA (Yes for parent trial)	Yes for parent trial	Yes	Overall NR/CND 14% of white subjects in the parent trial who received NTX or placebo were not included in this study (604/706 were included); 56% of subjects randomized in COMBINE were not included in this study (604/1383 were included); Overall attrition in COMBINE was 6%	NR/CND (but was very low in overall COMBINE, and unlikely to be much different)	No	No	Yes
Lucey, 2008 ³⁰ NA DBRCT	Yes	NR/CND	Yes	25 (loss to follow up and withdrawal of consent)	7	No	No	Yes
Anton, 2008 ³¹ NA DBRCT	NR/CND	NR/CND	Yes for most characteristics; more males in aripiprazole group (75% vs. 62%)	<i>Aripiprazole</i> vs. <i>Placebo</i> Loss to follow-up: 7.4% vs. 9.6% Did not complete treatment phase: 41% vs. 26.7%	Loss to follow up: 2.2% Did not complete: 14%	Yes	No	Yes
Addolorato, 2007 ³² NA DBRCT	Yes	Yes	Yes	Loss to follow-up: 14% Total dropouts: 23%	Loss to follow-up: 9% Total dropouts: 17%	Yes, differential	No	NR/CND
Laaksonen, 2008 ³³ NA OLRCT	Yes	NR/CND	Yes for most variables; no for smoking	25% at 12 weeks (continuous med phase); 52% at 52 weeks (after targeted med phase)	7% at 12 weeks; 5% at 52 weeks	No at 12 weeks; Yes at 52 weeks	Np	Yes

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Johnson, 2007 ³⁴ Johnson, 2008 ³⁵ NA DBRCT	Yes	Yes	Yes	Loss to follow-up: 6% Non-completers: 31%	Loss to follow-up: 4% Non-completers: 15%	Yes	No	Yes
Pettinati, 2008 ³⁶ NA DBRCT	NR/CND	NR/CND	Yes	36% did not complete	10	Yes	No	Yes
Kampman, 2007 ³⁷ NA DBRCT	NR/CND	NR/CND	No	23	2	No	No	Yes (considering that dose reductions were allowed)
De Sousa, 2008 ³⁸ NA OLRCT	NR/CND	NR/CND	Yes	8	0	No	No	Yes
Karhuvaara, 2007 ³⁹ NA DBRCT	Yes	Yes	Yes	37 noncompleters; 9% lost to follow-up	8; 1	Yes	CND	Yes
Book, 2008 ⁴⁰ , Thomas, 2008 ⁴¹ NA DBRCT	Yes	Yes	Yes	About 37% (from Figure) did not provide data at weeks 12 and 16; % lost to followup/missing data NR	CND (appears <2% from Figure)	Yes	No	Yes
O'Malley, 2007 ⁴² NA DBRCT	NR/CND	NR/CND	Yes	23	1.2	No	No	NR/CND
Gelernter, 2007 ⁴³ VACS 425 SSGA	NA	NA	NR/CND	65 (just 220/627 subjects in the main trial were included in this sample)	NR/CND	Yes	No	Yes
Nava, 2006 ⁴⁴ NA OLRCT	Yes	NR/CND	Yes	31	17	Yes	NR/CND	CND

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Morley, 2006 ⁴⁵ Morley, 2010 ⁴⁶ NA DBRCT	Yes	NR/CND	Yes	Loss to follow-up or unwilling to continue: 12% Non-completers: 31%	Loss to follow-up or unwilling to continue: 5% Non-completers: 9%	No	NR/CND	Yes
Anton, 2006 ⁴⁷ Donovan, 2008 ⁴⁸ LoCastro, 2009 ⁴⁹ Greenfield, 2010 ⁵⁰ Fucito, 2012 ⁵¹ COMBINE DBRCT	Yes	Yes	Yes	6 (16 wks) 18 (1 year)	7 (1 year)	No	No	Yes
Mason, 2006 ⁵² NA DBRCT	Yes	Yes	Yes	Loss to follow-up: 13% Non-completers: 51%	Loss to follow-up: 6% Non-completers: 14%	No	No	Yes
Hutchison, 2006 ⁵³ NA SBRCT	NR/CND	NR/CND	No	20	5	No	NR/CND	Yes
Huang, 2005 ⁵⁴ NA DBRCT	NR/CND	NR/CND	Yes	No data for primary outcome: 20 Non-completers: 40	No data for primary outcome: 10% Non-completers: 10%	No	NR/CND	NR/CND
De Sousa, 2005 ⁵⁵ NA OLRCT	NR/CND	No	Yes	7	2	No	No	Yes
Anton, 2005 ⁵⁶ NA DBRCT	NR/CND	NR/CND	Yes	19% did not complete trial; 15% did not have complete 12 week drinking data	9 to 11	No	No	Yes

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Petrakis, 2005 ⁵⁷ Ralevski, 2007 ⁵⁸ Petrakis, 2007 ⁵⁹ Petrakis, 2006 ⁶⁰ VA MIRECC DBRCT	NR/CND	NR/CND	Yes for most characteristic; No outcome data for number of other psych meds	11% without 12-week outcome data	2 to 7	No	Yes: some concern for contamination from additional psychiatric medications	NR/CND
Garbutt, 2005 ⁶¹ NA DBRCT	Yes	Yes	Yes	39% did not complete; 13% lost to follow-up	1%; 3%	Yes	NR/CND	NR/CND
Brady, 2005 ⁶² NA DBRCT	Yes	NR/CND	Yes	34 % (from consent to randomization); NR/CND for loss to follow-up	6	NR/CND	NR/CND	NR/CND
Salloum, 2005 ⁶³ NA DBRCT	Yes	NR/CND	Yes	62% non-completers; on average, 86% underwent assessment at each point; 100% underwent assessment at week 24	12; CND	Yes, but not high concern	No	NR/CND
Killeen, 2004 ⁶⁴ NA DBRCT	Yes	NR/CND	No	28	9	No	NR/CND	NR/CND
De Sousa, 2004 ⁶⁵ NA OLRCT	Yes	No	Yes	3	2	No	NR/CND	NR/CND
Schmitz, 2004 ⁶⁶ NA DBRCT	NR/CND	NR/CND	No	69% did not complete 12 weeks of treatment; lost to follow- up/missing data NR; mean sessions attended: 10.3	NR/CND	Yes	NR/CND	NR/CND

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Johnson, 2004 ⁶⁷ NA DBRCT	NR/CND	NR/CND	No	30	12	Yes	NR/CND	NR/CND
Johnson, 2004 ⁶⁸ NA DBRCT	NR/CND	NR/CND	NR/CND	35	11	Yes	NR/CND	NR/CND
Kranzler, 2004 ⁶⁹ NA DBRCT	NR/CND	Yes	Yes	22	5	No	NR/CND	Yes
Anton, 2004 ⁷⁰ NA DBRCT	NR/CND	NR/CND	Yes	26 did not complete 12 weeks; smaller number for lost-to follow up (6 to 16%) and missing data	1 to 9	No	NR/CND	NR/CND
Guardia, 2004 ⁷¹ NA DBRCT	Yes	NR/CND	Yes	32% non-completers; % missing data NR	19	Yes	NR/CND	NR/CND
Chick, 2004 ⁷² NA DBRCT	Yes	Yes	Yes	64% non-completers; 5.6% post-randomization exclusions (nont in ITT sample); 21% of the ITT sample lost to follow-up	17; 1; 1	Yes	No	NR/CND
Baltieri, 2004 ⁷³ NA DBRCT	NR/CND	NR/CND	Yes	23	5	No	NR/CND	NR/CND
Petrakis, 2004 ⁷⁴ ; Ralevski, 2006 ⁷⁵ NA DBRCT	NR/CND	NR/CND	Yes	19	12	No	No	NR/CND
Gual, 2003 ⁷⁶ NA DBRCT	NR/CND	NR/CND	Yes	45% did not complete; 13% lost to follow-up	2	Yes	No	NR/CND

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Moak, 2003 ⁷⁷ NA DBRCT	Yes	NR/CND	Yes	28% did not complete; 18 missing data NR		No	No	Yes
Balldin, 2003 ⁷⁸ NA DBRCT	Yes	Yes	Yes	22% terminated the study early; 9% had missing drinking data	NR/CND	No	No	Yes
Johnson, 2003 ⁷⁹ Ma, 2006 ⁸⁰ , Johnson, 2004 ⁶⁸ NA DBRCT	Yes	NR/CND	Yes	35% did not complete; 9; 2; unclear for 5% not assessed for outcomes at all; unclear amount of missing data	missing data	CND	No	NR/CND
Kiefer, 2003 ⁸¹ Kiefer, 2005 ⁸² NA DBRCT	Yes	Yes	Yes, for most characteristics; Drug arms had slightly more severe problems on some alcohol measures	0 lost to follow-up; 11% dropout; 53% did not complete trial (most because of relapse)	0 for lost to follow-up; No 40% for completion of trial (because 75% of the placebo group relapsed and did not complete)		No	NR/CND
Gastpar, 2002 ⁸³ NA DBRCT	NR/CND	NR/CND	Yes	36% did not complete (19% failed to return/lost to follow-up, 8% withdrew consent, 4% AEs, 1% protocol violations, 4% other reasons)	5	Yes	No	NR/CND
Guardia, 2002 ⁸⁴ NA DBRCT	NR/CND	NR/CND	Yes	5% did not have assessable data; 26% dropout, treatment refusal, or other reasons for not completing; 41% total did not complete the study for any reason	0%; 7%; 2	Yes	Possible contamination due to allowed SSRIs	NR/CND
Brady, 2002 ⁸⁵ NA DBRCT	NR/CND	NR/CND	Yes	26% non-completers; 6.5% not included in analyses	NR/CND	No	NR/CND	Yes
Latt, 2002 ⁸⁶ NA DBRCT	Yes	NR/CND	Yes	31% lost to follow-up; 0% excluded from analyses	3%; 0%	Yes	No	NR/CND

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Morris, 2001 ⁸⁷ NA DBRCT	NR/CND	NR/CND	No	36% did not complete; 20% dropout for reasons other than relapse	10%; 3%	No	No	NR/CND
Krystal, 2001 ⁸⁸ VACS 425 DBRCT	NR/CND	NR/CND	Yes	27% did not complete; 22% did not have complete data and 10% did not have complete or partially complete data for drinking at week 13	NR/CND; 2%; 1%	No	No	NR/CND
Monti, 2001 ⁸⁹ ; Rohsenow, 2007 ⁹⁰ ; Rohsenow, 2000 ⁹¹ NA DBRCT	NR/ CND	NR/CND	NR/CND	9 to 13	NR/CND	No	No	Yes
Monterosso, 2001 ⁹² NA DBRCT	NR/CND	NR/CND	NR/CND	17	NR/CND	No	No	Yes
Rubio, 2001 ⁹³ NA SBRCT	Yes	NA (open-label trial)	Yes	17	13	No	Yes	Yes
Heinala, 2001 ⁹⁴ NA DBRCT	NR/CND	NR/CND	NR/CND	32% did not complete study	NR/CND	Yes	NR/CND	NR/CND
Pettinati, 2001 ⁹⁵ NA DBRCT	NR/CND	NR/CND	Yes	42% did not complete the study; NR/CND for loss to follow-up; unclear how much missing data for alcohol outcomes among those	12%; NR/CND	Yes	No	NR/CND
Chick, 2000 ⁹⁶ NA DBRCT	NR/CND	NR/CND	Yes	19% lost to follow-up; 59% did not complete 12 weeks36	1% for lost to follow up and for completing 12 weeks	Yes	No	Yes

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Fawcett, 2000 ⁹⁷ NA DBRCT	NR/CND	NR/CND	Yes	53% (93/175 did not complete 3 months); 11% post-randomization exclusions; missing data/lost to follow-up NR, but Table 2 suggests very low among the 156/175 used for their ITT sample	19; 2 (for missing data first 3 mths for alcohol consumption outcomes)	Yes	No	Yes
Tempesta, 2000 ⁹⁸ NA DBRCT	NR/CND	NR/CND	Yes, for most characteristic; not for previous treatment for alcoholism (see comment)	26% did not complete; 9% for lost to follow-up	2%; 0% for lost to follow-up	No	No	Yes
Chick, 2000 ⁹⁹ NA DBRCT	NR/CND	NR/CND	Yes	16% not interviewed at end of medication phase; 32% lost to follow up or missed many appointments; 65% did not complete 6-month study	5% for lost to follow up or missed many appointments	No	No	Yes
Anton, 1999 ¹⁰⁰ , Anton, 2001 ¹⁰¹ NA DBRCT	NR/CND	NR/CND	Yes	17 (but all but 2 subjects, 1.5%, had week 12 drinking data)	9	No	No	Yes
George, 1999 ¹⁰² NA DBRCT	NR/CND	Yes	NR/CND	42% completed 1 year; 34% lost to follow up	NR/CND	Yes	No	Yes
Besson, 1998 ¹⁰³ NA DBRCT	NR/CND	NR/CND	Yes	30 at 90 days; 65 at 360 days	6 at 90 days; 0 at 360 days	No at 90 days; Yes by 360 days	Yes	NR/CND

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Poldrugo, 1997 ¹⁰⁴ NA DBRCT	NR/CND	NR/CND	Yes	4% lost to follow-up; 55% did not complete 6 months (top reasons were severe relapse, non-compliance, and refusal to continue	0 for lost to follow-up; 15% for completing 6 months (most of difference accounted for by higher severe relapse rate in placebo group)	No	No	NR/CND
Oslin, 1997 ¹⁰⁵ NA DBRCT	NR/CND	NR/CND	Yes	39% did not complete; 20% with some missing data (lost to follow-up or dropped out)	10%; 7%	No	No	Yes
Volpicelli, 1997 ¹⁰⁶ NR DBRCT	Yes	NR/CND	Yes	27% did not complete	0	No	No	NR/CND (for therapy co-intervention); Yes for medication
Cornelius, 1997 ¹⁰⁷ , Cornelius, 1995 ¹⁰⁸ NA DBRCT	NR/CND	NR/CND	No	10	NR/CND	No	No	Yes
Pelc, 1997 ¹⁰⁹ NA DBRCT	NR/CND	NR/CND	NR/CND ("no statistical differences" was reported, but data not provided)	37% did not complete the study; 14% lost to follow-up	18% for not completing; 14.7% for lost to follow-up	No	No	Yes
Sass, 1996 ¹¹⁰ NA DBRCT	NR/CND	Yes	Yes	20% lost to follow-up; 51% did not complete 48 weeks	1.5% for lost to follow-up; 18% for completing	No	No	Yes
Kabel, 1996 ¹¹¹ NA DBRCT	NR/CND	NR/CND	NR/CND	42% did not complete 12 weeks (including those who dropped out before discharge); loss to follow-up NR	10	Yes	No	Yes
Whitworth, 1996 ¹¹² NA DBRCT	Yes	Yes	Yes	15% for loss to follow-up; 60% did not complete double-blind treatment	1.4% for lost to follow-up	No	No	Yes

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

[illegible]

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Kranzler, 1994 ¹²¹ NA DBRCT	NR/CND	NR/CND	No	31% did not complete 12 weeks; loss to follow-up not totally clear, but was 13% or less (based on review of reasons for not completing)	30.6% for not completing; NR for loss to follow-up	Yes	No	Yes
Malcolm, 1992 ¹²² NA DBRCT	NR/CND	NR/CND	Yes	10% lost to follow-up	3	No	No	Yes
O'Malley, 1992 ¹²³ ; O'Malley, 1996 ¹²⁴ NA DBRCT	NR/CND	NR/CND	No	35% did not complete; 7% were not included in analyses	8% for did not complete (NTX vs. placebo); 9.6% for inclusion in analyses (NTX vs. placebo)	No	No	Yes
Lhuintre, 1990 ¹²⁵ NA DBRCT	NR/CND	NR/CND	Yes	37% drop-outs	<1% drop-outs	Yes	No	NR/CND
Fuller, 1986 ¹²⁶ NA DBRCT	Yes	Yes	Yes	5	<5% across three groups	No	No	Yes
Lhuintre, 1985 ¹²⁷ NA DBRCT	NR/CND	NR/CND	NR/CND; only age, ggt and MCV level reported	11% lost to follow-up; 18% did not complete	2% for lost to follow- up; 7% for did not complete	No	Yes	NR/CND
Ling, 1983 ¹²⁸ NA DBRCT	NR/CND	NR/CND	NR/CND	57% did not complete 12 week study; 55% lost to follow-up	3% for completion of study; 22% for lost to follow-up	Yes	No	NR/CND
Fuller, 1979 ¹²⁹ NA DBRCT	Yes	NR/CND	NR/CND (no data provided; per authors, groups were similar at baseline)	2% for final assessment after 1 year; 18% for regular bimonthly and final assessments	NR/CND	No	No	Yes
Gual, 2001 ¹³⁰ NA DBRCT	NR/CND	NR/CND	Yes	16% lost to follow-up; 35% non-completers	4% lost to follow-up; 7% non-completers	No	No	Yes

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Coskunol, 2002 ¹³¹ NA DBRCT	Yes	NR/CND	Yes	0% lost to follow-up (3 0 left study because they developed depression; appears they were included in the analysis)		No	No	Yes
Kranzler, 2013 ¹³² NA SSGA	NR/CND	NR/CND	NR/CND (data provided for sample as a whole, and authors report that there was no differences between groups but data not given)	15% of initial sample did not complete trial, but outcomes were available on entire sample. For this analysis, 74.6% of possible person days of drinking were included. Incomplete drinking data was not included.	NR/CND	No	No	Yes
Ahmadi, 2002 ¹³³ , Ahmadi, 2004 ¹³⁴ NA DBRCT	NR/CND	NR/CND	NR/CND (no data provided; per authors, groups were similar at baseline)	NR/CND	NR/CND	NR/CND	NR/CND	NR/CND
Mason, 1999 ¹³⁵ NA DBRCT	NR/CND	NR/CND	Yes	35% noncompleters; about 10% lost to follow-up	1	No	No	Yes
Geerlings, 1997 ¹³⁶ NA DBRCT	NR/CND	NR/CND	Yes	15% lost to follow up; 64% did not complete the study (most common reason was relapse leading to hospitalization)	1% lost to follow up; 10% for completing the study	No	No	Yes

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Mason, 1994 ¹³⁷ NA DBRCT	NR/CND	NR/CND	NR/CND; study gives means for demographic and lab values for sample as a whole and reports none were statistically significant, but does not provide data.	62% did not complete; missing data and lost to follow up NR	14%	Yes	No	Yes
Tollefson, 1991 ¹³⁸ , Tollefson, 1992 ¹³⁹ NA DBRCT	NR/CND	NR/CND	Yes, in regards to alcohol related items. Those randomized to the active drug was reported to be more likely to have used prior benzodiazepines but rates of prior usage were not provided.	73% "dropped out"; 1% were lost to follow-up; 16% were excluded from the analysis because they did not complete at least 4 weeks of treatment	Twenty two % "dropped out" (more in placebo group); there was no differential loss to follow-up or number of participants completing 4 weeks.	No	No	Yes
Lee, 2001 ¹⁴⁰ NA DBRCT	NR/CND	NR/CND	Yes	66% did not complete 12 weeks; 26% did not have any drinking data	18%; 15%	Yes	No	Yes
Carroll, 1993 ¹⁴¹ NA OLRCT	NR/CND	NR/CND	NR/CND- study says groups were comparable, but data not presented.	67	22	Yes	No	Yes
Morgenstern, 2012 ¹⁴² NA DBRCT	Yes	NR/CND	No, but relatively small differences	16% discontinued treatment; 7% were unavailable for follow-up.	4	No	No	Yes

Table C-1. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Was Trial Name Design	Was randomization adequate?	Was allocation concealment adequate?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Did the study have cross- overs or contamination raising concern for bias?	Was intervention fidelity adequate?
Pelc, 1996 ¹⁴³ , Pelc, 1992 ¹⁴⁴	NR/CND	NR/CND	Yes	45% lost to follow-up by day 90; 65% by day 180	17%; 21%	Yes	No	NR/CND

^a Unable to determine exact differential attrition because they don't report number by group for all 4 groups for how many completed the trial; the flowchart provides number that completed all visits and number that completed week 12 assessments or were on meds for at least 10 weeks (but does not separate the latter group). 24% differential attrition is based on 14/22 vs. 16/20 vs. 16/22 vs. 21/24 (who completed all visits/assessments or were on meds for at least 10 weeks). Article reports data in another place suggesting differential attrition of 20% between all those on desipramine (65% completed the trial) and those on paroxetine (45% completed the trial).

Abbreviations: CND = cannot determine; DBRCT = double-blind randomized controlled trial; NA = not applicable; NR = not reported; OLRCT = open-label randomized controlled trial; SBRCT = single-blind randomized controlled trial; SSGA = secondary or subgroup analysis

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Petrakis, 2012 ¹ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	NR ^a /CND	High	High risk of attrition bias with almost 45% attrition and over 20% differential attrition, along with method of handling missing data; method of randomization and allocation concealment NR
Kranzler, 2012 ² NA SSGA	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Some information in companion Kranzler ⁸
Fogaca, 2011 ³ NA DBRCT	NR/CND	NR/CND	Yes	Yes	NR/CND	No	No	High	High risk of attrition bias, completer's analysis (excluded 37/80 patients after randomization); methods of randomization and allocation concealment NR; unclear method of measurement for consumption outcomes
Ralevski, 2011 ⁴ ; Ralevski, 2001 ⁵ NA DBRCT	No	NR/CND	Yes	Yes	Yes	NR/CND	NR/CND	High	High risk of attrition bias; some baseline differences in sex (all females in the placebo group) and very small sample size of 23; methods of randomization and allocation concealment NR; unclear how missing data was handled; no reporting of masking outcome assessors
Wolwer, 2011 ⁶ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Anton, 2011 ⁷ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	No	No	Medium	Note on statistical methods and missing data: 4 post-randomizations excluded; missing data due to dropout censored, but very low percentage of subjects
Kranzler, 2011 ⁸ DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Florez, 2011 ⁹ NA OLRCT	Yes	NR/CND	No	No	Yes	Yes	NR/CND	High	Open-label trial of topiramate and naltrexone; no masking of patients, providers or outcome assessors; unclear method of randomization and allocation concealment; For missing data, they report assuming that subjects resumed heavy drinking, but not what was done for the quality of life outcomes that we would be interested in from this article (it's not eligible for our KQ 1b because it's open label)
Garbutt, 2010 ¹⁰ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Stedman, 2010 ¹¹ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	High	Very high attrition (57% in this 12 week study); over half of the subjects did not complete the study; no reporting of methods of randomization or allocation concealment or masking of outcome assessors; some concern for contamination and methods of handling missing data (used LOCF for some outcomes and used available data for some others)

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Kiefer, 2011 ¹² NA SSGA	NR/CND	NR/CND	Yes	Yes	Yes	NR/CND	NR/CND	High	Exploratory, hypothesis-generating study; secondary analysis of data from PREDICT (N=430), a German RCT of ACA, NTX, and placebo designed similar to COMBINE; this study used data from 374/430 (87%) of the subjects; those for whom genotype data was available, but unclear how many of those also provided outcome data; study provides unadjusted association between GATA4 genotype (SNP rs13273672) and relapse over 90 days, and associated the finding with response to ACA; high risk of selection bias and confounding; no reporting of baseline characteristics of the groups being compared (across the genotypes or the medications) other than saying they were not different for sex, age, and age of dependence onset and giving p values for those.
Pettinati, 2010 ¹³ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	Medium	Methods of allocation concealment and masking of outcome assessors NR; some risk of attrition bias; Did not impute anything for missing data, but 84.1% of patients provided drinking reports that were 100% complete, and analyses are time to event analyses
Rubio, 2009 ¹⁴ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	No	No	High	Completer's analysis (N=63 analyzed), not ITT; no approach to handling missing data; methods of randomization and allocation concealment and masking of outcome assessors NR

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Schmitz, 2009 ¹⁵ NA DBRCT	No	Yes	Yes	Yes	Yes	Yes	No	High	High risk of selection bias, primarily due to attrition; only 40.5% of subjects completed at least 6 weeks of treatment and just 24% completed all 12 weeks; median follow up prior to dropout was around 30 days; some baseline differences between groups for sex (lower percentage of males in the naltrexone+CBT+CM group); adherence ranged from 50 to 80%; missing data due to dropout were handled as missing (indicating that nothing was done for missing data due to dropout)
Brown, 2009 ¹⁶ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	No	No	High	High risk of selection bias and confounding; 7 out of 50 post-randomization exclusions; 48% of subjects did not complete the study; inadequate handling of missing data; Groups similar at baseline for demographics, but higher proportion of anticonvulsant, antidepressant, and sedative/hypnotic use in the naltrexone group; methods of randomization and allocation concealment NR; allowed adjustment of medications or addition of new medications raising some concern for contamination

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Longabaugh, 2009 ¹⁷ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	No	Medium	Moderate risk of selection bias and confounding; inadequate handling of missing data; Excluded 32/174 (18.4%) randomized subjects from analyses, although non-differential; some baseline differences between the four groups for marital status, education, abstinent days and heavy drinking days in previous 90 days (possibly a result of not using the sample that was randomized, which may have undermined the randomization); methods of allocation concealment NR
Kranzler, 2009 ¹⁸ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Methods of randomization and allocation concealment NR; unclear if outcome assessors were masked; very little baseline information reported to allow comparing the two groups at baseline
Baltieri, 2008 ¹⁹ ; Baltieri, 2009 ²⁰ NA DBRCT	NR/CND	Yes	Yes	Yes	NR/CND	Yes	Yes	High	High risk of selection bias and confounding; high overall attrition (45% did not complete the 12-week study) and differential attrition; Concern for contamination as the groups had differences in rates of AA participation (the authors provide some adjusted analyses to attempt to address this); Those with insufficient adherence were dropped from the study; Some concern for measurement bias as the study did not report using TLFB method to ascertain drinking outcomes (used self-report to ascertain quantity and frequency, but further details of method NR)

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Florez, 2008 ²¹ NA SSGA	NR/CND	NR/CND	NR/CND	NR/CND	Yes	Yes	Yes	High	Exploratory, hypothesis-generating study; secondary analysis of data from a trial; the analyses really focus on whether the outcomes differ by genotype, combining subjects receiving different treatments for main analyses (so not that directly relevant to our questions); evaluates 6 polymorphisms; relatively small sample to attempt this many exploratory genotype analyses (N=90); high risk of selection bias and confounding; no reporting of baseline characteristics of the groups being compared (across the genotypes); study provides unadjusted associations; no adjustment for potential confounders
Oslin, 2008 ²² NA DBRCT	No	Yes	Yes	Yes	Yes	Yes	Yes	Medium	Methods of randomization and allocation concealment NR; some baseline differences between groups (race), but analyses adjusted for age, race, gender, pretreatment percent of HDDs; only 50% adhered to medication across conditions

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Arias, 2008 ²³ NA SSGA	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes (from parent article)	High	Secondary analysis of data from a trial; evaluates 5 polymorphisms; high risk of selection bias and confounding; used 67% of the subjects from the parent trial (those with complete data and genotype information available); does not report baseline characteristics for the comparisons of interest to this article (the different genotypes); some baseline differences in alcohol consumption for those receiving nalmefene compared with those receiving placebo in this sample (statistical methods did make adjustment for these); inadequate consideration of potential confounding
Martinotti, 2009 ²⁴ NA DBRCT	NR/CND	NR/CND	Yes	Yes	CND	Yes	No	Medium	Data not provided to allow assessment of comparison of groups at baseline (text reports no differences for demographics, etc.); used self-report, but not TLFB to gather consumption data; this head-to-head study used LOCF for missing data, but attrition was not too high and was non-differential
Florez, 2008 ²⁵ NA OLRCT	Yes	No	No	No	Yes	Yes	Yes, for consumption and composite measure (assumed relapse); No, for quality of life measures and other outcomes (nothing done to handle missing data)	High	Open label; no masking; some baseline differences between groups that may bias results in favor of topiramate—including more nicotine addiction in the naltrexone group, higher proportions of family history of alcoholism, personality disorders, and higher alcohol intake; baseline means on some scales show trends toward worse scores for naltrexone (Fagerstrom, OCDS, most EuropASI subscales, EQ-5D); methods of randomization and allocation concealment NR

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
O'Malley, 2008 ²⁶ NA DBRCT	No	Yes	Yes	Yes	Yes	Yes	Yes	Medium	Methods of randomization and allocation concealment NR; 33% did not complete study; adherence was 59 to 67% across groups
Wilens, 2008 ²⁷ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	No	High	High risk of attrition bias; 54% did not complete the study; high differential attrition of 20%; inadequate handling of missing data; results for drinking outcomes reported with censoring of missing data (authors report that they also ran analyses counting lost to follow up as relapsed, but data is not shown); methods of randomization and allocation concealment NR
Brown, 2008 ²⁸ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	No	High	High risk of selection bias, attrition bias; inadequate handling of missing data (used LOCF for early withdrawals); poor reporting of methods; 13/115 randomized subjects (11%) excluded after randomization, and information about attrition not reported for the remaining 102 analyzed; methods of randomization and allocation concealment NR; some baseline differences between groups
Anton, 2008 ²⁹ COMBINE SSGA	Yes	Yes for meds; no for psychosocial treatment	Yes	Yes	Yes	Yes	Yes	Medium	Subgroup analysis of data from COMBINE, by genotype; some risk of selection bias and confounding; subjects not randomized by genotype; missing genotype data for some; nevertheless, key variables seem to be distributed similarly across genotype groups; several strengths in design, conduct, and analyses
Lucey, 2008 ³⁰ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	NR/CND	Medium	63 to 65% received all 6 injections; 74% received at least 4 injections
Anton, 2008 ³¹ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Addolorato, 2007 ³² NA DBRCT	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Medium	Some concern for attrition bias due to differential attrition, and because most subjects counted as relapses in the placebo group were those who dropped out or didn't follow up (accounted for 10/21 relapses) rather than those with actual outcome data confirming relapse
Laaksonen, 2008 ³³ NA OLRCT	Yes for NTX and ACA during continuous phase; No for DIS (67.5%)	NR/CND	No	No	Yes	Yes for some outcomes; no for others (see comments)	No	High for quality of life/KQ 2 outcomes	Open label trial; no masking; Quality of life outcomes were reported for the 52 week timepoint (with less than 50% of subjects reaching that timepoint); inadequate handling of missing data for AUDIT, SADD, QL measures (per-protocol analysis including patients that completed the study); used ITT for primary outcomes (consumption outcomes) but study is not eligible for KQ 1 because it is open label.
Johnson, 2007 ³⁴ Johnson, 2008 ³⁵ NA DBRCT	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Low	High differential attrition, with 61.2% completing the trial in the topiramate group compared with 76.6% in the placebo group, but not concerned that introduces significant risk of bias because they have outcome information for most of the non-completers and imputed missing data with baseline values (which were all heavy drinking), so the analysis would be likely to underestimate the benefit of topiramate, if anything; also, few subjects were actually lost to follow up; statistical analysis methods and approach to handling missing data were good.

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Pettinati, 2008 ³⁶ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Methods of randomization and allocation concealment NR; moderate risk of selection bias due to attrition; <50% had adequate adherence (over 80%) to medication; unclear if outcome assessors were masked
Kampman, 2007 ³⁷ NA DBRCT	No	NR/CND	Yes	Yes	Yes	NR/CND	NR/CND	High	Methods of randomization, allocation concealment, and handling of missing data NR; some baseline differences between groups for race, sex, drinks/drinking day, HAM-D and HAM-A scores; inadequate adherence to medication 70 to 77%; small pilot study
De Sousa, 2008 ³⁸ NA OLRCT	CND	No	No	No	No	Yes	Yes	High	Methods of randomization (by the "qualified statistician") and allocation concealment NR; High risk of ascertainment bias; no masking; Open label trial comparing disulfiram and topiramate; potentially had more effort to ensure adherence in the disulfiram group
Karhuvaara, 2007 ³⁹ NA DBRCT	CND	Yes	Yes	Yes	Yes	Yes	Yes	Medium	
Book, 2008 ⁴⁰ ; Thomas, 2008 ⁴¹ NA DBRCT	Yes	Yes	Yes	Yes	Yes	Yes	CND	Medium	Some concern for attrition bias and missing data in this small (N=42) trial, but attrition was non-differential, and study used mixed model analysis considered robust to non-informative missing data
O'Malley, 2007 ⁴² NA DBRCT	Yes, when calculation based on number of days in treatment	Yes	Yes	Yes	Yes	Yes	Yes	Medium	Methods of randomization and allocation concealment NR; our attrition calculations based on having complete timeline data

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

[illegible]

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Hutchison, 2006 ⁵³ NA SBRCT	Yes	NR/CND	No	Yes	Yes	NR/CND	NR/CND	High	No masking of providers; unclear whether outcome assessors were masked; unclear whether used ITT analysis for the drinking outcomes and how missing data were handled (used ITT for craving outcomes and LOCF for missing data); methods or randomization and allocation concealment NR; baseline differences between the groups being compared (i.e., since the analyses were by genotype subgroups and not by the full groups that subjects were randomized to; e.g. those with DRD4 genotype randomized to olanzapine vs. placebo); no adjustment for baseline differences in the comparison by genotype; would consider the genotype findings to be hypothesis generating exploratory analyses
Huang, 2005 ⁵⁴ NA DBRCT	NR/CND	NR/CND	Yes	Yes	NR/CND	CND	CND	High	High risk of measurement bias and confounding; statistical methods don't report whether they used an ITT or completer's analysis; no description of approach to handling missing data; methods of randomization and allocation concealment NR; no description of ascertainment methods for drinking quantity and frequency; relatively few subjects with missing data because they interviewed those who did not complete the study visits and were able to determine that many of them relapsed, they ultimately had outcome data for 80% of subjects

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
De Sousa, 2005 ⁵⁵ NA OLRCT	NR/CND	No	No	No	NR/CND	Yes	Yes	High	Methods of randomization (by the "qualified statistician") NR; no allocation concealment; High risk of ascertainment bias; no masking; Open label trial comparing disulfiram and acamprosate; potentially had more effort to ensure adherence in the disulfiram group
Anton, 2005 ⁵⁶ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	NR/CND	Medium	Therapists were blind to drug assignment but not therapy type, and since the drug is our treatment of interest, we considered the care providers masked.
Petrakis, 2005 ⁵⁷ Ralevski, 2007 ⁵⁸ Petrakis, 2007 ⁵⁹ Petrakis, 2006 ⁶⁰ VA MIRECC DBRCT	Yes	NR/CND	Mixed (yes for NTX, no for DIS)	Mixed (yes for NTX, no for DIS)	Yes	Yes	NR/CND	Medium for NTX vs. pbo High for DIS vs. NTX or pbo	For the DIS comparisons, high risk of ascertainment bias, with no masking; DIS was open-label.
Garbutt, 2005 ⁶¹ NA DBRCT	See comment	Yes	Yes	Yes	Yes	Yes	No	Medium	64% received all 6 injections; 74% received at least 4 injections. Moderate risk of attrition bias due to dropouts, but non-differential.
Brady, 2005 ⁶² NA DBRCT	CND	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Salloum, 2005 ⁶³ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Medium to low risk of bias
Killeen, 2004 ⁶⁴ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	NR/CND	Medium	

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
De Sousa, 2004 ⁶⁵ NA OLRCT	Yes	No	No	No	Yes	Yes	Drop-out considered relapse	High	
Schmitz, 2004 ⁶⁶ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Restricted Maximum Likelihood Estimation, repeated ANCOVA & survival analyses	High	High risk of selection bias and confounding; high overall attrition, unclear differential attrition and missing data, methods of randomization, allocation concealment, and masking of outcome assessors NR; unclear why patients dropped out and if they were included in the analysis
Johnson, 2004 ⁶⁷ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	NR/CND	NR/CND	High	High risk of selection bias and confounding. Groups were not similar at baseline, with differences for sex and higher baseline heavy drinking days for the placebo group. Not surprising that groups were different at baseline in this small, pilot study with 25 NTX subjects and 5 placebo subjects. High attrition. Methods of statistical analyses and handling of missing data NR.
Johnson, 2004 ⁶⁸ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	generalized estimating equations	Medium	
Kranzler, 2004 ⁶⁹ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Anton, 2004 ⁷⁰ NA DBRCT	Yes	Yes	Yes	Yes	Yes	Yes	No	Medium	
Guardia, 2004 ⁷¹ NA DBRCT	No	Yes	Yes	Yes	Yes	Yes	No	Medium	Moderate risk of attrition bias and inadequate handling of missing data. Missing data were not replaced, but amount of missing data may be very low.

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Chick, 2004 ⁷² NA DBRCT	NR/CND	Yes	Yes	Yes	Yes	Yes	Mixed	Medium	Moderate risk of attrition bias; some LOCF used for missing data that might introduce bias, but study found trend for fluvoxamine group to do worse than placebo.
Baltieri, 2004 ⁷³ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	All missing data entered as nonabstinent; the ASI includes a field for "Alcohol-any use at all" allowing a reasonably valid and reliable ascertainment
Petrakis, 2004 ⁷⁴ ; Ralevski, 2006 ⁷⁵ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Methods of randomization and allocation concealment NR; some baseline differences between groups for drinking; low adherence; masking of outcome assessors NR
Gual, 2003 ⁷⁶ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Moak, 2003 ⁷⁷ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Ballidin, 2003 ⁷⁸ NA DBRCT	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Low	
Johnson, 2003 ⁷⁹ Ma, 2006 ⁸⁰ ; Johnson, 2004 ⁶⁸ NA DBRCT	Yes	Yes	Yes	Yes	Yes	Yes	CND	Medium	No completely clear how much missing data for consumption outcomes there was; methods of handling missing data—used data reduction technique taking mean of weeks 1 through 12, weighted by number of study weeks completed with non-missing data; unclear how this would compare with imputing heavy drinking for missing data
Kiefer, 2003 ⁸¹ Kiefer, 2005 ⁸² NA DBRCT	NR/CND	Yes	Yes	Yes	Yes	Yes	Yes	Low	

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Gastpar, 2002 ⁸³ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	Medium	
Guardia, 2002 ⁸⁴ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	No	Medium	Risk of attrition bias, but non-differential; some were excluded post-randomization and not evaluated; apparently censored dropouts in the survival analysis.
Brady, 2002 ⁸⁵ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	Medium	Some concern with approach to handling missing data; some LOCF using previous week's drinking data was used for some missing data; for other missing data (collected monthly), they used monthly group means
Latt, 2002 ⁸⁶ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	NR/CND	Medium	Moderate risk of attrition bias; unclear how missing values were imputed for some analyses
Morris, 2001 ⁸⁷ NA DBRCT	Yes	Yes	Yes	Yes	Yes	Yes	No	Medium	Some baseline differences, with NTX patients drinking 15 more drinks/wk than placebo; inadequate handling of missing data
Krystal, 2001 ⁸⁸ VACS 425 DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	Medium	Inadequate handling of missing data, but relatively low % without complete or partial data (10%) that were not included in the analyses, and non-differential missing data.
Monti, 2001 ⁸⁹ ; Rohsenow, 2007 ⁹⁰ ; Rohsenow, 2000 ⁹¹ NA DBRCT	No	Yes	Yes	Yes	Yes	Yes	Yes	Medium	Rated on basis of medication part of the study (not the preceding psychological treatment part)
Monterosso, 2001 ⁹² NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Rubio, 2001 ⁹³ NA SBRCT	Yes	Yes	No	No	Yes	Yes	Yes	High	Significantly more patients in the acamprosate group were prescribed disulfiram during the course of the study.
Heinala, 2001 ⁹⁴ NA DBRCT	NR/CND	NR/CND	NR/CND	Yes	Yes	NR/CND	NR/CND	High	High risk of selection bias, attrition bias, and confounding. No description of randomization, allocation concealment, outcome assessor masking, or details of statistical methods. Methods section does not include any information on statistical analyses. Patient characteristics according to treatment group NR. High rate of overall attrition with no reporting of differential attrition and inadequate description of how missing data was handled.
Pettinati, 2001 ⁹⁵ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	NR/CND	Unclear	High overall attrition (42%) and 12% differential attrition for study completion; and degree of missing data/loss to follow-up NR for alcohol consumption outcomes; unclear methods of handling missing data for alcohol consumption outcomes.
Chick, 2000 ⁹⁶ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Fawcett, 2000 ⁹⁷ NA DBRCT	Yes	No	Yes	Yes	Yes	Yes	Yes	Medium	
Tempesta, 2000 ⁹⁸ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Chick, 2000 ⁹⁹ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Anton, 1999 ¹⁰⁰ , Anton, 2001 ¹⁰¹ NA DBRCT7	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
George, 1999 ¹⁰² NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	No	High	High risk of selection bias; inadequate handling of missing data; censored those lost to follow up (34% of subjects); differential loss to follow-up NR
Besson, 1998 ¹⁰³ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Use of disulfiram (voluntary, not randomized) was allowed; randomization was stratified by disulfiram use. Missing data was assumed to be relapse.
Poldrugo, 1997 ¹⁰⁴ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Oslin, 1997 ¹⁰⁵ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	NR/CND	Medium	Unclear handling of missing data, but non-differential missing data; methods of randomization and allocation concealment NR
Volpicelli, 1997 ¹⁰⁶ NR DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	CND	Medium	
Cornelius, 1997 ¹⁰⁷ , Cornelius, 1995 ¹⁰⁸ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	Medium	LOCF used for missing data, but just 5 subjects
Pelc, 1997 ¹⁰⁹ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Slightly high differential loss to follow-up, but overall loss to follow-up was low and the higher loss to follow-up was in the placebo group, who also had higher rate of severe relapse
Sass, 1996 ¹¹⁰ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Kabel, 1996 ¹¹¹ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	Yes	High	High overall attrition (but unclear how many of those were lost to follow-up and had missing data) and high risk of confounding; 15% post-enrollment exclusions (of an already very small sample); Unable to determine comparability of groups at baseline--along with small sample size raises concern for selection bias/confounding.
Whitworth, 1996 ¹¹² NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Malec, 1996 ¹¹³ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	High	High overall attrition; 13% differential attrition; and inadequate handling of missing data; completer's analysis.
Mason, 1996 ¹¹⁴ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	High	High risk of attrition bias, inadequate handling of missing data; 28% of subjects that dropped out in the first 2 weeks were not included in analyses. Methods of randomization and allocation concealment NR.
McGrath, 1996 ¹¹⁵ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Some baseline differences for % married and days of drinking heavily and drinks per drinking day. Missing data was handled with LOCF; but participants were not required to be abstinent before study entry. Abstinence of > 2 weeks before randomization was an exclusion criteria.
Tiihonen, 1996 ¹¹⁶ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	High	High overall and differential attrition; inadequate handling of missing data; unclear methodology for randomization, allocation concealment
Kranzler, 1995 ¹¹⁷ NA DBRCT	No	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Paille, 1995 ¹¹⁸ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Study counted those lost to follow-up as not abstinent.
Naranjo, 1995 ¹¹⁹ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	High	Completer's analysis (62/99); high attrition; inadequate handling of missing data
Volpicelli, 1995 ¹²⁰ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	NR/CND	Unclear	Unclear risk of selection bias, confounding, and attrition bias. Baseline characteristics are not reported by treatment group. Inadequate description of handling of missing data. No information is provided regarding attrition or differential attrition. Methods of randomization, allocation concealment, and masking outcome assessors NR.
Kranzler, 1994 ¹²¹ NA DBRCT	Yes	Yes	Yes	Yes	Yes	Yes	NR/CND	Medium	.95% of subjects were interviewed at study completion to obtain information on consumption outcomes.
Malcolm, 1992 ¹²² NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
O'Malley, 1992 ¹²³ ; O'Malley, 1996 ¹²⁴ NA DBRCT	Yes	Yes	Yes (to medication, not therapy)	Yes	Yes	Yes	Mixed	Medium	Subjects randomized to supportive therapy had more severe alcohol problems and drank more alcohol per occasion during baseline compared to those randomized to supportive psychotherapy; inadequate handling of missing data for some analyses; methods of randomization and allocation concealment NR.

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Lhuintre, 1990 ¹²⁵ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	NR/CND	NR/CND	Unclear	Unclear analytic methods and methods of handling missing data; some indications that this is a completers analysis, but unclear; 37% of study participants dropped-out; although non-differential attrition. Methods of randomization, allocation concealment, and masking of outcome assessors NR.
Fuller, 1986 ¹²⁶ NA DBRCT	No	Yes	Yes	Partially	Yes	Yes	No	Medium	Subjects receiving 250 or 1 mg doses of disulfiram were masked to the dose they received, but told they were receiving disulfiram (aim was to control for implied threat of the disulfiram-ethanol reaction); subjects receiving riboflavin were told they were being given a vitamin; missing data censored (if no interview obtained, they were considered to be abstinent until censored) and did not impute assumed lapse/relapse, but relatively little missing data.
Lhuintre, 1985 ¹²⁷ NA DBRCT	NR/CND	NR/CND	Yes	Yes	NR/CND	No	No	High	High risk of selection bias and confounding; medium to high risk of ascertainment bias; completers-only analysis (70/85 randomized subjects in the analysis); methods of randomization, allocation concealment, and consumption outcome assessment NR; inadequate handling of missing data; some concern for contamination because of the use of meprobamate; unable to assess similarity of groups at baseline

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Ling, 1983 ¹²⁸ NA DBRCT	NR/CND	NR/CND	Yes	Yes	NR/CND	Yes	No for most outcomes; Yes for return to heavy drinking	High	High risk of selection bias and confounding, primarily due to attrition; very high overall and differential loss to follow-up; inadequate handling of missing data for most outcomes (e.g., completers analysis for everything in the Table); methods of randomization, allocation concealment, and masking outcome assessors NR; unclear whether consumption outcomes used valid and reliable measures (just reports that it was self-report, but no description of timeline follow back or other details).
Fuller, 1979 ¹²⁹ NA DBRCT	No	Yes	Yes	Partially	Yes	Yes	NR/CND	Medium	Subjects receiving 250 or 1 mg doses of disulfiram were masked to the dose they received, but told they were receiving disulfiram (aim was to control for implied threat of the disulfiram-ethanol reaction); subjects receiving riboflavin were told they were not receiving disulfiram
Gual, 2001 ¹³⁰ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Coskunol, 2002 ¹³¹ NA DBRCT	Yes	Yes	Yes	Yes	Yes	Yes	NR/CND	Medium	
Kranzler, 2013 ¹³² NA SSGA	Yes	NR/CND; daily drinking data was recorded electronically by participants.	Yes	Yes	Yes	Yes	No	Medium	This analysis looks at genetic variation and the effect on craving and subsequent drinking. Data that was incomplete was not included for some outcomes.

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Ahmadi, 2002 ¹³³ , Ahmadi, 2004 ¹³⁴ NA DBRCT	NR/CND	NR/CND	Yes	Yes	Yes	Yes	NR/CND	Unclear	Unclear risk of bias due to limited reporting of methods; methods of randomization, allocation concealment, and handling of missing data NR; baseline characteristics of groups and loss-to-follow up data NR. Primary outcome was abstinence (completers); those who relapsed were non-completers. It is not clearly stated whether outcome data is available for all participants, or whether those who were not available for follow-up were considered to be relapsed.
Mason, 1999 ¹³⁵ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	
Geerlings, 1997 ¹³⁶ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	Yes	Medium	Although his study had a high rate of non-completers, they have follow-up information for most of those subjects, and all subjects were considered to be non-abstinent for the period during which there was missing data.
Mason, 1994 ¹³⁷ NA DBRCT	NR/CND; patients were discontinued from the study if they were not adherent	NR/CND	Yes	Yes	Yes	Yes	No	High	Overall attrition very high, and may have substantially affected the findings given the small sample size (N=21). inadequate handling of missing data, and unclear how much missing data for consumption outcomes
Tollefson, 1991 ¹³⁸ , Tollefson, 1992 ¹³⁹ NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	No	No	High	For participants who did not complete 4 weeks of study duration, a LOCF analysis was used for drinking outcomes. It is unclear whether there was an attempt to determine drinking outcomes after participants dropped out of the study.

Table C-2. Risk of bias assessment for RCTs and related secondary/subgroup analyses (continued)

Author, Year Trial name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were care providers masked?	Were patients masked?	Were outcome measures equal, valid, and reliable?	Did the study use acceptable statistical methods?	Was an appropriate method used to handle missing data?	Risk of bias	Comments
Lee, 2001 ¹⁴⁰ NA DBRCT	NR/CND	NR/CND	Yes, but NTX and placebo pills not identical	Yes	Yes	No	No	High	High risk of selection bias and confounding; high rate of overall and differential attrition; inadequate handling of missing data; methods of randomization and allocation concealment NR; LOCF analysis used which included some, but not all non-completers
Carroll, 1993 ¹⁴¹ NA OLRCT	NR/CND	Yes	No	No	Yes	Yes	No	High	Very high rate of attrition; inadequate description of how missing data was handled.
Morgenstern, 2012 ¹⁴² NA DBRCT	Yes	NR/CND	Yes	Yes	Yes	Yes	No	Medium	
Pelc, 1996 ¹⁴³ , Pelc, 1992 ¹⁴⁴	NR/CND	NR/CND	Yes	Yes	Yes	Yes	Yes	High	High risk of selection bias and confounding, primarily due to potential attrition bias due to high overall (65% loss to follow-up) and high differential attrition; methods of randomization and allocation concealment NR

^a Used mixed effects model, assuming that missing data were missing at random, but unable to determine if that is true from the article, and the study had high differential attrition

Abbreviations: CND = cannot determine; DBRCT = double-blind randomized controlled trial; NA = not applicable; NR = not reported; OLRCT = open-label randomized controlled trial; SBRCT = single-blind randomized controlled trial; SSGA = secondary or subgroup analysis

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
Petrakis, 2012 ¹ NA DBRCT	No	No	Yes	Yes	High	Used a modified version of the Systematic Assessment for Treatment Emergent Events, but don't describe details or what was modified; Some of the same reasons for high risk of bias as for benefits questions, primarily related to attrition bias
Fogaca, 2011 ³ NA DBRCT	No	No	NR/CND	Yes	High	
Ralevski, 2011 ⁴ ; Ralevski, 2001 ⁵ NA DBRCT	No	No	Yes	Yes	High	
Anton, 2011 ⁷ NA DBRCT	No	No	NR/CND	Yes	Medium	
Kranzler, 2011 ⁸ DBRCT	No	No	NR/CND	Yes	Medium	Assessed AEs at every visit with self-reported questionnaire; no further details reported
Florez, 2011 ⁹ NA OLRCT	No	Yes	Yes	Yes	High	See comments for effectiveness risk of bias assessment
Garbutt, 2010 ¹⁰ NA DBRCT	No	Yes	NR/CND	Yes	Medium	
Stedman, 2010 ¹¹ NA DBRCT	Mixed	No	Yes	Yes	High	Unclear how most AEs were identified (implication is voluntary self-report); used specific instruments for EPS and to classify AEs that were reported; same concerns as with efficacy assessment regarding attrition, contamination, etc.
Pettinati, 2010 ¹³ NA DBRCT	No	No	Yes	Yes	Medium	Used Systematic Assessment for Treatment Emergent Effects, no other details reported
Rubio, 2009 ¹⁴ NA DBRCT	Yes	Yes	Yes	Yes	Medium	Describe using an interview that assessed 38 specific AEs and open-ended questions to assess unexpected AEs; in the results, only withdrawals due to AEs are reported (no specific AEs); unlike the benefits analyses for alcohol consumption (which were only of completers), the AEs reported do include the full sample
Schmitz, 2009 ¹⁵ NA DBRCT	No	No	NR/CND	Yes	High	See comments for efficacy assessment. AEs were evaluated by study nurse and physician; article reports that it included a "standardized reporting system when appropriate", but no further details

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms (continued)

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
Brown, 2009 ¹⁶ NA DBRCT	No	No	NR/CND	Yes	High	Only information reported is that side effect assessments were repeated at each weekly appointment
Longabaugh, 2009 ¹⁷ NA DBRCT	No	Yes	NR/CND	Yes	Medium	
Kranzler, 2009 ¹⁸ NA DBRCT	Yes	Yes	Yes	Yes	Medium	Self report screening questionnaire was followed by a nurse's inquiry concerning the presence of 11 AEs commonly associated with naltrexone
Baltieri, 2008 ¹⁹ ; Baltieri, 2009 ²⁰ NA DBRCT	No	No	NR/CND	Yes	High	See comments for efficacy/effectiveness risk of bias also
Oslin, 2008 ²² NA DBRCT	No	NR/CND	NR/CND	Yes	Medium	Minimal description; AEs were monitored by the research physician's probing for side effects commonly associated with NTX
Martinotti, 2009 ²⁴ NA DBRCT	No	Yes, for EKG, UA, blood tests; No, for symptoms	NR/CND	Yes	Medium	
Florez, 2008 ²⁵ NA OLRCT	Yes	No	NR/CND	Yes	High	Used UKU Side Effect Rating Scale (which prespecifies a list of potential harms), but unclear how it was used (who assessed the side effects or completed the scale; whether it was a structured interview or just relied on medical records, whether the person completing this was blinded [likely not, in this open label trial], how involved the patients were in the process, etc.)
O'Malley, 2008 ²⁶ NA DBRCT	NR/CND	No	NR/CND	Yes	Medium	
Wilens, 2008 ²⁷ NA DBRCT	No	No	NR/CND	Yes	High	
Brown, 2008 ²⁸ NA DBRCT	No	No	NR/CND	Yes	High	

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms (continued)

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
Lucey, 2008 ³⁰ NA DBRCT	Yes, for liver tests, vital signs, lab measuremen ts; No, for symptoms and physical exam	Yes, for liver tests, vital signs, lab measurements; No, for symptoms and physical exam	Yes, for liver tests, vital signs, lab measurements; No, for symptoms and physical exam	Yes	Medium	
Anton, 2008 ³¹ NA DBRCT	Yes for EPS; No for others	Yes for EPS; No for others	NR/CND	Yes	Medium	
Addolorato, 2007 ³² NA DBRCT	NR/CND	No	NR/CND	Yes	Medium	
Laaksonen, 2008 ³³ NA OLRCT	No	No	NR/CND	Yes	High	No masking; open label trial; only reports that harms were elicited at each visit and recorded in the drinking diary; labs were drawn at wk 0, 6, and 52, but very high attrition by week 52
Johnson, 2007 ³⁴ Johnson, 2008 ³⁵ NA DBRCT	Yes for vital signs and lab tests; No for symptoms	No	Yes for vital signs and lab tests; NR/CND for symptoms	Yes	Medium	
Pettinati, 2008 ³⁶ NA DBRCT	NR/CND	Yes	Yes	Yes	Medium	Harms were prespecified; not clear if they were defined; see comments for effectiveness assessment of risk of bias
Kampman, 2007 ³⁷ NA DBRCT	No	No	No	Yes	High	Results describe that adverse events were assessed at each visit as NPs asked if there were any changes in their health since the last visit
De Sousa, 2008 ³⁸ NA OLRCT	No	No	No	Yes	High	
Karhuvaara, 2007 ³⁹ NA DBRCT	NR/CND	No	NR/CND	Yes	Medium	
Book, 2008 ⁴⁰ , Thomas, 2008 ⁴¹ NA DBRCT	NR/CND	No	NR/CND	Yes	Medium	

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms (continued)

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
O'Malley, 2007 ⁴² NA DBRCT	No	Yes for some (depression, liver enzymes); No for self- reported adverse effects	NR/CND	Yes	Medium	
Nava, 2006 ⁴⁴ NA OLRCT	No	No	NR/CND	Yes	High	
Morley, 2006 ⁴⁵ Morley, 2010 ⁴⁶ NA DBRCT	No	Yes	NR/CND	Yes	Medium	
Anton, 2006 ⁴⁷ Donovan, 2008 ⁴⁸ LoCastro, 2009 ⁴⁹ Greenfield, 2010 ⁵⁰ Fucito, 2012 ⁵¹ COMBINE DBRCT	Yes	Yes	Yes	Yes	Low	
Mason, 2006 ⁵² NA DBRCT	NR/CND	No	NR/CND	Yes	Medium	Only report that adverse drug events were assessed at every study visit by an open-ended question and coded with the Coding Symbols for Thesaurus of Adverse Reaction Terms (COSTART)
De Sousa, 2005 ⁵⁵ NA OLRCT	No	No	NR/CND	Yes	High	
Petrakis, 2005 ⁵⁷ Ralevski, 2007 ⁵⁸ Petrakis, 2007 ⁵⁹ Petrakis, 2006 ⁶⁰ VA MIRECC DBRCT	Yes	Yes	Yes	Yes	Medium for NTX vs. pbo; High for DIS vs. NTX or vs. pbo	
Garbutt, 2005 ⁶¹ NA DBRCT	NR/CND	Yes	Yes	Yes	Medium	
Salloum, 2005 ⁶³ NA DBRCT	Yes	Yes	Yes	Yes	Low	Somatic symptoms checklist, weekly

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms (continued)

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
Killeen, 2004 ⁶⁴ NA DBRCT	NR/CND	Yes	Yes	Yes	Medium	
De Sousa, 2004 ⁶⁵ NA OLRCT	No	No	NR/CND	Yes	High	
Schmitz, 2004 ⁶⁶ NA DBRCT	NR/CND	Yes	NR/CND	Yes	High	Used preset list of harms, but not clear if those were defined. See comments for efficacy assessment; no usable harms data reported in results
Johnson, 2004 ⁶⁷ NA DBRCT	No	No	NR/CND	Yes	High	Also see comments for efficacy risk of bias
Johnson, 2004 ⁶⁸ NA DBRCT	No	No	NR/CND	Yes	High	
Kranzler, 2004 ⁶⁹ NA DBRCT	No	No	NR/CND	Yes	Medium	Very few details about harms data collection; specific harms were only reported if overall frequency $\geq 10\%$ or significant group difference
Anton, 2004 ⁷⁰ NA DBRCT	No	No	NR/CND	Yes	Medium	
Guardia, 2004 ⁷¹ NA DBRCT	Yes	Yes	Yes	Yes	Medium	See comments for efficacy
Chick, 2004 ⁷² NA DBRCT	No	No	NR/CND	Yes	Medium	
Baltieri, 2004 ⁷³ NA DBRCT	Yes	Yes	Yes	Yes	Medium	
Petrakis, 2004 ⁷⁴ , Ralevski, 2006 ⁷⁵ NA DBRCT	Yes	Yes	Yes	Yes	Medium	See comments for efficacy assessment
Gual, 2003 ⁷⁶ NA DBRCT	No	Yes	No	Yes	High	AEs were spontaneously reported or observed by investigator, then classified.
Moak, 2003 ⁷⁷ NA DBRCT	No	No	NR/CND	Yes	Medium	

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms (continued)

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
Balldin, 2003 ⁷⁸ NA DBRCT	No	Yes	Equal but not valid/reliable	Yes	Medium	
Johnson, 2003 ⁷⁹ Ma, 2006 ⁸⁰ , Johnson, 2004 ⁶⁸ NA DBRCT	Yes	Yes	Yes	Yes	Medium	See comments for efficacy assessment
Kiefer, 2003 ⁸¹ Kiefer, 2005 ⁸² NA DBRCT	No	No	NR/CND	Yes	Medium	Ascertainment techniques for lab measures adequately described, but nothing reported for subjective AEs (e.g., fatigue, diarrhea, etc.)
Gastpar, 2002 ⁸³ NA DBRCT	No	No	NR/CND	Yes	Medium	
Guardia, 2002 ⁸⁴ NA DBRCT	No	No	NR/CND	Yes	Medium	
Latt, 2002 ⁸⁶ NA DBRCT	No	Yes	NR/CND	Yes	Medium	
Morris, 2001 ⁸⁷ NA DBRCT	No	No	NR/CND	Yes	Medium	
Krystal, 2001 ⁸⁸ VACS 425 DBRCT	NR/CND	No	NR/CND	Yes	Medium	
Monti, 2001 ⁸⁹ , Rohsenow, 2007 ⁹⁰ , Rohsenow, 2000 ⁹¹ NA DBRCT	NR/CND	Yes	NR/CND	Yes	Medium	open-ended description of specific symptoms
Rubio, 2001 ⁹³ NA SBRCT	No	No	No	Yes	High	
Heinala, 2001 ⁹⁴ NA DBRCT	No	No	NR/CND	Yes	High	
Pettinati, 2001 ⁹⁵ NA DBRCT	No	No	NR/CND	Yes	Unclear	

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms (continued)

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
Chick, 2000 ⁹⁶ NA DBRCT	No	No	NR/CND	Yes	Medium	
Fawcett, 2000 ⁹⁷ NA DBRCT	No (aside from monitoring for lithium toxicity)	Yes – labs only, not for other harms	Yes- labs only, not for other harms	Yes	Medium	Monitoring for lithium toxicity was prespecified and described. Other harms are reported but no information is given on ascertainment techniques.
Tempesta, 2000 ⁹⁸ NA DBRCT	No	No	Yes	Yes	Medium	Harms were not defined; recorded by spontaneous reporting and by a questionnaire, but it is unclear what the questionnaire asks.
Chick, 2000 ⁹⁹ NA DBRCT	No	No	NR/CND	Yes	Medium	
Anton, 1999 ¹⁰⁰ , Anton, 2001 ¹⁰¹ NA DBRCT	No	Yes	Yes	Yes	Medium	
Besson, 1998 ¹⁰³ NA DBRCT	No	No	NR/CND	Yes	Medium	
Poldrugo, 1997 ¹⁰⁴ NA DBRCT	NR/CND	No	NR/CND	Yes	Medium	Reports using a systematic questionnaire for evaluation of adverse events; details NR
Oslin, 1997 ¹⁰⁵ NA DBRCT	NR/CND	Yes	NR/CND	Yes	Medium	Harms prespecified, used checklist, but not clear if defined
Volpicelli, 1997 ¹⁰⁶ NR DBRCT	NR/CND	Yes	NR/CND	Yes	Medium	Used a side effects checklist, so harms were prespecified, but unclear if they were defined and how they were defined
Cornelius, 1997 ¹⁰⁷ , Cornelius, 1995 ¹⁰⁸ NA DBRCT	No	No	NR/CND	Yes	Medium	
Pelc, 1997 ¹⁰⁹ NA DBRCT	No	Yes	Yes	Yes	Medium	
Sass, 1996 ¹¹⁰ NA DBRCT	No	No	NR/CND	Yes	Medium	

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms (continued)

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
Whitworth, 1996 ¹¹² NA DBRCT	Yes	Yes	NR/CND	Yes	Medium	Asked about 44 AEs (details of the list of 44 and their definitions NR) and rated for severity, and classified into one of seven categories
Malec, 1996 ¹¹³ NA DBRCT	No	No	NR/CND	Yes	High	
Mason, 1996 ¹¹⁴ NA DBRCT	No	No	NR/CND	Yes	High	
McGrath, 1996 ¹¹⁵ NA DBRCT	No	No	NR/CND	Yes	Medium	Harms are reported, but no information is given on ascertainment techniques.
Tiihonen, 1996 ¹¹⁶ NA DBRCT	No	No	NR/CND	Yes	High	
Kranzler, 1995 ¹¹⁷ NA DBRCT	NR/CND	Yes	Yes	Yes	Medium	This study did not prespecify harms, but described using a standardized questionnaire to assess harms.
Paille, 1995 ¹¹⁸ NA DBRCT	No	Yes	Yes	Yes	Medium	
Naranjo, 1995 ¹¹⁹ NA DBRCT	No	No	NR/CND	Yes	High	
Volpicelli, 1995 ¹²⁰ NA DBRCT	No	No	NR/CND	Yes	Unclear	See comments on efficacy assessment
Kranzler, 1994 ¹²¹ NA DBRCT	No	No	NR/CND	Yes	Medium	
Malcolm, 1992 ¹²² NA DBRCT	No	Yes	Yes	Yes	Medium	
O'Malley, 1992 ¹²³ ; O'Malley, 1996 ¹²⁴ NA DBRCT	No	No	NR/CND	Yes	Medium	

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms (continued)

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
Lhuintre, 1990 ¹²⁵ NA DBRCT	Yes	Yes	Yes	Yes	Medium	Used a 44-item questionnaire of somatic complaints; AEs assessment includes those who dropped out due to AEs (whereas it was unclear whether efficacy outcomes only included completers)
Fuller, 1986 ¹²⁶ NA DBRCT	No	Yes	Yes	Yes	Medium	
Lhuintre, 1985 ¹²⁷ NA DBRCT	No	No	NR/CND	Yes	High	
Ling, 1983 ¹²⁸ NA DBRCT	No	No	NR/CND	Yes	High	
Gual, 2001 ¹³⁰ NA DBRCT	No	No	NR/CND	Yes	Medium	
Ahmadi, 2002 ¹³³ , Ahmadi, 2004 ¹³⁴ NA DBRCT	No	No	NR/CND	Yes	Unclear	See comments for efficacy risk of bias assessment
Mason, 1999 ¹³⁵ NA DBRCT	No	No	NR/CND	Yes	Medium	
Geerlings, 1997 ¹³⁶ NA DBRCT	No	No	NR/CND	Yes	Medium	
Mason, 1994 ¹³⁷ NA DBRCT	Only weight loss. No other harms prespecified.	No	NR/CND	Yes	High	Harms were not prespecified and ascertainment techniques were not described.
Tollefson, 1991 ¹³⁸ , Tollefson, 1992 ¹³⁹ NA DBRCT	No	No	NR/CND	Yes	High	
Lee, 2001 ¹⁴⁰ NA DBRCT	No	No (a questionnaire was used, but not described)	NR/CND	Yes	High	

Table C-3. Additional risk of bias questions for RCTs and related secondary/subgroup analyses that report harms (continued)

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias	Comments
Morgenstern, 2012 ¹⁴² NA DBRCT	NR/CND	NR/CND	NR/CND	Yes	High	
Pelc, 1996 ¹⁴³ ; Pelc, 1992 ¹⁴⁴	No	Yes	NR/CND	Yes	High	High risk of selection bias and confounding due to attrition bias. AEs prespecified (checklist used) but not defined. Harms rates only reported for AEs with >5% occurrence. With relatively small Ns, this could be an issue.

Abbreviations: AE = adverse effect; CND = cannot determine; DBRCT = double-blind randomized controlled trial; NA = not applicable; NR = not reported; OLRCT = open-label randomized controlled trial; SBRCT = single-blind randomized controlled trial; SSGA = secondary or subgroup analysis

Table C-4. Risk of bias assessment for observational studies

Author, Year Trial Name Design	Was the sample size adequate?	Were groups recruited from the same source population?	Were groups recruited over the same time period?	Were inc/exc criteria applied equally for all groups?	Were groups similar at baseline?	What was the overall attrition?	What was the differential attrition?	Did the study have overall high attrition or differential attrition raising concern for bias?	Was intervention fidelity adequate?
Coller, 2001 ¹⁴⁵ Prospective cohort	Yes	NR/CND	NR/CND	Yes	Yes	32	<1	Yes	NR/CND
Kim, 2009 ¹⁴⁶ Prospective cohort	No	Yes	NR/CND	Yes	No	49	22	Yes	NR/CND
Narayana, 2008 ¹⁴⁷ Prospective cohort	Yes	Yes	Yes	Yes	Yes, for the few characteristics reported	29	CND exact number, but appears to be about 20% higher in the NTX and ACA groups than the TOP group	Yes	NR/CND
Mutschler, 2012 ¹⁴⁸ Prospective cohort	Yes	Yes	NR/CND	NR/CND	Yes	NR/CND	NR/CND	NR/CND	NR/CND

Abbreviations: ACA = acamprosate; CND = cannot determine; NR = not reported; NTX = naltrexone; TOP = topiramate

Table C-5. Risk of bias assessment for observational studies

Author, Year Trial Name Design	Was adherence to the intervention adequate?	Were outcome assessors masked?	Were outcome measures equal, valid, and reliable?	Were differences between groups taken into account in statistical analysis?	Was confounding adequately accounted for either through study design or statistical analysis?	Was an appropriate method used to handle missing data? Which?	Was time of follow-up equal in both groups?	RISK OF BIAS	Notes; explain “high” ratings
Coller, 2001 ¹⁴⁵ Prospective cohort	Yes	NR/CND	NR/CND	Yes	NR/CND	NR/CND	Yes	Medium	Moderate risk of attrition bias and confounding
Kim, 2009 ¹⁴⁶ Prospective cohort	No	NR/CND	Yes	Yes	Yes	No	Yes	High	High risk of selection bias and confounding; analysis of 32/63 patients in the original cohort who finished the trial and were at least 80% adherent to NTX. The 6 excluded for non-adherence all came from one group.
Narayana, 2008 ¹⁴⁷ Prospective cohort	NR/CND	No	No	NR/CND	NR/CND	NR/CND	Yes	High	Very high differential attrition, completers-only analysis. Inadequate handling of missing data; high risk of selection bias and confounding.
Mutschler, 2012 ¹⁴⁸ Prospective cohort	NR/CND	NR/CND	NR/CND	NR/CND	No	NR/CND	NR/CND	High	High risk of selection bias and confounding

Abbreviations: CND = cannot determine; NR = not reported; NTX = naltrexone

Table C-6. Additional risk of bias questions for observational studies that report harms

Author, Year Trial Name Design	Were harms prespecified and defined?	Were ascertainment techniques for harms adequately described?	Were ascertainment techniques for harms equal, valid and reliable?	Was the duration of follow-up adequate for harms assessment?	Risk of bias
Narayana, 2008 ¹⁴⁷ Prospective cohort	No	No	NR/CND	Yes	High

Abbreviations: CND = cannot determine; NR = not reported;

Table C-7. Risk of bias assessment for systematic reviews and meta-analyses

Author, year	Was the review based on a focused question of interest?	Was the literature search strategy clearly described?	Was there evidence of a substantial effort to search for all relevant research?	Were there explicit inclusion/ exclusion criteria for the selection of studies?	Did at least 2 people independently review studies?	Was the validity of included studies adequately assessed?	Was publication bias assessed?	Was heterogeneity assessed and addressed?
Mason, 2012 ¹⁴⁹	Yes	Yes	Yes	Yes	NR	Yes	No	Mixed
Jorgensen, 2011 ¹⁵⁰	Yes	Yes	No	Yes	NR	Yes	No	Yes
Rosner, 2010 ¹⁵¹	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes
Rosner, 2010 ¹⁵²	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes
Lobmaier, 2008 ¹⁵³	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes
NICE, 2011 ¹⁵⁴	Yes	Yes	Yes	Yes	No	Yes	No	Yes

Abbreviations: NR = not reported

Table C-8. Risk of bias assessment for systematic reviews and meta-analyses

Author, Year	Was the approach used to synthesize the information adequate and appropriate?	Were the author's conclusions supported by the evidence they presented?	RISK OF BIAS	Notes; explain "high" ratings
Mason, 2012 ¹⁴⁹	Yes	Yes	Medium	Unclear whether they had dual independent review for study selection.
Jorgensen, 2011 ¹⁵⁰	Yes	Yes	Medium	Did not search for unpublished studies; did not assess publication bias; unsure about dual review.
Rosner, 2010 ¹⁵¹	Yes	Yes	Medium	Did not have dual independent review of abstracts and full texts
Rosner, 2010 ¹⁵²	Yes	Yes	Medium	Did not have dual independent review of abstracts and full texts
Lobmaier, 2008 ¹⁵³	Yes	Yes	Low	The ROB is low, but only harms data for the subset of alcohol-dependent studies are useful for the purposes of our report (the rest is beyond our scope).
NICE, 2011 ¹⁵⁴	Yes	Yes	Medium	Did not have dual independent review of abstracts and full texts

References for Appendix C

1. Petrakis IL, Ralevski E, Desai N, et al. Noradrenergic vs serotonergic antidepressant with or without naltrexone for veterans with PTSD and comorbid alcohol dependence. *Neuropsychopharmacology*. 2012 Mar;37(4):996-1004. PMID: 22089316.
2. Kranzler HR, Armeli S, Tennen H. Post-treatment outcomes in a double-blind, randomized trial of sertraline for alcohol dependence. *Alcohol Clin Exp Res*. 2012 Apr;36(4):739-44. PMID: 21981418.
3. Fogaca MN, Santos-Galduroz RF, Eserian JK, et al. The effects of polyunsaturated fatty acids in alcohol dependence treatment-a double-blind, placebo-controlled pilot study. *BMC Clin Pharmacol*. 2011;11:10. PMID: 21787433.
4. Ralevski E, O'Brien E, Jane JS, et al. Effects of acamprosate on cognition in a treatment study of patients with schizophrenia spectrum disorders and comorbid alcohol dependence. *J Nerv Ment Dis*. 2011 Jul;199(7):499-505. PMID: 21716064.
5. Ralevski E, O'Brien E, Jane JS, et al. Treatment with acamprosate in patients with schizophrenia spectrum disorders and comorbid alcohol dependence. *J Dual Diagn*. 2011;7(1-2):64-73.
6. Wolwer W, Frommann N, Janner M, et al. The effects of combined acamprosate and integrative behaviour therapy in the outpatient treatment of alcohol dependence: a randomized controlled trial. *Drug Alcohol Depend*. 2011 Nov 1;118(2-3):417-22. PMID: 21621929.
7. Anton RF, Myrick H, Wright TM, et al. Gabapentin combined with naltrexone for the treatment of alcohol dependence. *Am J Psychiatry*. 2011 Jul;168(7):709-17. PMID: 21454917.
8. Kranzler HR, Armeli S, Tennen H, et al. A double-blind, randomized trial of sertraline for alcohol dependence: moderation by age of onset [corrected] and 5-hydroxytryptamine transporter-linked promoter region genotype. *J Clin Psychopharmacol*. 2011 Feb;31(1):22-30. PMID: 21192139.
9. Florez G, Saiz PA, Garcia-Portilla P, et al. Topiramate for the treatment of alcohol dependence: comparison with naltrexone. *Eur Addict Res*. 2011;17(1):29-36. PMID: 20975274.
10. Garbutt JC, Kampov-Polevoy AB, Gallop R, et al. Efficacy and safety of baclofen for alcohol dependence: a randomized, double-blind, placebo-controlled trial. *Alcohol Clin Exp Res*. 2010 Nov;34(11):1849-57. PMID: 20662805.
11. Stedman M, Pettinati HM, Brown ES, et al. A double-blind, placebo-controlled study with quetiapine as adjunct therapy with lithium or divalproex in bipolar I patients with coexisting alcohol dependence. *Alcohol Clin Exp Res*. 2010 Oct;34(10):1822-31. PMID: 20626727.
12. Kiefer F, Witt SH, Frank J, et al. Involvement of the atrial natriuretic peptide transcription factor GATA4 in alcohol dependence, relapse risk and treatment response to acamprosate. *Pharmacogenomics J*. 2011 Oct;11(5):368-74. PMID: 20585342.
13. Pettinati HM, Oslin DW, Kampman KM, et al. A double-blind, placebo-controlled trial combining sertraline and naltrexone for treating co-occurring depression and alcohol dependence. *Am J Psychiatry*. 2010 Jun;167(6):668-75. PMID: 20231324.
14. Rubio G, Martinez-Gras I, Manzanares J. Modulation of impulsivity by topiramate: implications for the treatment of alcohol dependence. *J Clin Psychopharmacol*. 2009 Dec;29(6):584-9. PMID: 19910725.
15. Schmitz JM, Lindsay JA, Green CE, et al. High-dose naltrexone therapy for cocaine-alcohol dependence. *Am J Addict*. 2009 Sep-Oct;18(5):356-62. PMID: 19874153.
16. Brown ES, Carmody TJ, Schmitz JM, et al. A randomized, double-blind, placebo-controlled pilot study of naltrexone in outpatients with bipolar disorder and alcohol dependence. *Alcohol Clin Exp Res*. 2009 Nov;33(11):1863-9. PMID: 19673746.

17. Longabaugh R, Wirtz PW, Gulliver SB, et al. Extended naltrexone and broad spectrum treatment or motivational enhancement therapy. *Psychopharmacology (Berl)*. 2009 Oct;206(3):367-76. PMID: 19639303.
18. Kranzler HR, Tennen H, Armeli S, et al. Targeted naltrexone for problem drinkers. *J Clin Psychopharmacol*. 2009 Aug;29(4):350-7. PMID: 19593174.
19. Baltieri DA, Daro FR, Ribeiro PL, et al. Comparing topiramate with naltrexone in the treatment of alcohol dependence. *Addiction*. 2008 Dec;103(12):2035-44. PMID: 18855810.
20. Baltieri DA, Daro FR, Ribeiro PL, et al. Effects of topiramate or naltrexone on tobacco use among male alcohol-dependent outpatients. *Drug Alcohol Depend*. 2009 Nov 1;105(1-2):33-41. PMID: 19595518.
21. Florez G, Saiz P, Garcia-Portilla P, et al. Association between the Stin2 VNTR polymorphism of the serotonin transporter gene and treatment outcome in alcohol-dependent patients. *Alcohol Alcohol*. 2008 Sep-Oct;43(5):516-22. PMID: 18552399.
22. Oslin DW, Lynch KG, Pettinati HM, et al. A placebo-controlled randomized clinical trial of naltrexone in the context of different levels of psychosocial intervention. *Alcohol Clin Exp Res*. 2008 Jul;32(7):1299-308. PMID: 18540910.
23. Arias AJ, Armeli S, Gelernter J, et al. Effects of opioid receptor gene variation on targeted nalmefene treatment in heavy drinkers. *Alcohol Clin Exp Res*. 2008 Jul;32(7):1159-66. PMID: 18537939.
24. Martinotti G, Di Nicola M, Di Giannantonio M, et al. Aripiprazole in the treatment of patients with alcohol dependence: a double-blind, comparison trial vs. naltrexone. *J Psychopharmacol*. 2009 Mar;23(2):123-9. PMID: 18515460.
25. Florez G, Garcia-Portilla P, Alvarez S, et al. Using topiramate or naltrexone for the treatment of alcohol-dependent patients. *Alcohol Clin Exp Res*. 2008 Jul;32(7):1251-9. PMID: 18482157.
26. O'Malley SS, Robin RW, Levenson AL, et al. Naltrexone alone and with sertraline for the treatment of alcohol dependence in Alaska natives and non-natives residing in rural settings: a randomized controlled trial. *Alcohol Clin Exp Res*. 2008 Jul;32(7):1271-83. PMID: 18482155.
27. Wilens TE, Adler LA, Weiss MD, et al. Atomoxetine treatment of adults with ADHD and comorbid alcohol use disorders. *Drug Alcohol Depend*. 2008 Jul 1;96(1-2):145-54. PMID: 18403134.
28. Brown ES, Garza M, Carmody TJ. A randomized, double-blind, placebo-controlled add-on trial of quetiapine in outpatients with bipolar disorder and alcohol use disorders. *J Clin Psychiatry*. 2008 May;69(5):701-5. PMID: 18312058.
29. Anton RF, Oroszi G, O'Malley S, et al. An evaluation of mu-opioid receptor (OPRM1) as a predictor of naltrexone response in the treatment of alcohol dependence: results from the Combined Pharmacotherapies and Behavioral Interventions for Alcohol Dependence (COMBINE) study. *Arch Gen Psychiatry*. 2008 Feb;65(2):135-44. PMID: 18250251.
30. Lucey MR, Silverman BL, Illeperuma A, et al. Hepatic safety of once-monthly injectable extended-release naltrexone administered to actively drinking alcoholics. *Alcohol Clin Exp Res*. 2008 Mar;32(3):498-504. PMID: 18241321.
31. Anton RF, Kranzler H, Breder C, et al. A randomized, multicenter, double-blind, placebo-controlled study of the efficacy and safety of aripiprazole for the treatment of alcohol dependence. *J Clin Psychopharmacol*. 2008 Feb;28(1):5-12. PMID: 18204334.
32. Addolorato G, Leggio L, Ferrulli A, et al. Effectiveness and safety of baclofen for maintenance of alcohol abstinence in alcohol-dependent patients with liver cirrhosis: randomised, double-blind controlled study. *Lancet*. 2007 Dec 8;370(9603):1915-22. PMID: 18068515.

33. Laaksonen E, Koski-Jannes A, Salaspuro M, et al. A randomized, multicentre, open-label, comparative trial of disulfiram, naltrexone and acamprosate in the treatment of alcohol dependence. *Alcohol Alcohol*. 2008 Jan-Feb;43(1):53-61. PMID: 17965444.
34. Johnson BA, Rosenthal N, Capece JA, et al. Topiramate for treating alcohol dependence: a randomized controlled trial. *JAMA*. 2007 Oct 10;298(14):1641-51. PMID: 17925516.
35. Johnson BA, Rosenthal N, Capece JA, et al. Improvement of physical health and quality of life of alcohol-dependent individuals with topiramate treatment: US multisite randomized controlled trial. *Arch Intern Med*. 2008 Jun 9;168(11):1188-99. PMID: 18541827.
36. Pettinati HM, Kampman KM, Lynch KG, et al. Gender differences with high-dose naltrexone in patients with co-occurring cocaine and alcohol dependence. *J Subst Abuse Treat*. 2008 Jun;34(4):378-90. PMID: 17664051.
37. Kampman KM, Pettinati HM, Lynch KG, et al. A double-blind, placebo-controlled pilot trial of quetiapine for the treatment of Type A and Type B alcoholism. *J Clin Psychopharmacol*. 2007 Aug;27(4):344-51. PMID: 17632217.
38. De Sousa AA, De Sousa J, Kapoor H. An open randomized trial comparing disulfiram and topiramate in the treatment of alcohol dependence. *J Subst Abuse Treat*. 2008 Jun;34(4):460-3. PMID: 17629442.
39. Karhuvaara S, Simojoki K, Virta A, et al. Targeted nalmefene with simple medical management in the treatment of heavy drinkers: a randomized double-blind placebo-controlled multicenter study. *Alcohol Clin Exp Res*. 2007 Jul;31(7):1179-87. PMID: 17451401.
40. Book SW, Thomas SE, Randall PK, et al. Paroxetine reduces social anxiety in individuals with a co-occurring alcohol use disorder. *J Anxiety Disord*. 2008;22(2):310-8. PMID: 17448631.
41. Thomas SE, Randall PK, Book SW, et al. A complex relationship between co-occurring social anxiety and alcohol use disorders: what effect does treating social anxiety have on drinking? *Alcohol Clin Exp Res*. 2008 Jan;32(1):77-84. PMID: 18028529.
42. O'Malley SS, Sinha R, Grilo CM, et al. Naltrexone and cognitive behavioral coping skills therapy for the treatment of alcohol drinking and eating disorder features in alcohol-dependent women: a randomized controlled trial. *Alcohol Clin Exp Res*. 2007 Apr;31(4):625-34. PMID: 17374042.
43. Gelernter J, Gueorguieva R, Kranzler HR, et al. Opioid receptor gene (OPRM1, OPRK1, and OPRD1) variants and response to naltrexone treatment for alcohol dependence: results from the VA Cooperative Study. *Alcohol Clin Exp Res*. 2007 Apr;31(4):555-63. PMID: 17374034.
44. Nava F, Premi S, Manzato E, et al. Comparing treatments of alcoholism on craving and biochemical measures of alcohol consumption. *J Psychoactive Drugs*. 2006 Sep;38(3):211-7. PMID: 17165363.
45. Morley KC, Teesson M, Reid SC, et al. Naltrexone versus acamprosate in the treatment of alcohol dependence: A multi-centre, randomized, double-blind, placebo-controlled trial. *Addiction*. 2006 Oct;101(10):1451-62. PMID: 16968347.
46. Morley KC, Teesson M, Sannibale C, et al. Clinical predictors of outcome from an Australian pharmacological relapse prevention trial. *Alcohol Alcohol*. 2010 Nov-Dec;45(6):520-6. PMID: 20952764.
47. Anton RF, O'Malley SS, Ciraulo DA, et al. Combined pharmacotherapies and behavioral interventions for alcohol dependence: the COMBINE study: a randomized controlled trial. *JAMA*. 2006 May 3;295(17):2003-17. PMID: 16670409.
48. Donovan DM, Anton RF, Miller WR, et al. Combined pharmacotherapies and behavioral interventions for alcohol dependence (The COMBINE Study): examination of posttreatment drinking outcomes. *J Stud Alcohol Drugs*. 2008 Jan;69(1):5-13. PMID: 18080059.
49. LoCastro JS, Youngblood M, Cisler RA, et al. Alcohol treatment effects on secondary nondrinking outcomes and quality of life: the COMBINE study. *J Stud Alcohol Drugs*. 2009 Mar;70(2):186-96. PMID: 19261230.

50. Greenfield SF, Pettinati HM, O'Malley S, et al. Gender differences in alcohol treatment: an analysis of outcome from the COMBINE study. *Alcohol Clin Exp Res*. 2010 Oct;34(10):1803-12. PMID: 20645934.
51. Fucito LM, Park A, Gulliver SB, et al. Cigarette smoking predicts differential benefit from naltrexone for alcohol dependence. *Biol Psychiatry*. 2012;72(10):832-8.
52. Mason BJ, Goodman AM, Chabac S, et al. Effect of oral acamprosate on abstinence in patients with alcohol dependence in a double-blind, placebo-controlled trial: the role of patient motivation. *J Psychiatr Res*. 2006 Aug;40(5):383-93. PMID: 16546214.
53. Hutchison KE, Ray L, Sandman E, et al. The effect of olanzapine on craving and alcohol consumption. *Neuropsychopharmacology*. 2006 Jun;31(6):1310-7. PMID: 16237394.
54. Huang MC, Chen CH, Yu JM, et al. A double-blind, placebo-controlled study of naltrexone in the treatment of alcohol dependence in Taiwan. *Addict Biol*. 2005 Sep;10(3):289-92. PMID: 16109592.
55. de Sousa A, de Sousa A. An open randomized study comparing disulfiram and acamprosate in the treatment of alcohol dependence. *Alcohol Alcohol*. 2005 Nov-Dec;40(6):545-8. PMID: 16043433.
56. Anton RF, Moak DH, Latham P, et al. Naltrexone combined with either cognitive behavioral or motivational enhancement therapy for alcohol dependence. *J Clin Psychopharmacol*. 2005 Aug;25(4):349-57. PMID: 16012278.
57. Petrakis IL, Poling J, Levinson C, et al. Naltrexone and disulfiram in patients with alcohol dependence and comorbid psychiatric disorders. *Biol Psychiatry*. 2005 May 15;57(10):1128-37. PMID: 15866552.
58. Ralevski E, Ball S, Nich C, et al. The impact of personality disorders on alcohol-use outcomes in a pharmacotherapy trial for alcohol dependence and comorbid Axis I disorders. *Am J Addict*. 2007 Nov-Dec;16(6):443-9. PMID: 18058408.
59. Petrakis I, Ralevski E, Nich C, et al. Naltrexone and disulfiram in patients with alcohol dependence and current depression. *J Clin Psychopharmacol*. 2007 Apr;27(2):160-5. PMID: 17414239.
60. Petrakis IL, Poling J, Levinson C, et al. Naltrexone and disulfiram in patients with alcohol dependence and comorbid post-traumatic stress disorder. *Biol Psychiatry*. 2006 Oct 1;60(7):777-83. PMID: 17008146.
61. Garbutt JC, Kranzler HR, O'Malley SS, et al. Efficacy and tolerability of long-acting injectable naltrexone for alcohol dependence: a randomized controlled trial. *JAMA*. 2005 Apr 6;293(13):1617-25. PMID: 15811981.
62. Brady KT, Sonne S, Anton RF, et al. Sertraline in the treatment of co-occurring alcohol dependence and posttraumatic stress disorder. *Alcohol Clin Exp Res*. 2005 Mar;29(3):395-401. PMID: 15770115.
63. Salloum IM, Cornelius JR, Daley DC, et al. Efficacy of valproate maintenance in patients with bipolar disorder and alcoholism: a double-blind placebo-controlled study. *Arch Gen Psychiatry*. 2005 Jan;62(1):37-45. PMID: 15630071.
64. Killeen TK, Brady KT, Gold PB, et al. Effectiveness of naltrexone in a community treatment program. *Alcohol Clin Exp Res*. 2004 Nov;28(11):1710-7. PMID: 15547458.
65. De Sousa A, De Sousa A. A one-year pragmatic trial of naltrexone vs disulfiram in the treatment of alcohol dependence. *Alcohol Alcohol*. 2004 Nov-Dec;39(6):528-31. PMID: 15525790.
66. Schmitz JM, Stotts AL, Sayre SL, et al. Treatment of cocaine-alcohol dependence with naltrexone and relapse prevention therapy. *Am J Addict*. 2004 Jul-Sep;13(4):333-41. PMID: 15370932.
67. Johnson BA, Ait-Daoud N, Aubin HJ, et al. A pilot evaluation of the safety and tolerability of repeat dose administration of long-acting injectable naltrexone (Vivitrex) in patients with alcohol dependence. *Alcohol Clin Exp Res*. 2004 Sep;28(9):1356-61. PMID: 15365306.

68. Johnson BA, Ait-Daoud N, Akhtar FZ, et al. Oral topiramate reduces the consequences of drinking and improves the quality of life of alcohol-dependent individuals: a randomized controlled trial. *Arch Gen Psychiatry*. 2004 Sep;61(9):905-12. PMID: 15351769.
69. Kranzler HR, Wesson DR, Billot L. Naltrexone depot for treatment of alcohol dependence: a multicenter, randomized, placebo-controlled clinical trial. *Alcohol Clin Exp Res*. 2004 Jul;28(7):1051-9. PMID: 15252291.
70. Anton RF, Pettinati H, Zweben A, et al. A multi-site dose ranging study of nalmefene in the treatment of alcohol dependence. *J Clin Psychopharmacol*. 2004 Aug;24(4):421-8. PMID: 15232334.
71. Guardia J, Segura L, Gonzalvo B, et al. A double-blind, placebo-controlled study of olanzapine in the treatment of alcohol-dependence disorder. *Alcohol Clin Exp Res*. 2004 May;28(5):736-45. PMID: 15166648.
72. Chick J, Aschauer H, Hornik K. Efficacy of fluvoxamine in preventing relapse in alcohol dependence: a one-year, double-blind, placebo-controlled multicentre study with analysis by typology. *Drug Alcohol Depend*. 2004 Apr 9;74(1):61-70. PMID: 15072808.
73. Baltieri DA, De Andrade AG. Acamprosate in alcohol dependence: a randomized controlled efficacy study in a standard clinical setting. *J Stud Alcohol*. 2004 Jan;65(1):136-9. PMID: 15000513.
74. Petrakis IL, O'Malley S, Rounsaville B, et al. Naltrexone augmentation of neuroleptic treatment in alcohol abusing patients with schizophrenia. *Psychopharmacology (Berl)*. 2004 Mar;172(3):291-7. PMID: 14634716.
75. Ralevski E, Balachandra K, Gueorguieva R, et al. Effects of naltrexone on cognition in a treatment study of patients with schizophrenia and comorbid alcohol dependence. *J Dual Diagn*. 2006;2(4):53-69.
76. Gual A, Balcells M, Torres M, et al. Sertraline for the prevention of relapse in detoxicated alcohol dependent patients with a comorbid depressive disorder: a randomized controlled trial. *Alcohol Alcohol*. 2003 Nov-Dec;38(6):619-25. PMID: 14633652.
77. Moak DH, Anton RF, Latham PK, et al. Sertraline and cognitive behavioral therapy for depressed alcoholics: results of a placebo-controlled trial. *J Clin Psychopharmacol*. 2003 Dec;23(6):553-62. PMID: 14624185.
78. Balldin J, Berglund M, Borg S, et al. A 6-month controlled naltrexone study: combined effect with cognitive behavioral therapy in outpatient treatment of alcohol dependence. *Alcohol Clin Exp Res*. 2003 Jul;27(7):1142-9. PMID: 12878920.
79. Johnson BA, Ait-Daoud N, Bowden CL, et al. Oral topiramate for treatment of alcohol dependence: a randomised controlled trial. *Lancet*. 2003 May 17;361(9370):1677-85. PMID: 12767733.
80. Ma JZ, Ait-Daoud N, Johnson BA. Topiramate reduces the harm of excessive drinking: implications for public health and primary care. *Addiction*. 2006 Nov;101(11):1561-8. PMID: 17034435.
81. Kiefer F, Jahn H, Tarnaske T, et al. Comparing and combining naltrexone and acamprosate in relapse prevention of alcoholism: a double-blind, placebo-controlled study. *Arch Gen Psychiatry*. 2003 Jan;60(1):92-9. PMID: 12511176.
82. Kiefer F, Helwig H, Tarnaske T, et al. Pharmacological relapse prevention of alcoholism: clinical predictors of outcome. *Eur Addict Res*. 2005;11(2):83-91. PMID: 15785069.
83. Gastpar M, Bonnet U, Boning J, et al. Lack of efficacy of naltrexone in the prevention of alcohol relapse: results from a German multicenter study. *J Clin Psychopharmacol*. 2002 Dec;22(6):592-8. PMID: 12454559.
84. Guardia J, Caso C, Arias F, et al. A double-blind, placebo-controlled study of naltrexone in the treatment of alcohol-dependence disorder: results from a multicenter clinical trial. *Alcohol Clin Exp Res*. 2002 Sep;26(9):1381-7. PMID: 12351933.
85. Brady KT, Myrick H, Henderson S, et al. The use of divalproex in alcohol relapse prevention: a pilot study. *Drug Alcohol Depend*. 2002 Aug 1;67(3):323-30. PMID: 12127203.

86. Latt NC, Jurd S, Houseman J, et al. Naltrexone in alcohol dependence: a randomised controlled trial of effectiveness in a standard clinical setting. *Med J Aust*. 2002 Jun 3;176(11):530-4. PMID: 12064984.
87. Morris PL, Hopwood M, Whelan G, et al. Naltrexone for alcohol dependence: a randomized controlled trial. *Addiction*. 2001 Nov;96(11):1565-73. PMID: 11784454.
88. Krystal JH, Cramer JA, Krol WF, et al. Naltrexone in the treatment of alcohol dependence. *N Engl J Med*. 2001 Dec 13;345(24):1734-9. PMID: 11742047.
89. Monti PM, Rohsenow DJ, Swift RM, et al. Naltrexone and cue exposure with coping and communication skills training for alcoholics: treatment process and 1-year outcomes. *Alcohol Clin Exp Res*. 2001 Nov;25(11):1634-47. PMID: 11707638.
90. Rohsenow DJ, Miranda R, Jr., McGeary JE, et al. Family history and antisocial traits moderate naltrexone's effects on heavy drinking in alcoholics. *Exp Clin Psychopharmacol*. 2007 Jun;15(3):272-81. PMID: 17563214.
91. Rohsenow DJ, Colby SM, Monti PM, et al. Predictors of compliance with naltrexone among alcoholics. *Alcohol*. 2000;24(10):1542-9.
92. Monterosso JR, Flannery BA, Pettinati HM, et al. Predicting treatment response to naltrexone: the influence of craving and family history. *Am J Addict*. 2001 Summer;10(3):258-68. PMID: 11579624.
93. Rubio G, Jimenez-Arriero MA, Ponce G, et al. Naltrexone versus acamprosate: one year follow-up of alcohol dependence treatment. *Alcohol Alcohol*. 2001 Sep-Oct;36(5):419-25. PMID: 11524308.
94. Heinala P, Alho H, Kiianmaa K, et al. Targeted use of naltrexone without prior detoxification in the treatment of alcohol dependence: a factorial double-blind, placebo-controlled trial. *J Clin Psychopharmacol*. 2001 Jun;21(3):287-92. PMID: 11386491.
95. Pettinati HM, Volpicelli JR, Luck G, et al. Double-blind clinical trial of sertraline treatment for alcohol dependence. *J Clin Psychopharmacol*. 2001 Apr;21(2):143-53. PMID: 11270910.
96. Chick J, Anton R, Chercinski K, et al. A multicentre, randomized, double-blind, placebo-controlled trial of naltrexone in the treatment of alcohol dependence or abuse. *Alcohol Alcohol*. 2000 Nov-Dec;35(6):587-93. PMID: 11093966.
97. Fawcett J, Kravitz HM, McGuire M, et al. Pharmacological treatments for alcoholism: revisiting lithium and considering buspirone. *Alcohol Clin Exp Res*. 2000 May;24(5):666-74. PMID: 10832908.
98. Tempesta E, Janiri L, Bignamini A, et al. Acamprosate and relapse prevention in the treatment of alcohol dependence: a placebo-controlled study. *Alcohol Alcohol*. 2000 Mar-Apr;35(2):202-9. PMID: 10787398.
99. Chick J, Howlett H, Morgan MY, et al. United Kingdom Multicentre Acamprosate Study (UKMAS): a 6-month prospective study of acamprosate versus placebo in preventing relapse after withdrawal from alcohol. *Alcohol Alcohol*. 2000 Mar-Apr;35(2):176-87. PMID: 10787394.
100. Anton RF, Moak DH, Waid LR, et al. Naltrexone and cognitive behavioral therapy for the treatment of outpatient alcoholics: results of a placebo-controlled trial. *Am J Psychiatry*. 1999 Nov;156(11):1758-64. PMID: 10553740.
101. Anton RF, Moak DH, Latham PK, et al. Posttreatment results of combining naltrexone with cognitive-behavior therapy for the treatment of alcoholism. *J Clin Psychopharmacol*. 2001 Feb;21(1):72-7. PMID: 11199951.
102. George DT, Rawlings R, Eckardt MJ, et al. Buspirone treatment of alcoholism: age of onset, and cerebrospinal fluid 5-hydroxyindolacetic acid and homovanillic acid concentrations, but not medication treatment, predict return to drinking. *Alcohol Clin Exp Res*. 1999 Feb;23(2):272-8. PMID: 10069556.

103. Besson J, Aeby F, Kasas A, et al. Combined efficacy of acamprosate and disulfiram in the treatment of alcoholism: a controlled study. *Alcohol Clin Exp Res*. 1998 May;22(3):573-9. PMID: 9622434.
104. Poldrugo F. Acamprosate treatment in a long-term community-based alcohol rehabilitation programme. *Addiction*. 1997 Nov;92(11):1537-46. PMID: 9519495.
105. Oslin D, Liberto JG, O'Brien J, et al. Naltrexone as an adjunctive treatment for older patients with alcohol dependence. *Am J Geriatr Psychiatry*. 1997 Fall;5(4):324-32. PMID: 9363289.
106. Volpicelli JR, Rhines KC, Rhines JS, et al. Naltrexone and alcohol dependence. Role of subject compliance. *Arch Gen Psychiatry*. 1997 Aug;54(8):737-42. PMID: 9283509.
107. Cornelius JR, Salloum IM, Ehler JG, et al. Fluoxetine in depressed alcoholics. A double-blind, placebo-controlled trial. *Arch Gen Psychiatry*. 1997 Aug;54(8):700-5. PMID: 9283504.
108. Cornelius JR, Salloum IM, Ehler JG, et al. Double-blind fluoxetine in depressed alcoholic smokers. *Psychopharmacol Bull*. 1997;33(1):165-70. PMID: 9133770.
109. Pelc I, Verbanck P, Le Bon O, et al. Efficacy and safety of acamprosate in the treatment of detoxified alcohol-dependent patients. A 90-day placebo-controlled dose-finding study. *Br J Psychiatry*. 1997 Jul;171:73-7. PMID: 9328500.
110. Sass H, Soyka M, Mann K, et al. Relapse prevention by acamprosate. Results from a placebo-controlled study on alcohol dependence. *Arch Gen Psychiatry*. 1996 Aug;53(8):673-80. PMID: 8694680.
111. Kabel DI, Petty F. A placebo-controlled, double-blind study of fluoxetine in severe alcohol dependence: adjunctive pharmacotherapy during and after inpatient treatment. *Alcohol Clin Exp Res*. 1996 Jun;20(4):780-4. PMID: 8800399.
112. Whitworth AB, Fischer F, Lesch OM, et al. Comparison of acamprosate and placebo in long-term treatment of alcohol dependence. *Lancet*. 1996 May 25;347(9013):1438-42. PMID: 8676626.
113. Malec E, Malec T, Gagne MA, et al. Buspirone in the treatment of alcohol dependence: a placebo-controlled trial. *Alcohol Clin Exp Res*. 1996 Apr;20(2):307-12. PMID: 8730222.
114. Mason BJ, Kocsis JH, Ritvo EC, et al. A double-blind, placebo-controlled trial of desipramine for primary alcohol dependence stratified on the presence or absence of major depression. *JAMA*. 1996 Mar 13;275(10):761-7. PMID: 8598592.
115. McGrath PJ, Nunes EV, Stewart JW, et al. Imipramine treatment of alcoholics with primary depression: A placebo-controlled clinical trial. *Arch Gen Psychiatry*. 1996 Mar;53(3):232-40. PMID: 8611060.
116. Tihihonen J, Ryyanen OP, Kauhanen J, et al. Citalopram in the treatment of alcoholism: a double-blind placebo-controlled study. *Pharmacopsychiatry*. 1996 Jan;29(1):27-9. PMID: 8852531.
117. Kranzler HR, Burleson JA, Korner P, et al. Placebo-controlled trial of fluoxetine as an adjunct to relapse prevention in alcoholics. *Am J Psychiatry*. 1995 Mar;152(3):391-7. PMID: 7864265.
118. Paille FM, Guelfi JD, Perkins AC, et al. Double-blind randomized multicentre trial of acamprosate in maintaining abstinence from alcohol. *Alcohol Alcohol*. 1995 Mar;30(2):239-47. PMID: 7662044.
119. Naranjo CA, Bremner KE, Lanctot KL. Effects of citalopram and a brief psychosocial intervention on alcohol intake, dependence and problems. *Addiction*. 1995 Jan;90(1):87-99. PMID: 7888983.
120. Volpicelli JR, Clay KL, Watson NT, et al. Naltrexone in the treatment of alcoholism: predicting response to naltrexone. *J Clin Psychiatry*. 1995;56 Suppl 7:39-44. PMID: 7673104.
121. Kranzler HR, Burleson JA, Del Boca FK, et al. Buspirone treatment of anxious alcoholics. A placebo-controlled trial. *Arch Gen Psychiatry*. 1994 Sep;51(9):720-31. PMID: 8080349.
122. Malcolm R, Anton RF, Randall CL, et al. A placebo-controlled trial of buspirone in anxious inpatient alcoholics. *Alcohol Clin Exp Res*. 1992 Dec;16(6):1007-13. PMID: 1335217.

123. O'Malley SS, Jaffe AJ, Chang G, et al. Naltrexone and coping skills therapy for alcohol dependence. A controlled study. *Arch Gen Psychiatry*. 1992 Nov;49(11):881-7. PMID: 1444726.
124. O'Malley SS, Jaffe AJ, Chang G, et al. Six-month follow-up of naltrexone and psychotherapy for alcohol dependence. *Arch Gen Psychiatry*. 1996 Mar;53(3):217-24. PMID: 8611058.
125. Lhuintre JP, Moore N, Tran G, et al. Acamprosate appears to decrease alcohol intake in weaned alcoholics. *Alcohol Alcohol*. 1990;25(6):613-22. PMID: 2085344.
126. Fuller RK, Branchey L, Brightwell DR, et al. Disulfiram treatment of alcoholism. A Veterans Administration cooperative study. *JAMA*. 1986 Sep 19;256(11):1449-55. PMID: 3528541.
127. Lhuintre JP, Daoust M, Moore ND, et al. Ability of calcium bis acetyl homotaurine, a GABA agonist, to prevent relapse in weaned alcoholics. *Lancet*. 1985 May 4;1(8436):1014-6. PMID: 2859465.
128. Ling W, Weiss DG, Charuvastra VC, et al. Use of disulfiram for alcoholics in methadone maintenance programs. A Veterans Administration Cooperative Study. *Arch Gen Psychiatry*. 1983 Aug;40(8):851-4. PMID: 6347118.
129. Fuller RK, Roth HP. Disulfiram for the treatment of alcoholism. An evaluation in 128 men. *Ann Intern Med*. 1979 Jun;90(6):901-4. PMID: 389121.
130. Gual A, Leher P. Acamprosate during and after acute alcohol withdrawal: a double-blind placebo-controlled study in Spain. *Alcohol and alcoholism (Oxford, Oxfordshire)*; 2001. p. 413-8.
131. Coskunol H, Gökden O, Ercan ES, et al. Long-term efficacy of sertraline in the prevention of alcoholic relapses in alcohol-dependent patients: a single-center, double-blind, randomized, placebo-controlled, parallel-group study. *Current Therapeutic Research*. 2002;63(11):759-71. PMID: 2003140542.
132. Kranzler HR, Armeli S, Covault J, et al. Variation in OPRM1 moderates the effect of desire to drink on subsequent drinking and its attenuation by naltrexone treatment. *Addiction Biology*. 2013;18(1):193-201.
133. Ahmadi J, Ahmadi N. A double blind, placebo-controlled study of naltrexone in the treatment of alcohol dependence. *German Journal of Psychiatry*. 2002;5(4):85-9.
134. Ahmadi J, Babaebeigi M, Maany I, et al. Naltrexone for alcohol-dependent patients. *Ir J Med Sci*. 2004 Jan-Mar;173(1):34-7. PMID: 15732235.
135. Mason BJ, Salvato FR, Williams LD, et al. A double-blind, placebo-controlled study of oral nalmefene for alcohol dependence. *Arch Gen Psychiatry*. 1999;56(8):719-24.
136. Geerlings PJ, Ansoms C, Van Den Brink W. Acamprosate and prevention of relapse in alcoholics. Results of a randomized, placebo-controlled, double-blind study in out-patient alcoholics in the Netherlands, Belgium and Luxembourg. *Eur Addict Res*. 1997;3(3):129-37.
137. Mason BJ, Ritvo EC, Morgan RO, et al. A double-blind, placebo-controlled pilot study to evaluate the efficacy and safety of oral nalmefene HCl for alcohol dependence. *Alcohol*. 1994;18(5):1162-7.
138. Tollefson GD, Lancaster SP, Montague-Clouse J. The association of buspirone and its metabolite 1-pyrimidinylpiperazine in the remission of comorbid anxiety with depressive features and alcohol dependency. *Psychopharmacol Bull*. 1991;27(2):163-70.
139. Tollefson GD, Montague-Clouse J, Tollefson SL. Treatment of comorbid generalized anxiety in a recently detoxified alcoholic population with a selective serotonergic drug (buspirone). *J Clin Psychopharmacol*. 1992 Feb;12(1):19-26. PMID: 1552035.
140. Lee A, Tan S, Lim D, et al. Naltrexone in the treatment of male alcoholics—An effectiveness study In Singapore. *Drug and Alcohol Review*. 2001;20(2):193-9.
141. Carroll K, Ziedonis D, O'Malley SS, et al. Pharmacologic interventions for alcohol- and cocaine-abusing individuals: A pilot study of disulfiram vs. naltrexone. *Am J Addict*. 1993;2(1):77-9.

142. Morgenstern J, Kuerbis AN, Chen AC, et al. A randomized clinical trial of naltrexone and behavioral therapy for problem drinking men who have sex with men. *J Consult Clin Psychol.* 2012;80(5):863-75. PMID: 22612306.
143. Pelc I, Le Bon O, Lehert P, et al. Acamprosate in the Treatment of Alcohol Dependence: A 6-Month Postdetoxification Study. In: Soyka M, ed. *Acamprosate in Relapse Prevention of Alcoholism.* Springer Berlin Heidelberg; 1996:133-42.
144. Pelc I, Le Bon O, Verbanck P, et al. Calciumacetylhomotaurinate for maintaining abstinence in weaned alcoholic patients: a placebo-controlled double-blind multi-centre study. In: Naranjo CA, Sellers EM, eds. *Novel Pharmacological Interventions for Alcoholism.* New York: Springer-Verlag; 1992.
145. Collier JK, Cahill S, Edmonds C, et al. OPRM1 A118G genotype fails to predict the effectiveness of naltrexone treatment for alcohol dependence. *Pharmacogenet Genomics.* 2011 Dec;21(12):902-5. PMID: 21946895.
146. Kim SG, Kim CM, Choi SW, et al. A mu opioid receptor gene polymorphism (A118G) and naltrexone treatment response in adherent Korean alcohol-dependent patients. *Psychopharmacology (Berl).* 2009;201(4):611-8.
147. Narayana PL, Gupta AK, Sharma PK. Use of anti-craving agents in soldiers with alcohol dependence syndrome. *Medical Journal Armed Forces India.* 2008;64(4):320-4.
148. Mutschler J, Abbruzzese E, Witt SH, et al. Functional polymorphism of the dopamine β -hydroxylase gene is associated with increased risk of disulfiram-induced adverse effects in alcohol-dependent patients. *J Clin Psychopharmacol.* 2012;32(4):578-80. PMID: 22760354.
149. Mason BJ, Lehert P. Acamprosate for alcohol dependence: a sex-specific meta-analysis based on individual patient data. *Alcohol Clin Exp Res.* 2012 Mar;36(3):497-508. PMID: 21895717.
150. Jorgensen CH, Pedersen B, Tonnesen H. The efficacy of disulfiram for the treatment of alcohol use disorder. *Alcohol Clin Exp Res.* 2011 Oct;35(10):1749-58. PMID: 21615426.
151. Rosner S, Hackl-Herrwerth A, Leucht S, et al. Acamprosate for alcohol dependence. *Cochrane Database Syst Rev.* 2010(9):CD004332. PMID: 20824837.
152. Rösner S, Hackl-Herrwerth A, Leucht S, et al. Opioid antagonists for alcohol dependence. *Cochrane Database Syst Rev:* John Wiley & Sons, Ltd; 2010.
153. Lobmaier P, Kornor H, Kunoe N, et al. Sustained-Release Naltrexone For Opioid Dependence. *Cochrane Database Syst Rev:* John Wiley & Sons, Ltd; 2008.
154. National Collaborating Centre for Mental Health. Alcohol-Use Disorders. Diagnosis, Assessment and Management of Harmful Drinking and Alcohol Dependence. National Clinical Practice Guideline 115. National Institute for Health & Clinical Excellence. London: Psychiatrists TBPSaTRCo; 2011. <http://guidance.nice.org.uk/CG115/Guidance/pdf/English>.

Appendix D. Strength of Evidence Assessments

KQ 1 and KQ 2

Table D-1. Acamprosate compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	15 ^a ; 4,747	Medium; RCTs	Consistent ^b	Direct	Precise	RD: -0.10 (-0.15 to -0.05)	Moderate
Return to heavy drinking	6; 2,239	Low; RCTs	Consistent	Direct	Precise	RD: -0.01 (-0.05 to 0.03)	Moderate ^c
Drinking days	12 ^d ; 4,385	Medium; RCTs	Consistent	Direct	Precise	WMD: -9.36 (-13.75 to -4.96)	Moderate
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	1 ^d ; 116	Low; RCT	Unknown	Direct	Imprecise	WMD: 0.40 (-1.81 to 2.61)	Insufficient
Accidents	0 ^e ; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	1; 612	Low; RCT	Unknown	Direct	Unknown	NSD ^f	Insufficient
Mortality	7 ^g ; 2,477	Medium; RCTs	Unknown	Direct	Imprecise	7 (ACA) vs. 5 (PBO)	Insufficient

^a 2 additional studies were rated high risk of bias; 1 additional study was rated as unclear risk of bias

^b Although there was considerable statistical heterogeneity, fourteen of fifteen studies reported point estimates that favored acamprosate; differences were in magnitude of benefit

^c The relatively small number of studies reporting this outcome raises concern for potential reporting bias, hence the rating of moderate rather than high rating

^d 1 additional study was rated high risk of bias

^e The single study that reported this outcome was rated as unclear risk of bias. It reported that one patient in the placebo group died by “accident.” No other details on the cause or nature of the accident were provided.¹

^f Results were not reported for each treatment group separately, but there were no clinically significant differences across treatment groups

^g One additional study reported a death but did not specify in which treatment group it occurred.²

Abbreviations: ACA = acamprosate; CI = confidence interval; NA = not applicable; NSD = no statistically significant difference; PBO = placebo; RCT = randomized controlled trial; RD = risk difference; WMD = weighted mean difference

Table D-2. Disulfiram compared with control

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	2 ^a ; 492	Medium; RCTs	Consistent ^b	Direct	Imprecise	RD: 0.04 (-0.03 to 0.11)	Low
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	2; 290	Medium; RCTs	Inconsistent	Indirect ^c	Imprecise	1 study reported similar percentages and no significant difference; the other reported that DIS was favored among the subset of subjects who drank and had a complete set of assessment interviews (N=162/605 subjects), P=0.05	Insufficient
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a 1 additional study was rated high risk of bias.³

^b Inclusion of the study rated high risk of bias would have made this inconsistent, though it would not have changed the conclusion (the meta-analysis still found no statistically significant difference between groups).

^c We considered this indirect because the larger study did not report the outcome for the randomized sample; it only reported this outcome for the subset (162/605) who drank and who had a complete set of assessment interviews.

Abbreviations: CI = confidence interval; NA = not applicable; RCT = randomized controlled trial; RD = risk difference

Table D-3. Naltrexone compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	21 ^a ; 4,232	Medium; RCTs	Consistent	Direct	Precise	RD: -0.04 (-0.07 to -0.01)	Moderate
Return to heavy drinking	21 ^a ; 3,794	Medium; RCTs	Consistent	Direct	Precise	RD: -0.08 (-0.12 to -0.04)	Moderate
Drinking days	19 ^b ; 3,329	Medium; RCTs	Consistent	Direct	Precise	WMD: -4.57 (-6.61 to -2.53)	Moderate
Heavy drinking days	10 ^c ; 1,423	Medium; RCTs	Consistent	Direct	Precise	WMD: -3.62 (-5.86 to -1.38)	Moderate
Drinks per drinking day	11 ^d ; 1,422	Medium; RCTs	Consistent	Direct	Imprecise	WMD: -0.54 (-1.01 to -0.07)	Low
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life	4; 1,513	Medium; RCTs	Inconsistent	Direct	Imprecise	Unable to pool data, some conflicting results ^e	Insufficient
Mortality	6 ^f ; 1,738	Medium; RCTs	Unknown	Direct	Imprecise	1 (NTX) vs. 2 (PBO)	Insufficient

^a 2 additional studies were rated high risk of bias; 2 additional studies were rated as unclear risk of bias

^b 3 additional studies were rated high risk of bias

^c 2 additional studies were rated high risk of bias

^d 5 additional studies were rated high risk of bias

^e Two studies found no significant difference between naltrexone- and placebo-treated subjects.^{4,5} One study reported that patients receiving injectable naltrexone 380mg/day had greater improvement on the mental health summary score than those receiving placebo at 24 weeks (8.2 vs. 6.2, $p=0.044$).^{6,210} One study measured alcohol-related consequences (with the DrInC) and reported that more subjects who received placebo (N=34) had ≥ 1 alcohol-related consequence than those who received naltrexone (N=34): 76% vs. 45%, $P=0.02$.⁷

^f One additional study reported a death but did not specify in which treatment group it occurred.²

Abbreviations: CI, confidence interval; NA, not applicable; NSD, no significant difference; NTX, naltrexone; PBO, placebo; RCT, randomized controlled trial; RD, risk difference; WMD, weighted mean difference

Table D-4. Acamprosate compared with disulfiram

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient

^a The one study reporting this outcome was rated high risk of bias.⁸ It reported one traffic accident in the disulfiram group and none in the acamprosate group over 52 weeks.⁸ No details of the event were described, it was noted that the study coordinator determined that the event was not related to the study treatment. One person committed suicide and two persons drowned in the acamprosate group but there were no events in the disulfiram group. Quality of life improved for both groups over the 52 week follow-up compared with baseline with no difference between the acamprosate and disulfiram groups.⁸

Abbreviations: CI, confidence interval; NA, not applicable

Table D-5. Acamprosate compared with naltrexone

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	3; 800	Low; RCTs	Consistent	Direct	Imprecise	RD: 0.02 (-0.03 to 0.08) ^a	Moderate
Return to heavy drinking	3; 800	Low; RCTs	Consistent	Direct	Imprecise	RD: 0.01 (-0.06 to 0.07) ^a	Moderate
Drinking days	2; 720	Low; RCTs	Inconsistent	Direct	Imprecise	WMD: -2.98 (-13.42 to 7.45) ^a	Low
Heavy drinking days	1; 612	Low; RCT	Unknown	Direct	Unknown	Significant NTX by CBI interaction, P=0.006	Insufficient
Drinks per drinking day	2; 720	Low; RCTs	Inconsistent	Direct	Unknown	Unable to pool data ^b	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	1 ^c ; 612	Low; RCT	Unknown	Direct	Imprecise	NSD for all measures except SF-12v2 physical health, which favored NTX+CBI	Insufficient
Mortality	0 ^d ; 0	NA	NA	NA	NA	NA	Insufficient

^a Positive value indicates that naltrexone is favored

^b Two trials reported some information about drinks per drinking day, but not enough data for us to conduct quantitative synthesis. One trial conducted in Australia reported no statistically significant difference between acamprosate and naltrexone (mean, SD: 7.5, 6.1 vs. 5.9, 6.1; P not reported).^{9,10} The COMBINE study reported that analyses of alternative summary measures of drinking, including drinks per drinking day (P=0.03), were consistent with those for the co-primary end points (percent days abstinent from alcohol and time to first heavy drinking day), all showing a significant naltrexone by CBI interaction.²

^c One additional study was rated high risk of bias.⁸ It found that quality of life improved for both groups over the 52 week follow-up compared with baseline, but found no difference between the acamprosate and naltrexone groups.

^d One study that reported this outcome was rated high risk of bias; another reported one death but did not specify in which treatment group it occurred

Abbreviations: ACA = acamprosate; CBI = combined behavioral intervention; CI = confidence interval; NA = not applicable; NTX = naltrexone; RCT = randomized controlled trial; RD = risk difference; WMD = weighted mean difference

Table D-6. Disulfiram compared with naltrexone

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0 ^b ; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0 ^b ; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0 ^c ; 0	NA	NA	NA	NA	NA	Insufficient

^a The single study that reported this outcome was rated high risk of bias.³ The trial reported no statistically significant difference between disulfiram and naltrexone for number of subjects achieving total abstinence (51 vs. 38, $P=0.11$), the percentage of days abstinent (96.6 vs. 95.4, $P=0.55$), or the percentage of heavy drinking days (3.2 vs. 4, $P=0.65$).

^b The only study that reported this outcome was rated high risk of bias.⁸ It reported one traffic accident in the disulfiram group and no accident or injuries in the naltrexone group. No details of the event were described, it was noted that the study coordinator determined that the event was not related to the study treatment. Quality of life improved for both groups over the 52 week follow-up compared with baseline with no difference between the disulfiram and naltrexone groups.

^c The only study that reported this outcome was rated high risk of bias.³ One person died in the naltrexone group and no deaths were reported in the disulfiram group.

Abbreviations: CI = confidence interval; NA = not applicable; RCT = randomized controlled trial

Table D-7. Amitriptyline compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

Abbreviations: CI, confidence interval; NA, not applicable

Table D-8. Aripiprazole compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size	Strength of Evidence Grade
Return to any drinking	1; 288	Medium; RCT	Unknown	Direct	Imprecise	89% (ARI) vs. 78% (PBO); P=0.02	Insufficient
Return to heavy drinking	1; 288	Medium; RCT	Unknown	Direct	Imprecise	73% (ARI) vs. 73% (PBO); P=0.98	Insufficient
Drinking days	1; 288	Medium; RCT	Unknown	Direct	Imprecise	41% (ARI) vs. 37% (PBO); P=0.23	Insufficient
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	1; 288	Medium; RCT	Unknown	Direct	Imprecise	4.4 (ARI) vs. 5.5 (PBO); P<0.001	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

Abbreviations: ARI, aripiprazole; NA, not applicable; PBO, placebo; RCT, randomized controlled trial

Table D-9. Atomoxetine compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^aThe single study reporting this outcome was rated high risk of bias.¹¹ The trial found no significant difference between groups for return to heavy drinking (94% for atomoxetine vs. 96% for placebo), drinking days, or reduction in drinks per drinking day. It did report a 26% lower rate of cumulative heavy drinking days for atomoxetine compared with placebo (P=0.02).

Abbreviations: ATO = atomoxetine; CI = confidence interval; HDD = heavy drinking days; NA = not applicable; PBO = placebo; RCT = randomized controlled trial

Table D-10. Baclofen compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size	Strength of Evidence Grade
Return to any drinking	2; 164	Medium; RCTs	Inconsistent	Direct	Imprecise	Study 1: OR 6.3, 95% CI 2.4, 16.1 Study 2: No difference ^a	Insufficient
Return to heavy drinking	2; 164	Medium; RCTs	Inconsistent	Direct	Imprecise	Study 1: BAC significantly lower than PBO (data in Figure, P=0.0062) Study 2: HR 0.924, P=0.76	Insufficient
Drinking days	1; 80	Medium; RCT	Unknown	Direct	Imprecise	50.1% (BAC) vs. 49.4% (PBO); P=0.50	Insufficient
Heavy drinking days	1; 80	Medium; RCT	Unknown	Direct	Imprecise	25.9% (BAC) vs. 25.5% (PBO); P=0.73	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a One trial conducted in Italy (N=84) reported that a lower percentage of patients treated with baclofen returned to any drinking than with placebo (29% [12/42] vs. 71% [30/42], OR 6.3, 95% CI 2.4, 16.1).¹² One trial conducted in the U.S. (N=80) did not report numbers for rates of return to any drinking, but reported no difference between groups for time to first usage (P = 0.13), and included a figure for percentage abstinent that shows over 90 percent of subjects returned to any drinking over the course of the trial.¹³

Abbreviations: BAC = baclofen; CI = confidence interval; NA = not applicable; PBO = placebo; RCT = randomized controlled trial

Table D-11. Buspirone compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	1 ^a ; 54	Medium; RCT	Unknown	Direct	Imprecise	RD: 0.07 (-0.19 to 0.34)	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	2; 161	Medium; RCTs	Consistent	Direct	Imprecise	WMD: -3.39 (-9.23 to 2.44)	Low
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	1; 61	Medium; RCT	Unknown	Direct	Imprecise	0.7 vs. 2.1; P NS	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a1 additional study was rated high risk of bias.¹⁴ Neither trial found a statistically significant difference between groups, and point estimates favored placebo in both trials.^{14,15}

Abbreviations: CI, confidence interval; NA, not applicable; RCT, randomized controlled trial; RD, risk difference; weighted mean difference

Table D-12. Citalopram compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0 ^b ; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0 ^b ; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a One trial conducted in Finland, rated as high risk of bias, reported 25 of 31 citalopram-treated patients and 28 of 31 placebo-treated patients returned to any drinking (P=0.10).¹⁶

^b One trial rated as high risk of bias conducted in Canada found similar proportions of drinking days for those who received citalopram and those who received placebo (72.7% vs. 76.5%, P NS) and similar reductions in drinks per drinking day for those who received citalopram and those who received placebo (26.1% vs. 26.4%, P NS) over the 12 weeks of treatment.¹⁷

Abbreviations: CI, confidence interval; NA, not applicable; RCT, randomized controlled trial

Table D-13. Desipramine compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a One study rated as high risk of bias, reported that 12% of patients in the desipramine arm returned to heavy drinking, compared with 32 percent of patients taking placebo (P=NS).¹⁸ It also reported that non-depressed patients treated with desipramine (N=14) drank on a median of 68 percent of days; non-depressed patients treated with placebo (N=15) drank on 72 percent of days (P NS).

Abbreviations: CI = confidence interval; NA = not applicable

Table D-14. Fluoxetine compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	1 ^a ; 51	Medium; RCT	Unknown	Direct	Imprecise	RD: -0.13 (-0.35 to 0.10)	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	2; 146	Medium; RCTs	Inconsistent	Direct	Imprecise	WMD: -3.15 (-18.2 to 11.9)	Low
Heavy drinking days	1; 51	Medium; RCT	Unknown	Direct	Imprecise	4.8 (FLUOX) vs. 16 (PBO); P=0.04	Insufficient
Drinks per drinking day	2; 146	Medium; RCTs	Inconsistent	Direct	Imprecise	WMD: -1.20 (-4.63 to 2.23)	Low
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a 1 additional study (N=28) reporting this outcome was rated high risk of bias.¹⁹ Both trials found no statistically significant difference between fluoxetine and placebo.^{19,20}

Abbreviations: CI = confidence interval; FLUOX = fluoxetine; NA = not applicable; PBO = placebo; RCT = randomized controlled trial; RD = risk difference; WMD = weighted mean difference

Table D-15. Fluvoxamine compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking ^a	1; 492	Medium; RCT	Unknown	Direct	Imprecise	12 weeks: 58% (FLUV) vs. 54% (PBO); P=0.40 52 weeks: 71% (FLUV) vs. 71% (PBO); P=0.94	Insufficient
Return to heavy drinking	1; 492	Medium; RCT	Unknown	Direct	Imprecise	12 weeks: 46% (FLUV) vs. 40% (PBO); P=0.18 52 weeks: 64% (FLUV) vs. 64% (PBO); P=0.47	Insufficient
Drinking days ^a	1; 492	Medium; RCT	Unknown	Direct	Imprecise	12 weeks: 31% (FLUV) vs. 23% (PBO); P=0.009 52 weeks: 44% (FLUV) vs. 38% (PBO); P=0.13	Insufficient
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	1; 492	Medium; RCT	Unknown	Direct	Imprecise	52 weeks: 1 (FLUV) vs. 1 (PBO)	Insufficient

^aThe study reported return to drinking and percent drinking days since the previous assessment. At 12 weeks, the previous assessment was at week 8; at 52 weeks, the previous assessment was at week 40.²¹

Abbreviations: CI = confidence interval; FLUV = fluvoxamine; NA = not applicable; PBO = placebo; RCT = randomized controlled trial

Table D-16. Imipramine compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking ^a	1; 56	Medium; RCT	Unknown	Direct	Imprecise	69% (IMI) vs. 79% (PBO); P NS	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	1; 56	Medium; RCT	Unknown	Direct	Imprecise	28.3% (IMI) vs. 30.8% (PBO); P NS	Insufficient
Heavy drinking days	1; 56	Medium; RCT	Unknown	Direct	Imprecise	13.5% (IMI) vs. 9.0% (PBO); P NS	Insufficient
Drinks per drinking day	1; 56	Medium; RCT	Unknown	Direct	Imprecise	3.7 (IMI) vs. 4.1 (PBO); P NS	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^aThe study reported return to drinking within the previous 4 weeks.²²

Abbreviations: CI = confidence interval; IMI = imipramine; NA = not applicable; NS = not significant; PBO = placebo; RCT = randomized controlled trial

Table D-17. Nalmefene compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	1 ^a ; 105	Medium; RCT	Unknown	Direct	Imprecise	RD: -0.22 (-0.42 to -0.02) ^a	Insufficient
Drinking days	2; 508	Medium; RCTs	Inconsistent	Direct	Imprecise	WMD: -1.1 (-7.6 to 5.4)	Low
Heavy drinking days	1; 403	Medium; RCT	Unknown	Direct	Imprecise	18.1% (NALM) vs. 29.7% (PBO); P=0.024	Insufficient
Drinks per drinking day	3; 608	Medium; RCTs	Consistent	Direct	Precise	WMD: -1.02 (-1.77 to -0.28)	Moderate
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a 1 additional study reported return to heavy drinking—a pilot study rated as high risk of bias (N=21).²³ It found no difference between groups (RD -0.05, 95% CI -0.51, 0.41). Pooling both studies found a 19% reduction in return to heavy drinking with nalmefene (RD -0.19, 95% CI -0.37, -0.01).

Abbreviations: CI = confidence interval; NA = not applicable; NALM = nalmefene; PBO = placebo; RCT = randomized controlled trial; RD = risk difference; WMD = weighted mean difference

Table D-18. Olanzapine compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	1; 60	Medium; RCT	Unknown	Direct	Imprecise	37.9% (OLA) vs. 29.0% (PBO); P=0.50 ²⁴	Insufficient
Drinking days	1; 60	Medium; RCT	Unknown	Direct	Imprecise	13.1% (OLA) vs. 22.7% (PBO); P=0.18 ²⁴	Insufficient
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	1; 60	Medium; RCT	Unknown	Direct	Imprecise	1.79 (OLA) vs. 2.02 (PBO); P=0.71 ²⁴	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

Abbreviations: CI = confidence interval; NA = not applicable; OLA = olanzapine; PBO = placebo; RCT = randomized controlled trial

Table D-19. Paroxetine compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	1; 42	Medium; RCT	Unknown	Direct	Imprecise	34% (PAR) vs. 35% (PBO); P=NS ^{25,26}	Insufficient
Heavy drinking days ^a	1; 42	Medium; RCT	Unknown	Direct	Imprecise	54% (PAR) vs. 55% (PBO); P=NS ^{25,26}	Insufficient
Drinks per drinking day	1; 42	Medium; RCT	Unknown	Direct	Imprecise	5.9 (PAR) vs. 7.0 (PBO); P=NS ^{25,26}	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a These results indicate the percent of heavy drinking days within the number of any drinking days.

Abbreviations: CI = confidence interval; NA = not applicable; PAR = paroxetine; PBO = placebo; RCT = randomized controlled trial

Table D-20. Quetiapine compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0 ^b ; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0 ^b ; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0 ^c ; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0 ^c ; 0	NA	NA	NA	NA	NA	Insufficient

^a The one study that reported this outcome was rated as high risk of bias.²⁷ It did not enroll subjects with co-occurring bipolar disorder and it reported that more subjects treated with quetiapine achieved complete abstinence (9/29 vs. 2/32, $P=0.012$)—i.e., fewer subjects treated with quetiapine returned to any drinking (20/29 vs. 30/32).

^b Three studies reported this outcome; all three were rated as high risk of bias.²⁷⁻²⁹ Our meta-analysis of the three trials found no difference in drinking days between patients treated with quetiapine and those who received placebo (WMD -2.7 95% CI -12.8, 7.5). Our meta-analysis of the three trials found no difference in heavy drinking days between patients treated with quetiapine and those who received placebo (WMD -3.1 95% CI -10.1, 4.0).

^c One placebo-controlled trial of quetiapine rated as high risk of bias reported two deaths (one in each treatment group); one after a skull fracture caused by blunt trauma in the quetiapine group and one attributed to myocardial ischemia more than 30 days after treatment in the placebo group.²⁹ Both deaths were judged to be unrelated to the study medications by the study investigators. The trial also reported no difference between groups for quality of life and function.²⁹

Abbreviations: CI = confidence interval; NA = not applicable

Table D-21. Sertraline compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	1; 79	Medium; RCT	Unknown	Direct	Imprecise	29/40 (SER) vs. 30/39 (PBO), P NS.	Insufficient
Return to heavy drinking	2; 142	Medium; RCTs	Inconsistent	Direct	Imprecise	RD: -0.04 (-0.31 to 0.23)	Low
Drinking days	2 ^a ; 165	Medium; RCTs	Consistent	Direct	Imprecise	WMD: 0.03 (-11.0 to 11.1) ^a	Low
Heavy drinking days	1; 94	Medium; RCT	Unknown	Direct	Imprecise	10.4 (SER) vs. 8.9 (PBO); P NS ³⁰	Insufficient
Drinks per drinking day	2; 176	Medium; RCTs	Consistent	Direct	Imprecise	WMD: -0.9 (-2.2 to 0.5)	Low
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	1; 83	Medium; RCT	Unknown	Direct	Imprecise	Graph only, data NR (P=0.031) ^b	Insufficient
Mortality	1; 79	Medium; RCT	Unknown	Direct	Imprecise	0 (SER) vs. 0 (PBO) ³¹	Insufficient

^a One additional study reporting this outcome was rated as unclear risk of bias.³² Our meta-analysis found no significant difference between patients treated with sertraline and those who received placebo, both without and with including the trial rated as unclear risk of bias (when including that trial: WMD -0.7, 95% CI -8.2, 6.9).

^b The single study that reported this outcome enrolled patients with co-existing depression and measured QoL using the SF-36 at 24 weeks. Scores were presented in a figure only (bar graph, data not reported). QoL improved during treatment for both the placebo and sertraline groups; the authors noted that the sertraline group improved more than placebo in only the mental health summary score of the SF-36 (p=0.031).³³

Abbreviations: CI = confidence interval; NA = not applicable; PBO = placebo; RCT = randomized controlled trial; RD = risk difference; SER = sertraline; WMD = weighted mean difference

Table D-22. Topiramate compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	2 ^b ; 521	Low; RCTs	Consistent	Direct	Imprecise	Trial 1 ³⁴ : WMD: -8.5 (-15.9 to -1.1) ^b Trial 2 ³⁵ : mean difference -11.6 (-3.98 to -19.3)	Moderate ^b
Heavy drinking days	2 ^b ; 521	Low; RCTs	Consistent	Direct	Imprecise	WMD: -11.53 (-18.29 to -4.77)	Moderate ^b
Drinks per drinking day	2 ^b ; 521	Low; RCT	Consistent	Direct	Imprecise	WMD: -1.10 (-1.75 to -0.45)	Moderate ^b
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	1; 371	Low; RCT	Unknown	Direct	Imprecise	4.4% (TOP) vs. 11.7% (PBO); P=0.01 ³⁴	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	1; 371	Low; RCT	Unknown	Direct	Imprecise	0 (TOP) vs. 1 (PBO) ³⁴	Insufficient

^a One study conducted in Brazil, rated as high risk of bias, reported this outcome.³⁶ It reported that more patients treated with topiramate returned to any drinking than with placebo (24/52 vs. 15/54).

^b One additional study reporting this outcome was rated as high risk of bias.³⁷ Our meta-analysis found a lower percentage of drinking days for patients treated with topiramate than for those who received placebo both without and with including the trial rated as high risk of bias (WMD -9.7, 95% CI -16.4, -3.1). Our meta-analysis found a lower percentage of heavy drinking days for patients treated with topiramate than for those who received placebo both without and with including the trial rated as high risk of bias (WMD -11.4, 95% CI -20.4, -2.4). Our meta-analysis found no statistically significant difference between topiramate and placebo when only including the trial rated as low risk of bias, but found a statistically significant reduction of 1.2 drinks per drinking day when including the trial rated as high risk of bias (WMD -1.2, 95% CI -2.2, -0.2). We were unable to include “trial 2” (N=150),³⁵ rated as medium risk of bias, in our meta-analyses due to differences in the type of data reported, but its findings are shown in the SOE table, and were generally consistent with those of the low risk of bias trial (“trial 1”, N=371).³⁴

Abbreviations: CI = confidence interval; NA = not applicable; PBO = placebo; RCT = randomized controlled trial; TOP = topiramate; WMD = weighted mean difference

Table D-23. Valproic acid compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	1; 29	Medium; RCT	Unknown	Direct	Imprecise	81% (VAL) vs. 83% (PBO); P NS	Insufficient
Return to heavy drinking	2; 81	Medium; RCTs	Consistent	Direct	Imprecise	RD: -0.33 (-0.55 to -0.11)	Low
Drinking days	1; 29	Medium; RCT	Unknown	Direct	Imprecise	15.9 (VAL) vs. 19.6 (PBO); P NS	Insufficient
Heavy drinking days	2; 81	Medium; RCTs	Consistent	Direct	Imprecise	WMD: -8.5 (-15.9 to -1.1)	Low
Drinks per drinking day	2; 81	Medium; RCTs	Consistent	Direct	Imprecise	WMD: -2.6 (-5.0 to -0.2)	Low
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

Abbreviations: CI = confidence interval; NA = not applicable; PBO = placebo; RCT = randomized controlled trial; RD = risk difference; VAL = valproic acid; WMD = weighted mean difference

Table D-24. Aripiprazole compared with naltrexone

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	1; 57	Medium; RCT	Unknown	Direct	Imprecise	NSD ³⁸	Insufficient
Return to heavy drinking	1; 57	Medium; RCT	Unknown	Direct	Imprecise	NSD ³⁸	Insufficient
Drinking days	1; 57	Medium; RCT	Unknown	Direct	Imprecise	NSD ³⁸	Insufficient
Heavy drinking days	1; 57	Medium; RCT	Unknown	Direct	Imprecise	NSD ³⁸	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

Abbreviations: CI = confidence interval; NA = not applicable; NSD = no significant difference; RCT = randomized controlled trial

Table D-25. Desipramine compared with paroxetine

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a One included trial, rated as high risk of bias, randomized patients with PTSD and alcohol dependence to desipramine, paroxetine, desipramine plus naltrexone, or paroxetine plus naltrexone.³⁹ The trial found that patients treated with desipramine had fewer heavy drinking days ($P=0.009$) and drinks per drinking day ($P=0.027$) than those who received paroxetine.

Abbreviations: CI = confidence interval; NA = not applicable

Table D-26. Sertraline compared with naltrexone

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	1; 89	Medium; RCT	Unknown	Direct	Imprecise	72.5% (SER) vs. 78.7 (NTX); P NS ³¹	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	1; 89	Medium; RCT	Unknown	Direct	Imprecise	0 (SER) vs. 0 (NTX) ³¹	Insufficient

Abbreviations: CI = confidence interval; NA = not applicable; NTX = naltrexone; RCT = randomized controlled trial; SER = sertraline

Table D-27. Topiramate compared with naltrexone

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Return to any drinking	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Return to heavy drinking	0; 0	NA	NA	NA	NA	NA	Insufficient
Drinking days	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Heavy drinking days	0 ^a ; 0	NA	NA	NA	NA	NA	Insufficient
Drinks per drinking day	0; 0	NA	NA	NA	NA	NA	Insufficient
Accidents	0; 0	NA	NA	NA	NA	NA	Insufficient
Injuries	0; 0	NA	NA	NA	NA	NA	Insufficient
Quality of life or function	0 ^b ; 0	NA	NA	NA	NA	NA	Insufficient
Mortality	0; 0	NA	NA	NA	NA	NA	Insufficient

^a The only included trial that was eligible for KQ 1 and that reported these outcomes, rated as high risk of bias, reported no significant differences between topiramate and naltrexone for proportion of abstinent subjects, cumulative abstinence duration, time to first relapse, or heavy drinking weeks.³⁶ Significantly more subjects in the topiramate group participated in AA than in the naltrexone group (19.2% vs. 4.1%, P=0.04).

^b The two studies that reported this outcome were rated as high risk of bias.

Abbreviations: CI = confidence interval; NA = not applicable

KQ 3

Table D-28. Acamprosate compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Withdrawals due to AEs	11 ^a ; 4,069	Medium; RCTs	Consistent	Direct	Imprecise	RD 0.007 (-0.003 to 0.017)	Low
Anxiety	1 ^b ; 821	Medium; RCT	Unknown	Direct	Imprecise	RD 0.232 (0.174 to 0.290)	Insufficient
Diarrhea	11 ^c ; 3,264	Low to medium; RCTs	Inconsistent	Direct	Imprecise	RD 0.090 (0.019 to 0.160)	Low
Dizziness	0; 0	NA	NA	NA	NA	NA	Insufficient
Headache	5 ^b ; 1,039	Medium; RCTs	Inconsistent	Direct	Imprecise	RD -0.003 (-0.059 to 0.053)	Low
Insomnia	1 ^b ; 116	Medium; RCT	Unknown	Direct	Imprecise	RD 0.038 (-0.030 to 0.106)	Insufficient
Nausea	5 ^b ; 1,623	Low; RCTs	Consistent	Direct	Imprecise	RD 0.009 (-0.009 to 0.027)	Moderate
Numbness / tingling / paresthesias	1 ^b ; 262	Medium; RCT	Unknown	Direct	Imprecise	RD 0.008 (-0.013 to 0.029)	Insufficient
Rash	1 ^b ; 246	Medium; RCT	Unknown	Direct	Imprecise	RD -0.008 (-0.030 to 0.014)	Insufficient
Suicide attempts or suicidal ideation	1 ^c ; 581	Medium; RCT	Unknown	Direct	Imprecise	RD 0.007 (-0.005, 0.019)	Insufficient
Taste abnormalities	0; 0	NA	NA	NA	NA	NA	Insufficient
Vomiting	3 ^b ; 1,782	Medium; RCTs	Consistent	Direct	Precise	RD 0.024 (0.007 to 0.042)	Moderate

^a Three additional studies were rated high or unclear risk of bias

^b One additional study was rated high or unclear risk of bias

^c Two additional studies were rated high or unclear risk of bias

Abbreviations: AE = adverse effect; CI = confidence interval; RCT = randomized controlled trial; RD risk difference

Table D-29. Naltrexone compared with placebo

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI)	Strength of Evidence Grade
Withdrawals due to AEs	14 ^a ; 2,203	Medium; RCTs	Consistent	Direct	Precise	RD 0.024 (0.009 to 0.038)	Moderate
Anxiety	5 ^a ; 725	Medium; RCTs	Consistent	Direct	Imprecise	RD 0.033 (-0.012 to 0.078)	Low
Diarrhea	9 ^b ; 2,232	Medium; RCTs	Consistent	Direct	Imprecise	RD 0.011 (-0.018 to 0.041)	Moderate
Dizziness	11 ^c ; 2,549	Medium; RCTs	Consistent	Direct	Precise	RD 0.068 (0.037 to 0.099)	Moderate
Headache	14 ^d ; 3,102	Medium; RCTs	Inconsistent	Direct	Imprecise	RD 0.010 (-0.020 to 0.039)	Low
Insomnia	6 ^c ; 1,571	Medium; RCTs	Consistent	Direct	Imprecise	RD 0.015 (-0.016 to 0.046)	Low
Nausea	22 ^e ; 4,320	Medium; RCTs	Consistent	Direct	Precise	RD 0.114 (0.073 to 0.155)	Moderate
Numbness / tingling / paresthesias	1 ^b ; 246	Medium; RCT	Unknown	Direct	Imprecise	RD 0.032 (-0.093 to 0.157)	Insufficient
Rash	2 ^b ; 134	Medium; RCTs	Consistent	Direct	Imprecise	RD 0.056 (-0.128 to 0.241)	Low
Suicide	0; 0	NA	NA	NA	NA	NA	Insufficient
Taste abnormalities	0; 0	NA	NA	NA	NA	NA	Insufficient
Vomiting	7 ^a ; 2,103	Medium; RCTs	Consistent	Direct	Precise	RD 0.054 (0.028 to 0.079)	Moderate

^a Two additional studies were rated high or unclear risk of bias

^b One additional study was rated high or unclear risk of bias

^c Four additional studies were rated high or unclear risk of bias

^d Five additional studies were rated high or unclear risk of bias

^e Seven additional studies were rated as high or unclear risk of bias

Abbreviations: AE = adverse effect; CI = confidence interval; RD = risk difference

Table D-30. Acamprosate compared with Naltrexone

Outcome	Number of Studies; Number of Subjects	Risk of Bias; Design	Consistency	Directness	Precision	Summary Effect Size (95% CI) ^a	Strength of Evidence Grade
Withdrawals due to AEs	1 ^a ; 612	Medium; RCT	Unknown	Direct	Imprecise	RD -0.009 (-0.038 to 0.020)	Insufficient
Anxiety	0; 0	NA	NA	NA	NA	NA	Insufficient
Diarrhea	3 ^b ; 800	Low to medium; RCTs	Consistent	Direct	Imprecise	RD 0.202 (-0.019 to 0.422)	Moderate
Dizziness	1 ^b ; 108	Medium; RCT	Unknown	Direct	Imprecise	RD -0.020 (-0.082 to 0.043)	Insufficient
Headache	1 ^c ; 108	Medium; RCT	Unknown	Direct	Imprecise	RD -0.058 (-0.151 to 0.035)	Insufficient ^d
Insomnia	0; 0	NA	NA	NA	NA	NA	Insufficient
Nausea	3 ^c ; 800	Medium; RCTs	Consistent	Direct	Imprecise	RD -0.048 (-0.124 to 0.028)	Low
Numbness / tingling / paresthesias	0; 0	NA	NA	NA	NA	NA	Insufficient
Rash	0; 0	NA	NA	NA	NA	NA	Insufficient
Suicide	0; 0	NA	NA	NA	NA	NA	Insufficient
Taste abnormalities	0; 0	NA	NA	NA	NA	NA	Insufficient
Vomiting	1; 612	Low; RCT	Unknown	Direct	Imprecise	8.91% (ACA) vs. 14.6% (NTX); P NR	Insufficient

^a In this column, a positive value favors naltrexone

^b One additional study was rated high or unclear risk of bias

^c Two additional studies were rated high risk of bias

^d The two additional studies rated as high risk of bias found similar results as the medium risk of bias study. Meta-analysis including all three found a higher risk of headache with naltrexone than with acamprosate: RD -0.10 (-0.17, -0.03)

Abbreviations: ACA = acamprosate; AE = adverse effect; CI = confidence interval; NTX = naltrexone; RCT = randomized controlled trial; RD = risk difference

References for Appendix D

1. Lhuintre JP, Moore N, Tran G, et al. Acamprosate appears to decrease alcohol intake in weaned alcoholics. *Alcohol Alcohol*. 1990;25(6):613-22. PMID: 2085344.
2. Anton RF, O'Malley SS, Ciraulo DA, et al. Combined pharmacotherapies and behavioral interventions for alcohol dependence: the COMBINE study: a randomized controlled trial. *JAMA*. 2006 May 3;295(17):2003-17. PMID: 16670409.
3. Petrakis IL, Poling J, Levinson C, et al. Naltrexone and disulfiram in patients with alcohol dependence and comorbid psychiatric disorders. *Biol Psychiatry*. 2005 May 15;57(10):1128-37. PMID: 15866552.
4. Morgenstern J, Kuerbis AN, Chen AC, et al. A randomized clinical trial of naltrexone and behavioral therapy for problem drinking men who have sex with men. *J Consult Clin Psychol*. 2012;80(5):863-75. PMID: 22612306.
5. LoCastro JS, Youngblood M, Cisler RA, et al. Alcohol treatment effects on secondary nondrinking outcomes and quality of life: the COMBINE study. *J Stud Alcohol Drugs*. 2009 Mar;70(2):186-96. PMID: 19261230.
6. Pettinati HM, Gastfriend DR, Dong Q, et al. Effect of extended-release naltrexone (XR-NTX) on quality of life in alcohol-dependent patients. *Alcohol Clin Exp Res*. 2009 Feb;33(2):350-6. PMID: 19053979.
7. O'Malley SS, Robin RW, Levenson AL, et al. Naltrexone alone and with sertraline for the treatment of alcohol dependence in Alaska natives and non-natives residing in rural settings: a randomized controlled trial. *Alcohol Clin Exp Res*. 2008 Jul;32(7):1271-83. PMID: 18482155.
8. Laaksonen E, Koski-Jannes A, Salaspuro M, et al. A randomized, multicentre, open-label, comparative trial of disulfiram, naltrexone and acamprosate in the treatment of alcohol dependence. *Alcohol Alcohol*. 2008 Jan-Feb;43(1):53-61. PMID: 17965444.
9. Morley KC, Teesson M, Reid SC, et al. Naltrexone versus acamprosate in the treatment of alcohol dependence: A multi-centre, randomized, double-blind, placebo-controlled trial. *Addiction*. 2006 Oct;101(10):1451-62. PMID: 16968347.
10. Morley KC, Teesson M, Sannibale C, et al. Clinical predictors of outcome from an Australian pharmacological relapse prevention trial. *Alcohol Alcohol*. 2010 Nov-Dec;45(6):520-6. PMID: 20952764.
11. Wilens TE, Adler LA, Weiss MD, et al. Atomoxetine treatment of adults with ADHD and comorbid alcohol use disorders. *Drug Alcohol Depend*. 2008 Jul 1;96(1-2):145-54. PMID: 18403134.
12. Addolorato G, Leggio L, Ferrulli A, et al. Effectiveness and safety of baclofen for maintenance of alcohol abstinence in alcohol-dependent patients with liver cirrhosis: randomised, double-blind controlled study. *Lancet*. 2007 Dec 8;370(9603):1915-22. PMID: 18068515.
13. Garbutt JC, Kampov-Polevoy AB, Gallop R, et al. Efficacy and safety of baclofen for alcohol dependence: a randomized, double-blind, placebo-controlled trial. *Alcohol Clin Exp Res*. 2010 Nov;34(11):1849-57. PMID: 20662805.
14. Malec E, Malec T, Gagne MA, et al. Buspirone in the treatment of alcohol dependence: a placebo-controlled trial. *Alcohol Clin Exp Res*. 1996 Apr;20(2):307-12. PMID: 8730222.
15. Malcolm R, Anton RF, Randall CL, et al. A placebo-controlled trial of buspirone in anxious inpatient alcoholics. *Alcohol Clin Exp Res*. 1992 Dec;16(6):1007-13. PMID: 1335217.
16. Tiihonen J, Ryyanen OP, Kauhanen J, et al. Citalopram in the treatment of alcoholism: a double-blind placebo-controlled study. *Pharmacopsychiatry*. 1996 Jan;29(1):27-9. PMID: 8852531.
17. Naranjo CA, Bremner KE, Lanctot KL. Effects of citalopram and a brief psychosocial intervention on alcohol intake, dependence and problems. *Addiction*. 1995 Jan;90(1):87-99. PMID: 7888983.

18. Mason BJ, Kocsis JH, Ritvo EC, et al. A double-blind, placebo-controlled trial of desipramine for primary alcohol dependence stratified on the presence or absence of major depression. *JAMA*. 1996 Mar 13;275(10):761-7. PMID: 8598592.
19. Kabel DI, Petty F. A placebo-controlled, double-blind study of fluoxetine in severe alcohol dependence: adjunctive pharmacotherapy during and after inpatient treatment. *Alcohol Clin Exp Res*. 1996 Jun;20(4):780-4. PMID: 8800399.
20. Cornelius JR, Salloum IM, Ehler JG, et al. Fluoxetine in depressed alcoholics. A double-blind, placebo-controlled trial. *Arch Gen Psychiatry*. 1997 Aug;54(8):700-5. PMID: 9283504.
21. Chick J, Aschauer H, Hornik K. Efficacy of fluvoxamine in preventing relapse in alcohol dependence: a one-year, double-blind, placebo-controlled multicentre study with analysis by typology. *Drug Alcohol Depend*. 2004 Apr 9;74(1):61-70. PMID: 15072808.
22. McGrath PJ, Nunes EV, Stewart JW, et al. Imipramine treatment of alcoholics with primary depression: A placebo-controlled clinical trial. *Arch Gen Psychiatry*. 1996 Mar;53(3):232-40. PMID: 8611060.
23. Mason BJ, Ritvo EC, Morgan RO, et al. A double-blind, placebo-controlled pilot study to evaluate the efficacy and safety of oral nalmefene HCl for alcohol dependence. *Alcohol*. 1994;18(5):1162-7.
24. Guardia J, Segura L, Gonzalvo B, et al. A double-blind, placebo-controlled study of olanzapine in the treatment of alcohol-dependence disorder. *Alcohol Clin Exp Res*. 2004 May;28(5):736-45. PMID: 15166648.
25. Book SW, Thomas SE, Randall PK, et al. Paroxetine reduces social anxiety in individuals with a co-occurring alcohol use disorder. *J Anxiety Disord*. 2008;22(2):310-8. PMID: 17448631.
26. Thomas SE, Randall PK, Book SW, et al. A complex relationship between co-occurring social anxiety and alcohol use disorders: what effect does treating social anxiety have on drinking? *Alcohol Clin Exp Res*. 2008 Jan;32(1):77-84. PMID: 18028529.
27. Kampman KM, Pettinati HM, Lynch KG, et al. A double-blind, placebo-controlled pilot trial of quetiapine for the treatment of Type A and Type B alcoholism. *J Clin Psychopharmacol*. 2007 Aug;27(4):344-51. PMID: 17632217.
28. Brown ES, Garza M, Carmody TJ. A randomized, double-blind, placebo-controlled add-on trial of quetiapine in outpatients with bipolar disorder and alcohol use disorders. *J Clin Psychiatry*. 2008 May;69(5):701-5. PMID: 18312058.
29. Stedman M, Pettinati HM, Brown ES, et al. A double-blind, placebo-controlled study with quetiapine as adjunct therapy with lithium or divalproex in bipolar I patients with coexisting alcohol dependence. *Alcohol Clin Exp Res*. 2010 Oct;34(10):1822-31. PMID: 20626727.
30. Brady KT, Sonne S, Anton RF, et al. Sertraline in the treatment of co-occurring alcohol dependence and posttraumatic stress disorder. *Alcohol Clin Exp Res*. 2005 Mar;29(3):395-401. PMID: 15770115.
31. Pettinati HM, Oslin DW, Kampman KM, et al. A double-blind, placebo-controlled trial combining sertraline and naltrexone for treating co-occurring depression and alcohol dependence. *Am J Psychiatry*. 2010 Jun;167(6):668-75. PMID: 20231324.
32. Pettinati HM, Volpicelli JR, Luck G, et al. Double-blind clinical trial of sertraline treatment for alcohol dependence. *J Clin Psychopharmacol*. 2001 Apr;21(2):143-53. PMID: 11270910.
33. Gual A, Balcells M, Torres M, et al. Sertraline for the prevention of relapse in detoxicated alcohol dependent patients with a comorbid depressive disorder: a randomized controlled trial. *Alcohol Alcohol*. 2003 Nov-Dec;38(6):619-25. PMID: 14633652.
34. Johnson BA, Rosenthal N, Capece JA, et al. Topiramate for treating alcohol dependence: a randomized controlled trial. *JAMA*. 2007 Oct 10;298(14):1641-51. PMID: 17925516.
35. Johnson BA, Ait-Daoud N, Bowden CL, et al. Oral topiramate for treatment of alcohol dependence: a randomised controlled trial. *Lancet*. 2003 May 17;361(9370):1677-85. PMID: 12767733.

36. Baltieri DA, Daro FR, Ribeiro PL, et al. Comparing topiramate with naltrexone in the treatment of alcohol dependence. *Addiction*. 2008 Dec;103(12):2035-44. PMID: 18855810.
37. Rubio G, Martinez-Gras I, Manzanares J. Modulation of impulsivity by topiramate: implications for the treatment of alcohol dependence. *J Clin Psychopharmacol*. 2009 Dec;29(6):584-9. PMID: 19910725.
38. Martinotti G, Di Nicola M, Di Giannantonio M, et al. Aripiprazole in the treatment of patients with alcohol dependence: a double-blind, comparison trial vs. naltrexone. *J Psychopharmacol*. 2009 Mar;23(2):123-9. PMID: 18515460.
39. Petrakis IL, Ralevski E, Desai N, et al. Noradrenergic vs serotonergic antidepressant with or without naltrexone for veterans with PTSD and comorbid alcohol dependence. *Neuropsychopharmacology*. 2012 Mar;37(4):996-1004. PMID: 22089316.

Appendix E. Placebo-Controlled Trials of Medications Used Off-label, or Those Under Investigation for Which We Found Only 1 Trial Meeting Inclusion Criteria

Aripiprazole

Characteristics of Trials

Table E-1 summarizes characteristics of the one trial meeting our inclusion criteria.¹ It was conducted across 16 academic centers in the U.S. and compared aripiprazole, titrated from 2 mg/day to 30 mg/day over the initial four weeks, with placebo. All participants (N=295) received an enhanced form of cognitive behavioral therapy. The recruitment method was not reported. All patients were alcohol-dependent, and the proportions of smokers and of patients with co-occurring conditions were not reported.

Table E-1. Characteristics of included double-blind randomized placebo-controlled trials of aripiprazole

Author, Year	Arm Dose, mg/day (N)	Medication Duration (Followup)	Country	Setting	Age Years	Per-cent-age Non-White	Percent-age Female	Co-intervention(s)	Risk of Bias
Anton, 2008 ¹	Aripiprazole titrated from 2 to 30 over 4 wks (149) Placebo (146)	12	U.S.	16 academic centers	47	15 to 16	25 to 38	Enhanced CBT 100%	Med

Notes: Age (y) is the mean age in years, unless otherwise stated

Abbreviations: CBT = cognitive behavioral therapy; mg = milligrams; U.S. = United States; wks = weeks

Return to Any Drinking

More patients treated with aripiprazole returned to any drinking than with placebo (89 percent versus 78 percent, $p=0.02$).

Return to Heavy Drinking

The proportion returning to heavy drinking did not differ between groups (73 percent versus 73 percent).

Drinking Days

Patients treated with aripiprazole drank on a mean of 41 percent of days; patients treated with placebo drank on 37 percent of days. The difference was not statistically significant.

Drinks per Drinking Day

Patients treated with aripiprazole reported fewer drinks per drinking day than patients treated with placebo (4.4 versus 5.5, $p<0.001$).

Atomoxetine

Characteristics of Trials

Table E-2 summarizes characteristics of the one trial meeting our inclusion criteria.² It was a multi-institutional study conducted in the U.S. and Canada. Investigators compared atomoxetine, titrated from 25 mg/day to 100 mg/day, with placebo. The mean final dose for atomoxetine was 89.9mg. Twelve-step program attendance was allowed, but all other types of co-intervention were prohibited. The recruitment method was not reported. Slightly more than half of the patients met criteria for alcohol dependence, and all patients were diagnosed with attention deficit hyperactivity disorder (ADHD). The proportion of smokers was not reported. The study was rated as high risk of bias, primarily for high risk of attrition bias and inadequate handling of missing data (see Appendix C for details).

Table E-2 . Characteristics of included double-blind randomized placebo-controlled trials of atomoxetine

Author, Year	Arm Dose, mg/day (N)	Medication Duration (Follow-up)	Country	Setting	Age Years	Per-cent-age Non-White	Per-cent-age F	Percentage With Co-Occurring Condition(s)	Risk of Bias
Wilens, 2008 ²	Atomoxetine titrated from 25 to 100 (72) Placebo (75)	12	U.S. and Canada	Multi-institution	35	12	15	100 (ADHD)	High

Notes: Age (y) is the mean age in years, unless otherwise stated

Abbreviations: ADHD, attention deficit hyperactivity disorder; mg, milligrams; Rx = prescription; U.S., United States

Return to Heavy Drinking

The trial found no significant difference between groups. (94 percent for patients treated with atomoxetine versus 96 percent for patients taking placebo).

Drinking Days

Patients treated with atomoxetine drank on roughly 50 percent of days; patients treated with placebo drank on roughly 60 percent of days. The difference between groups was not statistically significant.

Heavy Drinking Days

Atomoxetine-treated patients had a 26 percent lower rate of heavy drinking days compared with placebo (event ratio=0.74, p=0.02).

Drinks per Drinking Day

The reduction in drinks per drinking day from baseline was not significantly different between groups (1.1 for the atomoxetine group versus 0.6 for the placebo group, p NS).

Desipramine

Characteristics of Trials

Table E-3 summarizes characteristics of the one trial meeting our inclusion criteria.³ It was conducted in the outpatient psychiatry departments at two urban medical centers in the U.S. Investigators compared desipramine (median dose=200 mg/day) with placebo. No co-interventions were required, though Alcoholics Anonymous attendance was encouraged. Patients were recruited through inpatient and outpatient psychiatric referrals and via public service announcements. All patients met criteria for alcohol dependence, and 39 percent were also diagnosed with depression. The proportion of smokers was not reported. The study was rated as high risk of bias, primarily for high risk of attrition bias and inadequate handling of missing data (see Appendix C for details).

Table E-3. Characteristics of included double-blind randomized placebo-controlled trials of desipramine

Author, Year	Arm Dose, mg/day (N)	Medication Duration (follow-up)	Country	Setting	Age Years	Percentage Non-White	Percentage Female	Percentage With Co-Occurring Condition(s)	Risk of Bias
Mason, 1996 ³	Desipramine median 200 (37) Placebo (34)	26	U.S.	Psychiatry outpatient departments at 2 urban medical centers	Median= 40	38	17	Depression 39%	High

Notes: Age (y) is the mean age in years, unless otherwise stated

Abbreviations: U.S. = United States; wks = weeks

Return to Heavy Drinking

In this study, a return to heavy drinking was defined as two heavy drinking days per week for two consecutive weeks. Twelve percent of patients in the desipramine arm returned to heavy drinking, compared with 32 percent of patients taking placebo. The difference between groups was not statistically significant.

Drinking Days

Non-depressed patients treated with desipramine (N=14) drank on a median of 68 percent of days; non-depressed patients treated with placebo (N=15) drank on 72 percent of days (p NS). Depressed patients treated with desipramine (N=12) drank on a median of 40 percent of days; depressed patients treated with placebo (N=10) drank on 64 percent of days (p NS).

Fluvoxamine

Characteristics of Trials

Table E-4 summarizes characteristics of the one trial meeting our inclusion criteria.⁴ It was conducted in ten outpatient sites in four European countries. Investigators compared fluvoxamine 100-300 mg/day with placebo. In addition, patients received each site's usual psychosocial treatment. Recruitment method was not reported. All patients met criteria for alcohol dependence; the study did not report the proportion of patients with co-occurring conditions. The proportions of smokers and non-white participants were not reported.

Table E-4. Characteristics of Included double-blind randomized placebo-controlled trials of fluvoxamine

Author, Year	Arm Dose, mg/day (N)	Medication Duration (Follow-up)	Country	Setting	Age Years	Percentage Female	Risk of Bias
Chick, 2004 ⁴	Fluvoxamine 100-300 (261) Placebo (260)	52	U.K., Eire, Austria, Switzerland	10 outpatient sites	42 (19-72)	35	Med

Notes: Age (y) is the mean age in years, unless otherwise stated

Abbreviations: mg = milligram; N = number; U.K. = United Kingdom

Return to Any Drinking

At 12 weeks, the trial found no significant difference between groups (58 percent of fluvoxamine-treated patients versus 54 percent of placebo-treated patients returned to drinking since the assessment at week 8, P NS). Similarly, at 52 weeks, there was no difference between groups in percentage of patients who returned to drinking since the previous assessment at week 40 (71 percent versus 71 percent).

Return to Heavy Drinking

At 12 weeks, the study did not find a difference between those treated with fluvoxamine and those who received placebo (46 percent versus 40 percent, P NS). Similarly, at 52 weeks, the study reported no difference between groups (64 percent versus 64 percent).

Drinking Days

At 12 weeks, fluvoxamine-treated patients had more drinking days since the previous (at week 8) assessment than placebo-treated patients (31 percent versus 23 percent, p=0.009). At 52 weeks, the study did not find a significant difference between groups for drinking days since the previous assessment (44 percent versus 38 percent, p NS).

Mortality

During the 52-week study, there was no difference between those treated with fluvoxamine and those who received placebo: one patient in each arm died.

Imipramine

Characteristics of Trials

Table E-5 summarizes characteristics of the one trial meeting our inclusion criteria.⁵ It was conducted in a university-based depression research clinic in the U.S. Investigators compared imipramine 50-300 mg/day (mean dose=262 mg/day) with placebo. In addition, all patients received individual relapse prevention counseling. Patients were recruited using advertisements and via referrals. Almost all patients met criteria for alcohol dependence (96 percent), and all had some form of depression. The proportion of smokers was not reported. Roughly 20 percent of enrollees were non-white, and about half were female.

Table E-5. Characteristics of included double-blind randomized placebo-controlled trials of imipramine

Author, Year	Arm Dose, mg/day (N)	Medication Duration (Follow-up)	Country	Setting	Age Years	Percentage Non-White	Percentage Female	Percentage With Co-Occurring Condition(s)	Risk of Bias
McGrath, 1996 ⁵	Imipramine 50-300; mean 262 (36) Placebo (33)	12	U.S.	University-based depression research clinic	37 imipramine, 22 placebo ^a	17 to 22	49 to 53	MDD 71 to 72 Bipolar 11 to 12 Atypical depression 70 to 72 Other substance abuse 16	Med

^aThe study reported 11 percent, but it was clearly a reporting error; likely 31 or 41 percent.

Notes: Age (y) is the mean age in years, unless otherwise stated

Abbreviations: MDD = major depressive disorder; mg = milligrams; U.S. = United States

Return to Any Drinking

The study found no significant difference between groups (69 percent of those receiving imipramine versus 79 percent of those receiving placebo, p NS).

Drinking Days

The study found no significant difference between groups (Imipramine-treated patients drank on 28 percent of days versus 31 percent for those who received placebo, p NS).

Heavy Drinking Days

The study found no significant difference between groups (Imipramine-treated patients drank heavily on 13.5 percent of days versus 9 percent for those who received placebo, p NS).

Drinks per Drinking Day

The mean number of drinks per drinking day was 3.7 for imipramine-treated patients and 4.1 for placebo-treated patients. The difference between groups was not statistically significant.

Olanzapine

Characteristics of Trials

Table E-6 summarizes characteristics of the one trial meeting our inclusion criteria.⁶ Patients were treatment-seekers in a psychiatry department-based addictive behavior unit in a hospital in Spain. Investigators compared olanzapine 5 to 15 mg/day with placebo. In addition, all patients received cognitive behavioral therapy. All patients met criteria for alcohol dependence. The proportions of smokers and non-white patients were not reported.

Table E-6. Characteristics of included double-blind randomized placebo-controlled trials of olanzapine

Author, Year	Arm Dose, mg/day (N)	Medication Duration (Follow-up)	Country	Setting	Age Years	Percentage Female	Risk of Bias
Guardia, 2004 ⁶	Olanzapine 5-15 (29) Placebo (31)	12 (16)	Spain	Addictive behavior unit of a hospital psychiatry department	43	23 to 27	Med

Notes: Age (y) is the mean age in years, unless otherwise stated

Abbreviations: mg = milligrams; N = number; Rx = prescription

Return to Heavy Drinking

Thirty-eight percent of olanzapine patients returned to heavy drinking by the end of the study compared with 29 percent of placebo-treated patients. The difference between groups was not statistically significant.

Drinking Days

Olanzapine-treated patients drank on 13 percent of days; those who received placebo drank on 23 percent of days. The difference between groups was not statistically significant.

Drinks per Drinking Day

The mean number of drinks per drinking day was 1.8 for olanzapine-treated patients and 2.0 for placebo-treated patients. This difference was not statistically significant.

Paroxetine

Characteristics of Trials

Table E-7 summarizes characteristics of the one trial meeting our inclusion criteria.^{7,8} It was conducted in the U.S., but the specific setting was not reported. Investigators compared paroxetine, titrated from 10 to 60 mg/day over four weeks (mean dose=45 mg/day) with placebo; no psychosocial or psychological therapy was provided. Patients were recruited using media advertisements. Most of the patients met criteria for alcohol dependence (79 percent), and all had social anxiety disorder. Roughly ten percent were also diagnosed with major depression. The proportion of smokers was not reported.

Table E-7. Characteristics of included double-blind randomized placebo-controlled trials of paroxetine

Author, Year	Arm Dose, mg/day (N)	Medication Duration (Follow-up)	Country	Age Years	Percentage Non-White	Percentage Female	Percentage With Co-Occurring Condition(s)	Risk of Bias
Book, 2008 ⁷ ; Thomas, 2008 ⁸	Paroxetine titration over 4 weeks 10-60; avg. 45 (20) Placebo (22)	16	U.S.	28 to 30	0 to 18	45 to 50	Social anxiety disorder 100%; MDD ~10	Med

Notes: Age (y) is the mean age in years, unless otherwise stated

Abbreviations: MDD = major depressive disorder; mg = milligrams; U.S. = United States

Drinking Days

The study found no significant difference between groups (paroxetine-treated patients versus placebo-treated: 34 percent versus 35 percent, p NS).

Heavy Drinking Days

The study found no significant difference between groups (paroxetine-treated patients versus placebo-treated: 54 percent versus 55 percent, p NS).

Drinks per Drinking Day

At week 16, the mean number of drinks per drinking day was 5.9 for paroxetine-treated patients and 7.0 for placebo-treated patients. The difference between groups was not statistically significant.


References for Appendix E

1. Anton RF, Kranzler H, Breder C, et al. A randomized, multicenter, double-blind, placebo-controlled study of the efficacy and safety of aripiprazole for the treatment of alcohol dependence. *J Clin Psychopharmacol*. 2008 Feb;28(1):5-12. PMID: 18204334.
2. Wilens TE, Adler LA, Weiss MD, et al. Atomoxetine treatment of adults with ADHD and comorbid alcohol use disorders. *Drug Alcohol Depend*. 2008 Jul 1;96(1-2):145-54. PMID: 18403134.
3. Mason BJ, Kocsis JH, Ritvo EC, et al. A double-blind, placebo-controlled trial of desipramine for primary alcohol dependence stratified on the presence or absence of major depression. *JAMA*. 1996 Mar 13;275(10):761-7. PMID: 8598592.
4. Chick J, Aschauer H, Hornik K. Efficacy of fluvoxamine in preventing relapse in alcohol dependence: a one-year, double-blind, placebo-controlled multicentre study with analysis by typology. *Drug Alcohol Depend*. 2004 Apr 9;74(1):61-70. PMID: 15072808.
5. McGrath PJ, Nunes EV, Stewart JW, et al. Imipramine treatment of alcoholics with primary depression: A placebo-controlled clinical trial. *Arch Gen Psychiatry*. 1996 Mar;53(3):232-40. PMID: 8611060.
6. Guardia J, Segura L, Gonzalvo B, et al. A double-blind, placebo-controlled study of olanzapine in the treatment of alcohol-dependence disorder. *Alcohol Clin Exp Res*. 2004 May;28(5):736-45. PMID: 15166648.
7. Book SW, Thomas SE, Randall PK, et al. Paroxetine reduces social anxiety in individuals with a co-occurring alcohol use disorder. *J Anxiety Disord*. 2008;22(2):310-8. PMID: 17448631.
8. Thomas SE, Randall PK, Book SW, et al. A complex relationship between co-occurring social anxiety and alcohol use disorders: what effect does treating social anxiety have on drinking? *Alcohol Clin Exp Res*. 2008 Jan;32(1):77-84. PMID: 18028529.

Appendix F. Meta-Analyses


Key Question 1 Meta-Analysis Results: Acamprosate versus Placebo

Acamprosate versus Placebo Return to Any Drinking by Risk of Bias Rating




Note: doses combined for Mason, 2006, Paille, 1995, and Pelc, 1997

Acamprosate versus Placebo Return to Any Drinking by Country




Note: doses combined for Mason, 2006, Paille, 1995, and Pelc, 1997

Acamprosate versus Placebo Return to Any Drinking by Duration of Treatment




Note: doses combined for Mason, 2006, Paille, 1995, and Pelc, 1997

Acamprosate versus Placebo Return to Heavy Drinking by Risk of Bias Rating




Note: doses combined for Mason, 2006


Acamprosate versus Placebo - Percent Drinking Days by Risk of Bias




Acamprosate versus Placebo - Percent Drinking Days by Country




Acamprosate versus Placebo - Percent Drinking Days by Duration of Treatment



Acamprosate versus Placebo - Drinks per Drinking Day by Risk of Bias



Key Question 1 Meta-Analysis Results: Disulfiram versus Control




Note: Control - Fuller, 1979 and Fuller, 1986 control = Disulfiram 1 mg; Petrakis, 2005 control = placebo


author/year	t_event	t_noevent	c_event	c_noevent	RD (95% CI)	Weight
Fuller, 1979	34	9	37	5	-0.09 (-0.25, 0.07)	18.43
Fuller, 1986	164	38	167	32	-0.03 (-0.10, 0.05)	81.57
Subtotal (I-squared = 0.0%, p = 0.475)					-0.04 (-0.11, 0.03)	100.00
Overall (I-squared = 0.0%, p = 0.475)					-0.04 (-0.11, 0.03)	100.00

NOTE: Weights are from random effects analysis


Naltrexone versus Placebo Return to Any Drinking by Risk of Bias Rating




Naltrexone versus Placebo Return to Any Drinking by Country



Naltrexone versus Placebo Return to Any Drinking by Duration of Treatment




Naltrexone versus Placebo Return to Any Drinking by NTX Dose




Note: Dose for Oslin, 1997 represents an average of 50 mg.day; study participants received 100 mg two times per week and 150 mg one time per week.


Naltrexone versus Placebo Return to Heavy Drinking by Risk of Bias Rating




Naltrexone versus Placebo Return to Heavy Drinking by Country



Naltrexone versus Placebo Return to Heavy Drinking by Duration of Treatment




Naltrexone versus Placebo Return to Heavy Drinking by NTX Dose




Note: Dose for Oslin, 1997 represents an average of 50 mg/day; study participants received 100 mg two times per week and 150 mg one time per week.

Naltrexone versus Placebo - Percent Drinking Days by Risk of Bias




Note: Brown, 2009 not included


Naltrexone versus Placebo - Percent Drinking Days by Country



Naltrexone versus Placebo - Percent Drinking Days by Duration of Treatment




Naltrexone versus Placebo - Percent Drinking Days by NTX Dose




Note: Dose for Oslin, 1997 represents an average of 50 mg.day; study participants received 100 mg two times per week and 150 mg one time per week.


Naltrexone v. Placebo - % Heavy Drinking Days by Risk of Bias




Naltrexone v. Placebo - % Heavy Drinking Days by Country




Naltrexone v. Placebo - % Heavy Drinking Days by Duration of Treatment



Naltrexone v. Placebo - % Heavy Drinking Days by NTX Dose




Naltrexone versus Placebo - Drinks per Drinking Day by Risk of Bias




Note: Monti, 2001 and Krystal, 2001 are rated High risk of bias for this outcome because of attrition


Naltrexone versus Placebo - Drinks per Drinking Day by Country




Naltrexone versus Placebo - Drinks per Drinking Day by Duration of Treatment




Naltrexone versus Placebo - Drinks per Drinking Day by NTX Dose




Key Question 1 Meta-Analysis Results: Acamprosate versus Naltrexone



Acamprosate versus Naltrexone Return to Heavy Drinking by Risk of Bias




Acamprosate versus Naltrexone - Percent Drinking Days by Risk of Bias




Key Question 1 Meta-Analysis Results: Off label medications versus placebo


Buspirone versus Placebo Return to Any Drinking by Risk of Bias Rating




Fluoxetine versus Placebo Return to Any Drinking by Risk of Bias Rating




Divalproex/Valproate versus Placebo Return to Heavy Drinking by Risk of Bias Rating




Nalmefene versus Placebo Return to Heavy Drinking by Risk of Bias Rating



Sertraline versus Placebo Return to Heavy Drinking by Risk of Bias Rating




Buspirone versus Placebo - Percent Drinking Days by Risk of Bias




Note: 12 week time point used for Fawcett study


Fluoxetine versus Placebo - Percent Drinking Days by Risk of Bias




Nalmefene versus Placebo - Percent Drinking Days by Risk of Bias




Quetiapine versus Placebo - Percent Drinking Days by Risk of Bias



Sertraline versus Placebo - Percent Drinking Days by Risk of Bias




Topiramate versus Placebo - Percent Drinking Days by Risk of Bias




Note: Johnson, 2003 not included


Divalproex/Valproate versus Placebo - Percent Heavy Drinking Days by Risk of Bias



Quetiapine versus Placebo - Percent Heavy Drinking Days by Risk of Bias




Topiramate versus placebo - % Heavy Drinking Days




Heterogeneity			
Q-value	df (Q)	P-value	I-squared
2.215796	2	0.330252	9.738993

Topiramate versus placebo - % Heavy Drinking Days - Sensitivity Analysis




Heterogeneity			
Q-value	df (Q)	P-value	I-squared
1.505758	1	0.219787	33.58828

Divalproex/Valproate versus Placebo - Drinks Per Drinking Day by Risk of Bias




Fluoxetine versus Placebo - Drinks Per Drinking Day by Risk of Bias




Note: Kranzler, 1995 – Ns for each group, are not the total randomized (total randomized: FLU = 51; PBO = 50)

Nalmefene versus Placebo - Drinks Per Drinking Day by Risk of Bias




Note: Karhuvarra, 2007 Ns for each group are not the total randomized (total randomized: NAL = 242; PBO = 161)

Sertraline versus Placebo - Drinks Per Drinking Day by Risk of Bias




Topiramate versus placebo - Drinks per Drinking Day



Heterogeneity			
Q-value	df (Q)	P-value	I-squared
1.043611	2	0.593448	0

Topiramate versus placebo - Drinks per Drinking Day - Sensitivity Analysis




Heterogeneity			
Q-value	df (Q)	P-value	I-squared
0.156202	1	0.692677	0

KQ3 Analyses – Harms

We found insufficient data for all included medications to perform meta-analyses for the following harms: Anorexia, Cognitive Dysfunction, Glaucoma, Metabolic Acidosis, Palpitations, Taste,


Anxiety – Acamprosate versus placebo

Acamprosate versus Placebo Anxiety by Risk of Bias Rating




Anxiety – Naltrexone versus placebo

Naltrex versus Placebo Anxiety by Risk of Bias Rating




Diarrhea – Acamprosate versus placebo

Acamp versus Placebo Diarrhea by Risk of Bias Rating




Diarrhea – Naltrexone versus placebo




Diarrhea – Acamprosate versus naltrexone

Acamprosate versus Naltrexone Diarrhea by Risk of Bias Rating




Dizziness – Naltrexone versus placebo

Naltrexone versus Placebo DIZZY by Risk of Bias Rating




Dizziness – Acamprosate versus naltrexone

Acamp versus Naltrex Dizzy by Risk of Bias Rating




Headache – Acamprosate versus placebo

Acamp versus Placebo headache by Risk of Bias Rating




Headache – Naltrexone versus placebo

Naltrexone versus Placebo Headache by Risk of Bias Rating




Headache – Acamprosate versus naltrexone

Acamp versus Naltrexone Headache by Risk of Bias Rating




Insomnia – Acamprosate versus placebo




Insomnia – Naltrexone versus placebo

Naltrexone versus Placebo Insomnia by Risk of Bias Rating




Nausea – Acamprosate versus placebo




Footnote: The Poldrugo, 1997 study reported nausea/vomiting; the risk difference in an analysis run without this study was also not statistically significant [RD: 0.01 (95% CI -0.02, 0.04)].

Nausea – Acamprosate versus placebo – Sensitivity Analysis



Nausea – Naltrexone versus placebo


Naltrexone versus Placebo Nausea by Risk of Bias Rating



Footnote: Three studies in the main analysis (Anton, 1999, Killeen, 2004, and Monti, 2001) reported nausea and/or vomiting as a single outcome. The risk difference in an analysis run without these three studies, the result was RD 0.10 (95% CI 0.06, 0.13), when the high risk of bias studies were also removed, the result was RD 0.11 (95% CI 0.07, 0.15).


Nausea – Naltrexone versus placebo – Sensitivity Analysis I

Naltrexone versus Placebo Nausea by Nausea alone or Nausea and/or Vomiting




Nausea – Naltrexone versus placebo – Sensitivity Analysis II (no high risk of bias studies)

Naltrexone versus Placebo Nausea by Nausea alone or Nausea and/or Vomiting




Nausea – Acamprosate versus naltrexone

Acamprosate versus Naltrexone Nausea by Risk of Bias Rating




Numbness – Acamprosate versus placebo

Acamp versus Placebo Numbness by Risk of Bias Rating




Numbness – Naltrexone versus placebo


Naltrex versus Placebo Numbness by Risk of Bias Rating




Rash – Acamprosate versus placebo



Rash – Naltrexone versus placebo




Suicide attempts/Suicidal ideation – Acamprosate versus placebo




Blurred vision – Naltrexone versus placebo

Naltrex versus Placebo Vision by Risk of Bias Rating




Vomiting – Acamprosate versus placebo




Vomiting – Naltrexone versus placebo

Naltrexone versus Placebo Vomiting by Risk of Bias Rating




Withdrawals due to Any Adverse Event – Acamprosate versus placebo

Acamprosate versus Placebo Withdrawal Due to Any Adverse Event by Risk of Bias Rating




Withdrawals due to Any Adverse Event – Naltrexone versus placebo




Withdrawals due to Any Adverse Event – Acamprosate versus naltrexone


Acamprosate versus Naltrexone Withdrawal Due to Any Adverse Event by Risk of Bias Rating



KQ 6 Analyses – NTX response by genotype, AA versus AG, GG




NTX response by genotype: Sensitivity Analysis I




Note: This SA is with imputing bad outcome (return to any drinking) for those dropped from Kim results that they reported.
Main analysis used Kim data reported by the article for the 32/63 who were adherent to NTX for 12 weeks; SA run with importing bad outcome for those lost (9 more for AA group and 22 more for the G carrier group)

NTX response by genotype: Sensitivity Analysis II




Note: Added Oslin 2003 (which did not meet inclusion criteria): Oslin 2003: Despite the main effect of genotype in the naltrexone-treated group, there was no medication by genotype interaction on relapse rates (OR 2.27 (95% CI: 0.44, 11.60, P 0.326). There was also no medication by genotype interaction for abstinence OR 0.89 (95% CI: 0.18, 4.38), P 0.889). Of note, there was a significant effect of naltrexone in reducing rates of relapse in the overall pooled sample even when genotype was included in the regression analysis (OR 2.42 (95% CI: 1.09, 5.39), p 0.030)

NTX response by genotype: Return to Heavy Drinking by RoB




NTX response by genotype: Return to Heavy Drinking: SA I



Note: This sensitivity analysis done with imputing bad outcome for Kim, 2009 missing data

NTX response by genotype: Return to Heavy Drinking: SA II



Note: Added Oslin, 2003 (which did not meet inclusion criteria)

Appendix G. Additional Studies of Genetic Polymorphisms Meeting Inclusion Criteria, but with Only 1 Study for a Drug-Polymorphism Pair

Characteristics of Trials

Table G-1 summarizes characteristics of the six included studies. Five were secondary or subgroup analyses from randomized controlled trials, and one was a prospective cohort study of patients taking disulfiram.¹ One of the trials compared naltrexone 50 mg/day with topiramate 50 to 400 mg/day;² one was a three-arm study that compared acamprosate 1,998 mg/day, naltrexone 50 mg/day and placebo;³ the others were placebo-controlled trials of nalmefene 20 mg/day,⁴ olanzapine 5 mg/day,⁵ and sertraline 200 mg/day.⁶ Duration of treatment ranged from 12 to 28 weeks; one trial also reported three- and six-month off-treatment follow-up data.⁶ Two were conducted in the U.S.,^{5,6} two in Germany,^{1,3} and one each in Finland⁴ and Spain.²

Mean age was very similar across studies, in the 40s. All patients met criteria for alcohol dependence. Enrollment of women and non-White subjects, when reported, was generally low. None of the studies reported information on smoking history at baseline. Two studies reported co-occurring psychiatric conditions: in one, 23 percent had a personality disorder,² and in the other, 26 percent had a concurrent drug use disorder.⁶ Co-interventions in the studies included BRENDA, combined behavioral interventions, and coping skills therapy. One study was rated medium risk of bias;⁶ the other five were rated high risk of bias.

In the three-arm study that compared acamprosate, naltrexone and placebo,³ the rs13273672 polymorphism in the *GATA4* gene was associated with relapse. At the end of 90 days of treatment, fewer patients with AA genotype relapsed than patients with AG or GG (45.7 percent versus 53.9 percent versus 69.0 percent, $p=0.0066$). The polymorphism was associated with relapse for patients treated with acamprosate, but not for those who received naltrexone or placebo.

In the trial that compared sertraline with placebo, secondary analyses examined the main and interaction effects with time of 3 factors—medication group, age of onset of alcohol dependence, and 5-HTTLPR genotype.⁶ The study reported differential effects for S' carriers and for L' homozygotes, with no significant effects in S' carriers. Late-onset (>25 years of age) alcoholics with the L'L' genotype who received sertraline had fewer drinking days at 12 weeks than those who received placebo ($p=0.007$), but there was no treatment difference in heavy drinking days. Early-onset L'L' individuals who received sertraline had more drinking days and more heavy drinking days ($p=0.002$ and $p=0.004$, respectively) at 12 weeks than those who received placebo. At three months off-treatment, late onset L'L' patients who had received sertraline continued to have fewer drinking days compared with placebo-treated patients ($p=0.027$).

The prospective cohort study of disulfiram revealed no significant gene-treatment interaction for time to relapse or cumulative abstinence between genotype groups based on the SNP rs1611115 of the *DBH* gene.¹

Table G1. Characteristics of included studies that assessed the association between genetic polymorphisms and medication response

Author, Year Design	Arm Dose, mg/day (N)	Genotypes Assessed	Medication Duration (F-u)	Setting	Age Years	Per-cent-age Non-White	Per-cent-age Fe-male	Cointervention(s)	Risk of Bias
Arias, 2008 ⁴ SSGA	Nalmefene 20 (166) Placebo (106)	<i>OPRM</i> <i>OPRD</i> <i>OPRK</i>	28	Finland; Outpatient 15 sites	49 to 50	0	20	BRENDA 100%	High
Florez, 2008 ² SSGA	Topiramate 50-400 (45) Naltrexone 50 (45)	<i>DRD2</i> <i>DRD3</i> <i>HTR2A</i> <i>SLC6A</i>	26	Spain; Outpatient	46	0	13	NR	High
Hutchison, 2006 ⁵ SSGA	Olanzapine 5 (33) Placebo (31)	<i>DRD4</i>	12	U.S.; Outpatient clinical research center	43 to 45	4 to 33	26 to 42	Brief structured psychosocial intervention 100%	High
Kiefer, 2011 ³ SSGA	Acamprosate 1998 (147) Naltrexone 50 (148) Placebo (74)	<i>GATA4</i>	12	Germany; Unclear	45	NR	NR	Medical management (CBI) 100%	High
Kranzler, 2012 ⁶ SSGA	Sertraline 200 intended, mean dose 169 (63) Placebo (71)	<i>5-HTTLPR</i> ^a	12 (26)	U.S.; Outpatient; university health center	48	8	19	CS 100%	Med
Mutschler, 2012 ¹ Prosp. cohort	Disulfiram NR (62)	<i>DBH</i> ^b	12	Germany; SA 48 treatment, Outpatient	48	NR	32	CBI 100%	High

^a5-HTTLPR is a polymorphism in the serotonin transporter gene. Variation at this locus includes higher-activity long (L) and lower-activity short (S) alleles.

^b SNP tested was rs1611115, located in the promoter region of the DBH gene.

Notes: Age (y) is the mean age in years, unless otherwise stated

Abbreviations: CBI, combined behavioral intervention; CS, coping skills; DBH, dopamine beta-hydroxylase; DR, dopamine receptor; follow-up in weeks; HTR2A, serotonin 2A receptor; mg, milligrams; N = Number; NR, not reported; OPRD, δ -opioid receptor; OPRK, κ -opioid receptor; OPRM, μ -opioid receptor; prosp., prospective; SLC6A, dopamine transporter; SSGA, secondary or subgroup analysis of a randomized controlled trial; U.S., United States

References for Appendix G

1. Mutschler J, Abbruzzese E, Witt SH, et al. Functional polymorphism of the dopamine β -hydroxylase gene is associated with increased risk of disulfiram-induced adverse effects in alcohol-dependent patients. *J Clin Psychopharmacol*. 2012;32(4):578-80. PMID: 22760354.
2. Florez G, Saiz P, Garcia-Portilla P, et al. Association between the Stin2 VNTR polymorphism of the serotonin transporter gene and treatment outcome in alcohol-dependent patients. *Alcohol Alcohol*. 2008 Sep-Oct;43(5):516-22. PMID: 18552399.
3. Kiefer F, Witt SH, Frank J, et al. Involvement of the atrial natriuretic peptide transcription factor GATA4 in alcohol dependence, relapse risk and treatment response to acamprosate. *Pharmacogenomics J*. 2011 Oct;11(5):368-74. PMID: 20585342.
4. Arias AJ, Armeli S, Gelernter J, et al. Effects of opioid receptor gene variation on targeted nalmefene treatment in heavy drinkers. *Alcohol Clin Exp Res*. 2008 Jul;32(7):1159-66. PMID: 18537939.
5. Hutchison KE, Ray L, Sandman E, et al. The effect of olanzapine on craving and alcohol consumption. *Neuropsychopharmacology*. 2006 Jun;31(6):1310-7. PMID: 16237394.
6. Kranzler HR, Armeli S, Tennen H. Post-treatment outcomes in a double-blind, randomized trial of sertraline for alcohol dependence. *Alcohol Clin Exp Res*. 2012 Apr;36(4):739-44. PMID: 21981418.